


**Metal Elements  
in Environment,  
Medicine and Biology**

**Tome VIII**



**ROUMANIAN ACADEMY**  
BRANCH TIMIȘOARA AND BRANCH CLUJ-NAPOCA



**IOAN SILAGHI-DUMITRESCU, ZENO GÂRBAN, PETRU DRĂGAN**

*EDITORS OF SYMPOSIA SERIES*

# Metal Elements in Environment, Medicine and Biology **Tome VIII**

*EDITED IN COLLABORATION WITH*

**CORNELIU DAVIDESCU, GABRIELA GARBAN, IOSIF GERGEN,  
SIMONA DRĂGAN, NICOLAE VASZILCSIN, ADINA AVACOVICI**

*Proceedings of the 8th International Symposium on  
Metal Elements in Environment, Medicine and Biology  
Timișoara, December 5-6, 2008*

Publishing House "Eurobit" Timișoara  
2008

**IOAN SILAGHI-DUMITRESCU, ZENO GÂRBAN, PETRU DRĂGAN**  
*EDITORS OF SYMPOSIA SERIES*

# **METAL ELEMENTS IN ENVIRONMENT, MEDICINE AND BIOLOGY**

**Tome VIII**  
**Second printing**

*Edited in collaboration with*  
**CORNELIU DAVIDESCU, GABRIELA GÂRBAN, IOSIF GERGEN, SIMONA DRĂGAN,  
NICOLAE VASZILCSIN, ADINA AVACOVICI**

Copyright (C) 2008, Publishing House "Eurobit" Timișoara, Roumania  
All Rights Reserved

*Technical Committee:*  
*Adina-Elena Avacovici*  
*Ariana-Bianca Velciov*  
*George-Daniel Ghibu*

**METAL ELEMENTS IN ENVIRONMENT, MEDICINE AND BIOLOGY 2008**  
**Tome VIII (Eds. Symp. Series I. Silaghi-Dumitrescu, Z. Gârban, P. Drăgan)**

*Proceedings of the 8<sup>th</sup> International Symposium on Metal Elements In Environment, Medicine and  
Biology, Timișoara, December 5-6, 2008*  
Publishing House "Eurobit" Timișoara, Roumania, 2008

ISSN 1583 – 4204

Ioan, Silaghi-Dumitrescu; Zeno, Gârban; Petru, Drăgan (Eds. Symp. Series)  
Corneliu Davidescu, Gabriela Gârban, Iosif Gergen, Simona Drăgan, Nicolae Vaszilcsin,  
Adina Avacovici (Eds. Tome VIII)

**ISSN 1583 – 4204**



# CONTENTS

|                                                                                                                                                                                                                                       |    |
|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|----|
| Foreward .....                                                                                                                                                                                                                        | 1  |
| Preface.....                                                                                                                                                                                                                          | 3  |
| 1. <b>Haiduc Ionel, Silaghi-Dumitrescu Ioan, Garban, Zeno Fischer-Fodor Eva, Silaghi-Dumitrescu Radu</b> (Cluj - Napoca - Roumania) - Metallomics .....                                                                               | 5  |
| 2. <b>Djujic Ivana</b> (Beograd - Serbia) – Effects of foliar selenium application on performances in plants/animals/man .....                                                                                                        | 15 |
| 3. <b>Nechifor Mihai</b> (Iasi - Roumania) – Normal and pathological involvement of bivalent cations in oro-maxilo-facial area .....                                                                                                  | 27 |
| 4. <b>Farkas Etelka, Csapó Edit, Buglyó Péter</b> (Debrecen – Hungary) - Potentially metalloenzyme inhibitor hydroxamic acids: effects of an imidazole donor on the metal binding capability .....                                    | 35 |
| 5. <b>Lăcătușu Radu, Iordache Virgil</b> (București - Roumania) - Integrated modeling of geochemical and biogeochemical processes across space-time scale .....                                                                       | 41 |
| 6. <b>Takács-Hajos Maria, Kastori Rudolf, Pucarevic Mira, Skoric Tijana Zeremski</b> (Szarvas - Hungary) - Mineral element content of leaves of table beet varieties and their role in nutrition .....                                | 55 |
| 7. <b>Avacovici Adina-Elena, Garban Zeno</b> (Timișoara - Roumania) - Characteristics of the metallogram in cystine urolithiasis – investigation means .....                                                                          | 61 |
| 8. <b>Papp Andras, Sárközi Leila</b> (Szeged - Hungary) - Consequences of subacute intratracheal exposure of rats to cadmium oxide nanoparticles: behavioral, electrophysiological and toxicological effects ..                       | 67 |
| 9. <b>Neagoe Aurora, Iordache Virgil, Bodescu Florin</b> (București - Roumania) - Bioaccumulation of metals in food chain of wetlands .....                                                                                           | 73 |
| 10. <b>Halasi Tibor, Lalosevic Dusan, Culic Milka, Sekulic Slobodan, Capo Ivan, Grbic Gordana</b> (Novi Sad - Serbia) - Aluminium neurotoxicity .....                                                                                 | 81 |
| 11. <b>Simescu Mihaela, De Benoist Bruno, Sava Mariana, Podia-Igna Claudia, Delange Francois</b> (București – Romania) - Correction of iodine deficiency by an iodized vegetable oil – floriol iodine product (partial results) ..... | 87 |
| 12. <b>Lalic Tatjana, Djujic Ivana</b> (Beograd - Serbia) - Mineral imbalance - reflection on blood vessels .....                                                                                                                     | 95 |

|                                                                                                                                                                                                                                                                                      |     |
|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----|
| 13. <b>Alexandrova Radostina, Timcheva Konstanta, Kalfin Reni, Mosoarca Elena-Maria, Tudose Ramona, Costisor Otilia, Culita C. Daniela, Marinescu Gabriela, Patron Luminita, Kantouri Maria-Lallia</b> (Sofia –Bulgaria) - Few words about metal compounds in cancer treatment ..... | 105 |
| 14. <b>Borcean Ioan, David Gheorghe, Niță Simona, Borcean Adrian, Băghină Narcis, Niță Lucian</b> (Timișoara - Roumania) - A study on the possible prevention of environmental pollution with mining wastes from copper mines through agricultural valorisation .....                | 113 |
| 15. <b>Contrea Aurel</b> (Timișoara - Roumania) - Study of the complexes formed between some metal ions with the condensed phosphates by means of gel-chromatography.....                                                                                                            | 117 |
| 16. <b>Creanga Adina, Henin Nathalie, Simescu Mihaela, De Benoist Bruno</b> (București - Roumania) - Refined sunflower oil : an excellent vehicle for iodine fortification .....                                                                                                     | 123 |
| 17. <b>Davidescu Corneliu-Mircea, Popa Adriana, Negrea Petru, Bandur Geza, Ilia Gheorghe, Ardelean Radu</b> (Timișoara - Roumania) - Phosphonate ligands grafted on styrene-divinylbenzene copolymers used in metal ions separation from water .....                                 | 127 |
| 18. <b>Drăgan Simona, Socaciu Carmen, Gergen Iosif, Yogi Lavanya, Dehelean A. Cristina</b> (Timișoara - Roumania) - Development of functional foods from soy extracts enriched with magnesium in breast cancer .....                                                                 | 133 |
| 19. <b>Fischer-Fodor Eva, Miclasova Natalia, Virag Piroska, Tatomir Corina, Brie Ioana, Perde Maria, Cernea Valentin, Silaghi-Dumitrescu Luminița</b> (Cluj-Napoca - Roumania) – Biologic evaluation of platinum and palladium complexes .....                                       | 141 |
| 20. <b>Garban Gabriela</b> (Timișoara - Roumania) – Food supplements with chromium and selenium content on the Roumanian market .....                                                                                                                                                | 147 |
| 21. <b>Garban Zeno</b> (Timișoara - Roumania) – Adducts of deoxyribonucleic acid with metals ad limina of interdisciplinarity metallomics and proteomics.....                                                                                                                        | 153 |
| 22. <b>Ghibu George-Daniel, Garban Gabriela, Baltă Cornel, Bura Marian, Petruse Cristina, Horhoi Doina-Elena, Garban Zeno</b> (Timișoara - Roumania) - Hepatic metallograms in rabbits after sodium nitrate excess in drinking water .....                                           | 163 |
| 23. <b>Hărmănescu Monica, Popovici Daniel, Moisuc Alexandru, Gergen Iosif</b> (Timișoara - Roumania) - Sodium, potassium, calcium and magnesium contents – are indicators for geographycal origin of bee's honey from Banat county? .....                                            | 169 |

|                                                                                                                                                                                                                                                                                                                                                     |     |
|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----|
| 24. <b>Hădărugă G. Nicoleta, Hădărugă I. Daniel, Negrea Adina, Ciopec E. Mihaela, Negrea Petru</b> (Timișoara – Roumania) - Minor and trace elements in parsley ( <i>Petroselinum sativum</i> hoffm.) from the west side of Roumania.....                                                                                                           | 173 |
| 25. <b>Ianoș Gheorghe</b> (Timișoara - Roumania) - Conclusions concerning the source and the environment concentration with heavy metals on the etno-historical Banat.....                                                                                                                                                                          | 181 |
| 26. <b>Koczka Noémi, Pétersz Dóra, Stefanovits-Bányai Éva</b> (Gödöllő – Hungary) - Metal ion composition of stinging nettle ( <i>Urtica dioica</i> L.) leaves in view of the nutritive value.....                                                                                                                                                  | 189 |
| 27. <b>Lăzureanu Aurel, Moigrădean Diana, Măruțoiu Constantin, Gogoasă Ioan, Gergen Iosif</b> (Timișoara - Roumania) - Study regarding the heavy metals contents in some vegetables cultivated in the field in romanian West area .....                                                                                                             | 195 |
| 28. <b>Nagy Ádám, Stefanovits-Bányai Éva, Neményi András, Hegedűs Attila</b> (Budapest - Hungary) - Microelement content in fruits of fig genotypes grown in hungary and in commercially available dried figs .....                                                                                                                                 | 199 |
| 29. <b>Papadopol Victoria, Popa Elena, Albu Gabriela, Cîmpeanu Cristina, Florescu Nicoleta, Palamaru Iliana</b> (Iași – Roumania) - Body zinc level, eating habits and some nutritional parameters in an elderly group..                                                                                                                            | 205 |
| 30. <b>Slezák Katalin, Kiss Sandor, Némethy Uzuni Hanna, Orosz Ferenc, Stefanovits-Bányai Éva</b> (Budapest - Hungary) - Effect of foliar magnesium fertilization on leaf and fruit mineral compositions of greenhouse grown cucumbers .....                                                                                                        | 211 |
| 31. <b>Bogdan T. Alexandru, Acatincăi Stelian, Avacovici Adina-Elena, Garban Zeno</b> (Timișoara - Roumania) - Biodiversity and xenobiotics related to foods of animal origin .....                                                                                                                                                                 | 217 |
| 32. <b>Szilvássy Blanka, Papp Nóra, Szabó Zoltán, Nyéki József, Tordai Enikő, Pedryc Andrzej, Stefanovits-Bányai Éva, Hegedűs Attila</b> (Budapest - Hungary) - Nutrient element of stone fruit species and cultivars.....                                                                                                                          | 225 |
| 33. <b>Staicu Andreea-Cristina, Postolache Carmen, Munteanu Maria-Cristina, Costache Marieta, Dinischiotu Anca</b> (București - Roumania) - Intestine and liver histopathological changes induced in fed and starvated goldfish ( <i>Carassius auratus auratus</i> , <i>Cyprinidae</i> , <i>Pisces</i> ) during chronic exposure to manganese ..... | 231 |
| 34. <b>Takács Szabolcs, Bankó Sarolta, Papp András</b> (Szeged - Hungary) - Acute electrophysiological effects of a metal and a non-metal mitochondrial toxin .....                                                                                                                                                                                 | 239 |

|                                                                                                                                                                                                                                                                                                |     |
|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----|
| 35. <b>Taubert Lidia, Rădulescu Hortensia, Princz Ecaterina, Stefanovits-Banyai Eva, Kiss A. Sandor</b> (Timișoara - Roumania) – Impact of some mineral elements from industrial waste on wheat plants .....                                                                                   | 243 |
| 36. <b>Teodorescu Daniela, Munteanu Maria-Cristina, Staicu Andreea-Cristina, Meșter Elena-Lotus, Dinischiotu Anca</b> (București - Roumania) – LDH activity and histological changes in copper intoxication of <i>Carassius auratus gibelio</i> liver after short and long time exposure ..... | 249 |
| 37. <b>Varvari Lidia, Dorneanu Sorin Aurel, Popescu Cătălin Ionel</b> (Cluj Napoca – Roumania) - K <sup>+</sup> -selective electrode based on a calix[6]arenic compound .....                                                                                                                  | 255 |
| 38. <b>Velciov Ariana-Bianca, Velciov Petru-Robert, Garban Gabriela, Avacovici Adina-Elena, Ahmadi Mirela, Sayti Ludovic</b> (Timisoara – Roumania) - In vivo effects of cis-platinum on magnesium levels in blood, muscles and heart samples.....                                             | 259 |
| 39. <b>Voroniuc Otilia, Gheorgheș Manuela, Florescu Nicoleta, Cristea Anca, Casaux-Voroniuc Iulia</b> (Iași - Roumania) - Magnesium in drinking water of some public systems and wells in Iași district and the daily magnesium requirement .....                                              | 265 |
| 40. <b>Silaghi-Dumitrescu Radu, Deac Florina</b> (Cluj-Napoca - Roumania) - The redox reactivity of globins: the chicken and egg paradox.....                                                                                                                                                  | 271 |
| 41. <b>Crețu Ana-Maria, Aschie Mariana, Preda Gabriela, Roșoiu Natalia</b> (București - Roumania) - The significance of cytomorphometry in peritoneal effusions cytology practice.....                                                                                                         | 277 |
| 42. <b>Hadnadjev Mila, Hadnadjev Darka</b> (Novi Sad - Serbia) - Ambrosia mowing in Novi Sad and the number of children with allergic diseases registered in primary care .....                                                                                                                | 285 |
| Authors index.....                                                                                                                                                                                                                                                                             | 289 |
| Retrospectives.....                                                                                                                                                                                                                                                                            | 291 |
| In memoriam Prof. Petru Drăgan MD, PhD (1932-2007) .....                                                                                                                                                                                                                                       | 297 |

## **FOREWORD**

*The 8th Symposium of the series „Metal Elements in Environment, Medicine and Biology” (M.E.E.M.B.) addresses to a distinctive domain of bioinorganic chemistry and life sciences: metallomics. It takes place in Timișoara under the auspices of Branch Timișoara and Branch Cluj-Napoca and of the Roumanian Academy,*

*Approached topics regard the complex problems of metallomics in different domains for interest in environment (geobiochemistry, ecology, contamination of habitual area etc.), medicine (applications in clinical chemistry, patobiochemistry, implantology, cardiology, nephrology, etc.), nutrition (distribution in processed food, obtainment of food supplements, methodology of foods fortification), pharmacy (the synthesis and testing of chemiotherapeutic drugs with metal content, obtainment of some vegetal extracts), biology (distribution of metals in the vegetal and animal reign etc.*

*First symposium had as honorary president Acad. Nicolae Cajal, vicepresident of the Roumanian Academy (1990-1994), who agreed the series of symposia to take place under the auspices of Branch Timișoara of the Roumanian Academy. Beginning with the 2nd symposium (1996) until now (2008), the symposia were under the patronage of the Roumanian Academy – Commission of Biochemistry, having as honorary president Acad. Ionel Haiduc.*

*The series of symposia „Metal Elements in Environment, Medicine and Biology” has been initiated in 1993 at the proposal of the „Working Group on Metal Research in Biological Systems” constituted of researchers from universities and some scientific institutions from Timișoara (1979).*

*Starting with 2008 the series of symposia take place under the auspices of Branch Cluj-Napoca of the Roumanian Academy and of Branch Timișoara of the Roumanian Academy with the perspective to extent the internal and foreign collaborations and the thematic area.*

*The aprovement of the Roumanian Academy given by President Acad. Ionel Haiduc and General secretary Acad. Ion Păun Otiman for the cooperation of Branch Cluj-Napoca and Branch Timișoara, can offer a benefit for the progresses in this domain.*

*In context of contemporary development focused on environmental, nutritional and public health problems, preoccupations in the domain of metallomics present a special interest.*

*It is well known the fundamental and applicative character of metallomics in this domain. It confers explanation for the continuity and even broadening of topics of this symposium series.*

*President of Branch Timișoara of Roumanian Academy,*

*Prof. Zeno Simon, PhD  
corresponding member of Roumanian Academy*



## PREFACE

The on-going biennial series of Symposia „Metal Elements in Environment, Medicine and Biology” (M.E.E.M.B.) and the publishing of Tome I-VIII (1994-2008) confirm the existing interest for problems referring to the applications of Inorganic biochemistry, generally, and of Metallomics, especially, in fundamental and applicative domains.

The interest shown to the presence of metals in the living area and of their role in life sciences is proved by the continuous development of metallomics.

The Series of M.E.E.M.B. symposia – taking place during one and a half decade – under the auspices of Branch Timișoara of the Roumanian Academy benefitted of the support offered by the president Prof. Zeno Simon, PhD – corresponding member of Roumanian Academy (editions 2006 and 2008) and former presidents Acad. Ion Păun Otiman (editions 2002 and 2004) and Acad. Toma Dordea (editions 1996, 1998 and 2000) to whom we express our thanks.

At each edition was remarkable the contribution of scientists from diverse universities and research institutes from inland and abroad.

As it is well known, implication of metals and their derivatives in environment, nutrition, medicine a.o. are of main interest nowadays. Between metal ions and bioactive compounds can occur complex interactions which have a beneficial or harmful action on the functioning of the organism. Also, metal ions can, in case of pollution, represent a danger for public health.

While traditional bioinorganic chemistry is focused on the role and interactions of a single metal in a protein or enzyme system, «metallomics studies global, multi-element interactions and relationships» (Koppendaal and Hieftje, 2007). One can appreciate the significant connexions between chemistry and life sciences and identify easily the important concepts of today's knowledge.

Actual meeting was supported by Prof. Nicolae Robu, Eng, PhD – rector of University „Politehnica” Timișoara and Prof. Alexandru Moisuc, Eng. PhD – rector of the University of Agricultural Sciences and Veterinary Medicine of Banat Timișoara. In the M.E.E.M.B. Symposia Series organisation and perpetuation a special contribution had Prof. Gheorghe Ciuhandu, Eng, PhD – mayor of Timișoara city.

As the series of M.E.E.M.B. Tomes have ISSN number, it is confirmed its perpetuation and the possibility of extend to an annual meeting (until now having a biennial character). In order to pursue the future evolution of the Symposium it is desirable to encourage young colleagues to continue their scientific involvement in the domain of metallomics.

*Editors*





# METALLOMICS

Ionel HAIDUC<sup>1</sup>, Ioan SILAGHI-DUMITRESCU<sup>1\*</sup>, Zeno GARBAN<sup>2</sup>,  
Eva FISCHER-FODOR<sup>3</sup>, Radu SILAGHI-DUMITRESCU<sup>1</sup>

1. Department of Chemistry, Faculty of Chemistry and Chemical Engineering Babeş-Bolyai University, Str. Arany Janos Nr. 11, RO-400 028 Cluj-Napoca, Roumania; 2. Department of Biochemistry and Molecular Biology, Faculty of Food Products Technology, University of Agricultural Sciences and Veterinary Medicine of Banat, Calea Aradului Nr.119, RO-300 645, Timişoara, Roumania; 3. Research Department, I. Chiricuţă Cancer Institute, Str. Republicii Nr. 34-36, RO-400 015 Cluj-Napoca, Roumania

\* E-mail : isi@chem.ubbcluj.ro

The chemical derivatives of metals were tested in medicine, with contradictory results, even from the middle ages. The toxicity of many metal compounds especially at uncontrolled concentrations, has made the possible medical use of metals to be regarded with reluctance by clinical doctors and significantly successful. The situation changed with the discovery [1] of the antitumoral activity of Cisplatin  $[\text{Pt}(\text{NH}_3)_2\text{Cl}_2]$ , a very active compound against some forms of cancer. Cisplatin was clinically tested in 1972 [2], and in 1979 it was approved by U.S. Food and Drug Administration as a drug in the treatment of cancer [3]. In the years to come, several thousands platinum compounds were screened for antitumoral activity, and more efficient and less toxic derivatives were discovered, now known under several trade names, such as Iproplatin, Carboplatin, Oxaliplatin, Lobaplatin, Nedaplatin, etc. The extraordinary efficacy of platinum derivatives against various forms of cancer (especially testicular, ovarian, bladder, brain and throat tumors) awakened a renewed interest for the possible medical applications of metal combinations.

On the other hand, the understanding of the role of essential metals in biological process lead to a novel **interdisciplinary field**, the BIOINORGANIC CHEMISTRY [5], based mainly on **fundamental research**, but with applications constantly growing in importance. Lately, a new subject is emerging, studying the action of metals on living cells, their effect on the bio-control of cellular processes, through the influence on the metabolism, the cell biologic functions and biosynthesis and the medical implications occurring, called METALLOMICS [6], designation similar to that of new subjects known as genomics, proteomics, etc. METALLOMICS, a new journal from RSC Publishing, will be launched in January 2009. The announcement was made recently at the International Forum of Proteins meeting on Health, Food and Environment by Joseph A. Caruso, who will chair the Editorial Board.

The biological role of some metals (essential elements) is determined by complex interactions between metallic ions and bioactive organic compounds. The lack of essential metals (but also the excess) may cause specific illnesses. For instance, the iron as a component of hemoglobin plays an essential biological role, but the excess of iron in the organism leads to diseases. The copper is essential in the development of embryos, temperature control and the functioning of the nerve cell, but it is also involved in the Menkes disease. Chromium is essential in the carbohydrate and lipid metabolism, its deficiency can provoke diabetes, but the excess is toxic. Some metals are toxic in any concentrations (mercury, cadmium, lead, thallium) and their elimination from the organism represents a very important topic.

The implications of metals and their derivatives, including organometallic compounds, in medicine are numerous and varied [7, 8]. The following aspects can be mentioned: (a) toxicity and homeostasis; (b) therapeutic agents already incorporated in clinical practice (for instance the platinum derivatives displaying antitumoral activity, gold compounds with antiarthritis action, antibiotics with metal-mediated activity (bleomycin/iron, copper)); (c) radiopharmaceuticals for diagnosis and treatment (technetium 99, other radioactive isotopes); (d) contrast agents for Roentgen imagistic (barium sulphate) and magnetic resonance (gadolinium complexes); (e) antibacterial, antiviral, antiparasitic agents; (f) psychotherapeutic agents, antidepressants; (g) mineral food supplements; (h) redox mediators for biosensors with medical applications.

The recent interest in the biological role of metals and their implications in medicine is illustrated by several examples: a) research programs financed by grants in USA (National Institutes of Health) [9], Europe (COST programs) [10], Australia and other countries; b) conferences, congresses and theme-symposiums, like the prestigious Gordon Conferences in USA [11]; c) a large number of books published in the last years (from which a short selection is presented [12]).

As mentioned before, the metal-based drugs are intensely employed in the chemotherapeutical treatment of cancer and numerous aspects of their biological activities have already been studied. In clinical practice, the use of metal compounds is current, and in the anticancer therapy, platinum-based drugs have a very important impact on the disease remission and survival rate [13, 14]. A constant improving of these therapies is necessary though, because their clinical use is limited by unwanted secondary effects and the worldwide trend is to individualize the treatment for each patient. The therapeutic index of the cytotoxic drugs – the ratio between positive effects and their toxicity - is low, and its improvement is a major objective [15].

Next to platinum, other metallic and organometallic derivatives have been tested in an attempt to discover novel antimitotic compounds. A large number of publications showed the interdependency between the structure, biologic effect and biomedical applications of the novel metal compounds in cancer therapy. The study of cytotoxicity, that of the mutation-inducing capacity in the tumor cell and of the antimitotic effect leads to the identification of structure-effect dependency for numerous compounds. The incorporation of different metals in the structure of cell components constitutes the basis of their biological activity. Many aspects of their biological effects have been studied: apoptosis-inducement, genotoxicity, DNA lesions and the repair mechanisms, reactive oxygen species inducement, immunological implications, the molecular basis of the cell resistance and the synergy with irradiation effects. Several complexes of platinum and palladium that contain ligands with a high antitumoral activity [16] were identified and characterized.

Newer cytotoxic agents have a broader spectrum of anticancer activity, showing toxicity towards both platinum-sensitive and resistant lines. Organotin compounds were identified as novel antitumoral agents that act in the apoptosis of cervical and ovarian carcinomas in human cells [17, 18]. Vanadium [19] and ruthenium [20] also form complexes with selective antitumoral and antimetastatic effects, especially towards platinum-resistant tumors, and they display a low systemic toxicity in therapeutic levels. The activity [21] of rhenium and rhodium complexes is related to their structural and kinetic properties. Apoptosis is induced by the inhibition of cellular tyrosin-phosphatases and the cleavage and fragmentation of cellular DNA. Gallium [22] displays tumor-growth inhibition properties, through mechanisms competing with cellular iron-acquiring. The arsines, arsenic trioxide and

organoarsenic compounds are efficient against tumoral cells. Gold (III) is isoelectronic with Pt (II); its target is not the DNA, but the proteins, and the effects are protease-inhibition and apoptosis-inducement [23]. The efficiency of titan and molibdenum organometallic derivatives is correlated to the presence of cyclopentadienyl rings in their structure, and their biological activity is determined by the interaction with thiolic groups from the cell [24]. Lanthanum and gadolinium are also employed in cancer therapy as single agents or neoadjuvants in radiotherapy [25, 26].

The antitumoral effect of platin-based drugs in mammals is related to their binding to DNA components. A selective method of determining Cisplatin-DNA adducts is based on the use of element-selective mass spectrometry [27, 28]. The most important DNA adduct with Cisplatin is formed between adjacent guanine units of the same DNA strand (1, 2 intrastrand cross-links). The repair of the DNA lesions contribute to clinical acquired resistance and is an effect of the resistance to drugs.

Assays which measure DNA mutation frequency are appropriate for the assessment of genotoxicity [29, 30]. The single cell gel electrophoresis assay ("Comet assay") detects lesions and DNA repair, cellular mechanisms such as drug transport and detoxification and it is a method for measuring deoxyribonucleic acid strand breaks, and DNA-protein crosslink in eukaryotic cells. New platinum compounds that form more extensive lesions in DNA [31] have been identified. The preferred nucleotides for the intercalation of metal complexes in the double helix of DNA in cancer cells, which causes irreparable lesions in the macromolecule [32-34] have been evidenced. The interaction of metal complexes with other biologic molecules like proteins and peptides, through the terminal amine, the imidazolic group of histidine or the thiol group of methionine was reported [35].

Reactive oxygen species (ROS) play an important role in cell signaling, and when present at high concentrations, they can induce oxidative stress. Nitrogen oxides (NO<sub>x</sub>) react with oxygen radicals to generate highly damaging reactive nitrogen species that may inhibit DNA repair mechanisms [36]. This is recognized as playing an important role in the evolution of the cancer cell and the NO<sub>x</sub> generated by the metal-based drugs can influence the result of cancer treatment [37]. It has been proved that the addition of an NO-releasing moiety in chemotherapeutic drug molecule increases the anti-neoplastic activity of the drug, while it provides protection against apoptosis induced by oxidative stress and it reduces its cytotoxicity against normal, non-neoplastic cells by blocking the oxidative stress that would lead to their apoptosis [38-39]. Nitrogen monoxide-releasing agents currently used in biology and medicine are organic (NONO-ates [40-42], nitrosyl-glutathione) or inorganic (nitrite, sodium nitroprusside, or other coordination compounds with NO coordinated to transition metals [45, 46]); depending on their electronic structure, these donors can act as direct NO, NO<sup>+</sup> or NO<sup>-</sup> donors, with different potentials that are not yet fully explored as far as the biomedical effect is concerned [43, 44].

Standard drugs like Cisplatin alter the nitric oxide synthetase (NOS) levels through regulation mechanisms and increase p53 protein levels inducing the apoptosis of tumor cells. The matrix metalloproteinases (MMPs) have been identified as the targets of the treatment in various pathological conditions, such as cancer, and their overexpression has been associated with the high potential of metastasis in human carcinomas. In order to optimize the therapeutic efficacy of any drug, it is useful to be able to monitor the MMPs *in vitro* as well as *in vivo*. Platinum complexes able to inhibit MMPs through a noncompetitive mechanism have been reported [28]. These results, along with the investigations of their mechanisms (DNA interaction

and tumor cell growth inhibition), demonstrate that ligand modifications of platinum compounds can be exploited to target biological substrates other than DNA.

In this context, it is obvious that the understanding of the nitro-oxidative stress mechanism and the metabolism of nitrogen oxides/oxoanions is necessary – these aspects involve as primary targets the metalloenzymes [40-42, 48-53].

Metal-based antitumoral therapy has a significant impact on the immune system. Cell-mediated immunity parameters, such as CD4+, CD8+, and CD 28 which modulate T lymphocyte and the natural killer cells antitumoral response have been monitored in several studies [54, 55]. In the case of cervical cancer patients, neoadjuvant platinum-based therapy enhances the immune response against tumoral cells [56]. The key moment is the activation of the T lymphocyte through a signal from a co-stimulatory molecule, and the role of costimulatory receptors like CD28 and the glucocorticoid induced tumor necrosis factor (GITR-) has been established [57-59]. Platin-based antitumor drugs have radiosensitizing properties; they are considered as among the most active drugs in both neoadjuvant and salvage treatments for patients with cervical cancer [60].

The major impact of their association is a significant decrease in the loco-regional evolution, recurrence rates (50%) and metastases rates. Risk of death was decreased with 40% [61, 62]. Randomized clinical trials evidenced the superiority of platin-based radiochemotherapy versus radiation alone; the optimal chemotherapeutic regimens remain to be defined [63].

Metallomics is of interest for domains that are connex to inorganic biochemistry regarding lithiasic formations (renal, hepatic, sialic, thyroidic ones), problems of implantology (in dentistry, ortopedics), colloid-osmotic balances, dishomeostasis of hydroelectrolytic metabolism (with impact in nutrition, cardiology, nephrology etc.), biogenesis of DNA-biometal adducts (genetics, biotechnology), DNA-chemiotherapeutic drugs adducts (e.g. cisplatin) and DNA-metals with toxicogen potential adducts (xenobiochemistry, e.g. Pb, Cd, Sn, etc.) [64-66]

Our interest and contribution in this field is illustrated by the two-volume treatise on the applications of organometallic chemistry in cancer chemotherapy [8], a number of articles relevant for the proposed topic from which some of them are presented [67-69].

## REFERENCES

- [1] B. Rosenberg, L. Van Camp, T. Krigas, Inhibition of cell division in *Escherichia coli* by electrolysis products from a platinum electrode [17], **Nature** **1965**, 205, 698; B. Rosenberg, L. Van Camp, J.E. Trosko, V.H. Mansoui, Platinum compounds: A new class of potent antitumour agents [24], **Nature** **1969**, 222, 385.
- [2] D.J. Highby, J.H. Wallace, Jr., J.F. Hollaand, Cis-diamminedichloroplatinum (NSC 119875): A phase 1 study, **Cancer Chemother. Repts.** **1973**, 57, 459.
- [3] B. Lippert (Editor), **Cisplatin. Chemistry and biochemistry of a leading anticancer drug**, Wiley-VCH, Weinheim, **2000**.
- [4] Z. Guo, P.J. Sadler, Metals in medicine, **Angew. Chem., Int. Ed.** **1999**, 38, 1512.
- [5] J.J.R. Fausto da Silva, R.J.P. Williams, **The biological chemistry of elements: The inorganic chemistry of life**, Clarendon Press, New York, **1991**.
- [6] (a) D.W. Koppenaal, G.M. Hieftje, Metallomics- an interdisciplinary and evolving bfield, **J. Analyt. Atomic Spectr.** **2007**, 22, 855; (b) H. Haraguchi, Metallomics as integrated biometal science, **J. Analyt. Atomic Spectr.** **2004**, 19, 5.

<sup>[7]</sup> R. M. Roat-Malone, **Bioinorganic chemistry: A short course**, John Wiley & Sons, Inc., New York, **2002**.

<sup>[8]</sup> I. Haiduc, C. Silvestru, **Organometallics in Cancer Chemotherapy. Volume 1. Main Group Metal Compounds**, CRC Press, Inc., Boca Raton, Florida, 1989 (254 pag.); **Volume 2. Transition Metal Compounds**, CRC Press, Inc., Boca Raton, Florida, 1990 (366 pag.).

<sup>[9]</sup> **Metals in Medicine**, PA-08-2008- (Department of Health and Human Services, National Institutes of Health (NIH): National Institute of General Medical Sciences, National Institute Environmental Health Sciences, Office of Dietary Supplements); continua programele **Metals in Medicine** PA-07-276, PA-05-001, si PA-01-071 (2001-2004).

<sup>[10]</sup> Action D8: **Metals in Medicine**, 1996-2001; Action D18: **Lanthanide Chemistry for Diagnosis and Therapy**, 1999-2006; Action D20: **Metal Compounds in the Treatment of Cancer and Viral Diseases**, 2000-2005; Action 21: **Metalloenzymes and Chemical Biomimetics**, 2000-2005.

<sup>[11]</sup> **Applications of Metals in Medicine and Healthcare**, 3-5 September 2007, Durham University, UK; **Metals in Medicine**, Gordon Research Conference, June 13-18, 2004, Colby College, Waterville, ME, USA; **Metals in Medicine**, Gordon Research Conference, June 29-July 4, 2008, Proctor Academy, Andover, NH, USA; **Metals in medicine. Targets, diagnostics, therapeutics**, 28-29 June 2000, National Institutes of Health, Bethesda, MD, USA; **Metals in Medicine**, University of Southern California, April 15, 2004; **International Workshop on Metallothioneins**, European Commission, Geel, Belgium, October 30-31, 1996.

<sup>[12]</sup> (a) M. Gielen, E.R.T. Tiekink (Editors), **Metallotherapeutic Drugs and Metal-based Diagnostic Agents**, John Wiley & Sons Inc., New York, **2005**; (b) A. Sigel, H. Sigel: **Metal ions and their complexes in medication**, New York : Marcel Dekker, **2004**; (c) R. Crichton, **Inorganic Biochemistry of Iron Metabolism: from Molecular Mechanisms to Clinical Consequences**, John Wiley & Sons, New York, **2001**; (d) L.D. Zardiackas, M.J. Kraay, H.L. Freese, **Titanium, niobium, zirconium, and tantalum for medical and surgical applications**, ASTM Pub., West Conshohocken, PA, 2006; (e) J.A. Helsen; H.J. Breme, **Metals as biomaterials**, John Wiley & Sons, Chichester, New York, **1998**; (f) **Advances in metals in medicine**. JAI Press, Greenwich, Conn., **1993**; (g) R.R. Brooks, **Noble Metals and Biological Systems: Their Role in Medicine, Mineral Exploration, and the Environment**, CRC Press, **1992**; (h) G. William K Beatty: **The precious metals of medicine**, Scribner Publishers, New York, **1975**.

<sup>[13]</sup> B.A. Teicher, Newer cytotoxic agents: attacking cancer broadly, **Clin. Cancer Res.** **2008**, 14, 1610-7.

<sup>[14]</sup> M.A. Jakupiec, M. Galanski, V.B. Arion, C.G. Hartinger, B.K. Keppler, Antitumour metal compounds: more than theme and variations, **Dalton Trans.**, **2008**, 83-94.

<sup>[15]</sup> A.G. Bosanquet, P.B. Bell, Ex vivo therapeutic index by drug sensitivity assay using fresh human normal and tumor cells, **J. Exp. Ther. Oncol.** **2004**, 4, 145-54.

<sup>[16]</sup> A.C.Caires, Recent advances involving palladium (II) complexes for the cancer therapy, **Anticancer Agents Med. Chem.**, **2007**, 7(5), 484-91.

<sup>[17]</sup> N. Höti, J. Ma, S. Tabassum, Y. Wang, M. Wu, Triphenyltin benzimidazolethiol, a novel antitumor agent, induces mitochondrial-mediated apoptosis in human cervical cancer cells via suppression of HPV-18 encoded E6, **J. Biochem.** **2003**, 134, 521-8.

<sup>[18]</sup> F. Barbieri, F. Sparatore, M. Cagnoli, C. Bruzzo, F. Novelli, A. Alama, Antiproliferative activity and interactions with cell-cycle related proteins of the

organotin compound triethyltin(IV)lupinylsulfide hydrochloride, **Chem. Biol. Interact.**, **2001**, 134, 27-39.

<sup>[19]</sup> A.M. Evangelou, Vanadium in cancer treatment, **Crit. Rev. Oncol. Hematol.**, **2002**, 42, 249-65.

<sup>[20]</sup> I. Kostova, Ruthenium complexes as anticancer agents, **Curr. Med. Chem.**, **2006**, 13, 1085-1107.

<sup>[21]</sup> P.C. Bruijninx, P.J. Sadler, New trends for metal complexes with anticancer activity, **Curr. Opin. Chem. Biol.**, **2008**, 12, 197-206.

<sup>[22]</sup> M.A. Jakupec, B.K. Keppler, Gallium and other main group metal compounds as antitumor agents, **Met. Ions Biol. Syst.**, **2004**, 42, 425-62.

<sup>[23]</sup> V. Milacic, D. Fregona, Q.P. Dou, Gold complexes as prospective metal-based anticancer drugs, **Histol. Histopathol.**, **2008**, 23, 101-8.

<sup>[24]</sup> P.M. Abeysinghe, M.M. Harding, Antitumour bis(cyclopentadienyl) metal complexes: titanocene and molybdocene dichloride and derivatives, **Dalton Trans.**, **2007**, 3474-82. <sup>[25]</sup> G.M. Richards, M.P. Mehta, Motexafin gadolinium in the treatment of brain metastases, **Expert Opin. Pharmacother.**, **2007**, 8, 351-9.

<sup>[26]</sup> A.J. Freemont, Lanthanum carbonate, **Drugs Today**, **2006**, 42, 759-70.

<sup>[27]</sup> A.R. Collins, The comet assay for DNA damage and repair. Principles, applications, limitations, **Mol. Biotechnol.**, **2004**, 26, 249-61.

<sup>[28]</sup> O. Merk, G. Speit, Detection of crosslinking with the comet assay in relationship to genotoxicity and cytotoxicity, **Environ. Mutat. Res**, **1999**, 33, 167-72.

<sup>[29]</sup> D. Garcia Sar, M. Montes- Bayon, L. Aguado Ortiz, E. Blanco- Gonzalez, L.M. Sierra, A. Sanz-Medel, In vivo detection of DNA adducts induced by cisplatin using capillary HPLC-ICP-MS and their correlation with genotoxic damage in *Drosophila Melanogaster*, **Anal. Bioanal. Chem**, **2008**, 390, 37-44.

<sup>[30]</sup> P. Wynne, C. Newton, J.A. Ledermann, A. Olaitan, T.A. Mould, J.A. Hartley, Enhanced repair of DNA intersatrand crosslinking in ovarian cancer cells from patients following treatment with platinum-based chemotherapy, **Brit. J. Cancer**, **2007**, 97, 927-33.

<sup>[31]</sup> J. Blasiak, J. Kowalik, E. Malecka Panas, J. Drzewoski, M. Wojewodska, DNA damage and repair in human lymphocytes exposed to three anticancer platinum drugs. **Teratog. Mutat. Res**. **2000**, 20(3), 119-31.

<sup>[32]</sup> J. Reedijk, New clues for platinum antitumor chemistry: kinetically controlled metal binding to DNA, **Proc. Natl. Acad. Sci. U.S.A.**, **2003**, 100, 3611-6.

<sup>[33]</sup> H.T. Chifotides, K.R. Dunbar, Interactions of metal-metal-bonded antitumor active complexes with DNA fragments and DNA, **Acc. Chem. Res.**, **2005**, 38, 146-56.

<sup>[34]</sup> A. Halamikova, P. Heringova, J. Kasparova, F.P. Intini, G. Natile, A. Nemirovski, D. Gibson, V. Brabec, Cytotoxicity, mutagenicity, cellular uptake, DNA and glutathione interactions of lipophilic trans-platinum complexes tethered to 1-adamantine, **J. Inorg. Biochem.**, **2008**, 102, 1077-89.

<sup>[35]</sup> R. Ranaldo, N. Margiotta, F.P. Intini, C. Pacifico, G. Natile, Conformer distribution in (cis-1,4-Dach) bis(guanosine-5-phosphate)platinum II adducts: A reliable model for DNA adducts of antitumoral cisplatin, **Inorg. Chem.**, **2008**, 47, 2820-30.

<sup>[36]</sup> E.L. Leung, M. Fraser, R.R. Fiscus, B.K. Tsang, Cisplatin alters nitric oxide synthase levels in human ovarian cancer cells: involvement in p53 regulation and cisplatin resistance, **Brit. J. Cancer**, **2008**, 98(11), 1803-9.

<sup>[37]</sup> K.A. Conklin, Cancer chemotherapy and antioxidants, **J Nutr**. **2004**, 134(11), 3201-4. <sup>[38]</sup> L. Santucci, A. Mencarelli & al., Nitric oxide modulates proapoptotic and

antiapoptotic properties of chemotherapy agents, **FASEB Journal**, **2006**, 20, 765-767.

<sup>[39]</sup> D.G. Hirst, T. Robson, Nitrosative stress in cancer therapy, **Front. Biosci.**, **2007**, 12, 3406-18.

<sup>[40]</sup> R. Silaghi-Dumitrescu, D. M. Kurtz, Jr., L. G. Ljungdahl, W. N. Lanzilotta, X-ray Crystal Structures of *Moorella thermoacetica* FprA. Novel Diiron Site Structure and Mechanistic Insights into a Scavenging Nitric Oxide Reductase, **Biochemistry**, **2005**, 44, 6492-501.

<sup>[41]</sup> R. Silaghi-Dumitrescu, K. Y. Ng, R. Viswanathan, D. M. Kurtz, Jr., A Flavodiiron protein from *Desulfovibrio vulgaris* with oxidase and nitric oxide reductase activities. Evidence for an *In vivo* nitric oxide scavenging function, **Biochemistry**, **2005**, 44, 3572-9.

<sup>[42]</sup> R. Silaghi-Dumitrescu, E. D. Coulter, A. Das, L. G. Ljungdahl, G. N. Jameson, B. H. Huynh, D. M. J. Kurtz, A flavodiiron protein and high molecular weight rubredoxin from *Moorella thermoacetica* with nitric oxide reductase activity, **Biochemistry**, **2003**, 42, 2806-2815.

<sup>[43]</sup> K. Karidi, A. Garoufis, A. Tsipis, N. Hadjiliadis, H. den Dulk, Reedijk, Synthesis, characterization, in vitro antitumor activity, DNA-binding properties and electronic structure (DFT) of the new complex cis-(Cl,Cl)[RuIICl<sub>2</sub>(NO<sup>+</sup>)(terpy)]Cl, **Dalton Trans.**, **2005**, (7), 1176-1187.

<sup>[44]</sup> K. Karidi, A. Garoufis, A. Tsipis, N. Hadjiliadis, H. den Dulk; J. Reedijk, Synthesis, characterization, in vitro antitumor activity, DNA-binding properties and electronic structure (DFT) of the new complex cis-(Cl,Cl)[RuIICl<sub>2</sub>(NO<sup>+</sup>)(terpy)]Cl, **Dalton Trans.**, **2005**, 1176-87.

<sup>[45]</sup> C. E. Cooper, Nitric oxide and cytochrome oxidase: Substrate, inhibitor or effector, **Trends Biochem. Sci**, **2002**, 27, 33-9.

<sup>[46]</sup> R. Andrew Colin, R. Kenton Rodgers, R.A. Eady Robert, Novel Kinetic Trap for NO Release from Cytochrome c': A Possible Mechanism for NO Release from Activated Soluble Guanylate Cyclase, **J. Am. Chem. Soc.**, **2003**, 125, 9548-9549.

<sup>[47]</sup> R. Sasanelli, A. Boccarelli, D. Giordano, M. Laforgia, F. Arnesano, G. Natile, C. Cardellicchio, M.A. Capozzi, M. Coluccia, Platinum complexes can inhibit matrix metalloproteinase activity: platinum-diethyl[(methylsulfinyl)methyl]phosphonate complexes as inhibitors of matrix metalloproteinases 2, 3, 9, and 12, **J. Med. Chem.**, **2007**, 50:3434-41.

<sup>[48]</sup> R. Silaghi-Dumitrescu, I. Silaghi-Dumitrescu, DFT and the electromerism in complexes of iron with diatomic ligands, **J. Inorg. Biochem.**, **2006**, 100, 161-6.

<sup>[49]</sup> R. Silaghi-Dumitrescu, Copper-containing nitrite reductase: a DFT study of nitrite and nitric oxide adducts, **J. Inorg. Biochem.**, **2006**, 100, 396-402.

<sup>[50]</sup> R. Silaghi-Dumitrescu, A density functional study of heme-peroxynitrite adducts, **J. Mol. Struct. THEOCHEM**, **2005**, 722, 233-237.

<sup>[51]</sup> R. Silaghi-Dumitrescu, Nitrile hydration by the cobalt-containing nitrile hydratase DFT investigation of the mechanism, **Rev. Chim.**, **2004**, 55, 496-498.

<sup>[52]</sup> R. Silaghi-Dumitrescu, Linkage isomerism in nitrite reduction by cytochrome cd<sub>1</sub> nitrite reductase, **Inorg. Chem.**, **2004**, 43, 3715-3718.

<sup>[53]</sup> R. Silaghi-Dumitrescu, Nitric oxide reduction by heme-thiolate enzymes (P450<sub>nor</sub>): A reevaluation of the mechanism **Eur. J. Inorg. Chem.**, **2003**, 1048-1052.

<sup>[54]</sup> S. Coleman, A. Clayton, M.D. Mason, B. Jasani, M. Adams, Z. Tabi, Recovery of CD8<sup>+</sup> T-cell function during systemic chemotherapy in advanced ovarian cancer, **Cancer Res.**, **2005**, 65, 7000-6.

<sup>[55]</sup> C.W. Tseng, C.F. Hung, R.D. Alvarez, C. Trimble, W.K. Huh, D. Kim, C.M.

Chuang, C.T. Lin, Y.C. Tsai, L. He, A. Monie, T.C. Wu, Pretreatment with Cisplatin Enhances E7-Specific CD8+ T-Cell-Mediated Antitumor Immunity Induced by DNA Vaccination, **Clin. Cancer Res.** **2008**, 4, 3185-92.

<sup>[56]</sup> H.R. Cosiski Marana, J. Santana da Silva, J. Moreira de Andrade, NK cell activity in the presence of IL-12 is a prognostic assay to neoadjuvant chemotherapy in cervical cancer, **Gynecol. Oncol.** **2000**, 78, 318-23.

<sup>[57]</sup> K. M. Baltz, M. Krusch, A. Bringmann, P. Brossart, F. Mayer, M. Kloss, T. Baessler, I. Kumbier, A. Peterfi, S. Kupka, S. Kroeber, D. Menzel, M. P. Radsak, H. G.Rammensee, H. R. Salih, Cancer immunoediting by GITR (glucocorticoid-induced TNF-related protein) ligand in humans: NK cell/tumor cell interactions, **FASEB Journal**, **2007**, 21, 2442-2454.

<sup>[58]</sup> L. H. Boise, A. J. Minn, P. J. Noel, C. H. June, M. A. Accavitti, T. Lindsten, C. B. Thompson, , 1995. CD28 costimulation can promote T cell survival by enhancing the expression of Bcl-XL. **Immunity** 3, 87-98.

<sup>[59]</sup> G. Frumento, T. Piazza, E. Di Carlo, S. Ferrini, Targeting tumor-related immunosuppression for cancer immunotherapy. **Endocr. Metab. Immune Disord. Drug Targets**, **2006**, 6, 233-237.

<sup>[60]</sup> N. Magné, E. Deutsch, C. Haie-Meder, Current data on radiochemotherapy and potential of targeted therapies for cervical cancers, **Cancer Radiother**, **2008**, 12, 31-6.

<sup>[61]</sup> T. Uno, A. Mitsuhashi, K. Isobe, S. Yamamoto, H. Kawakami, N.Ueno, H .Usui, S. Tate, T. Kawata, H. Ito, Concurrent daily cisplatin and extended-field radiation therapy for carcinoma of the cervix, **Int. J. Gynecol. Cancer**. **2008**, 18, 80-4.

<sup>[62]</sup> B. Bachtiary, I. Herbacek, T. Zideck, T.H. Knocke, J. Dimopoulos, R. Poetter, M. Micksche, Impact of radiotherapy with and without concurrent cisplatin on lymphocyte subpopulations in cervical cancer patients, **Anticancer Res**,. **2005**, 25, 4673-8.

<sup>[63]</sup> J. Kodama, M. Takemoto, N. Seki, K. Nakamura, A. Hongo, S. Moriya, S. Kanazawa, Y. Hiramatsu, Phase I study of chemoradiation with nedaplatin and ifosfamide in patients with advanced squamous cell carcinoma of the uterine cervix, **Int. J. Gynecol. Cancer**. **2008**, PMID: 18284451.

<sup>[64]</sup> (a) Garban Z., Carțiș Ioana, Avacovici Adina, Moldovan I. - Comparative aspects between the interaction of deoxyribonucleic acid with some cytostatic drugs: Particularisation for the action with cis-platinum and cyclophosphamide. I. Investigations in vivo on experi-mental animals, pp.1118-1125, in **Mengen- und Spurenelemente**, 20. Arbeitstagung 2000, Friedrich-Schiller-Universität Jena (Hrsg. Anke M. et al.); Verlag Harald Schubert, Leipzig, 2000., (b) Garban Z., Dancău Gh., Daranyi Gabriela, Erdelean R., Văcărescu G., Precob V., Eremia Iulia, Udriște C. – Implication of chronobiochemistry-metabolism relationship in the induction of homeostasis changes. I. The reaction of cis-platinum on hepatic Dna biosynthesis and on some serum metabolites in rats, **Rev. Roum. Biochim.**, 1989, 26(2), 107-117, (c) Garban Z., Daranyi Gabriela, Tămaș Marta, Miklos J, Borza Iconia, Nemeș Radița, Precob V. - Investigations on the action of cis-platinum on the deoxyribonucleic acid in vitro and in vivo, “**14<sup>th</sup> International Cancer Congress**”, Abstracts of Lectures, Symposia and Free Communications, Vol.3, p.1187, Budapest-Hungary, August 21-27, 1986, Publ.by S.Karger AG-Basel/Akadémiai Kiadó Budapest, 1986, (d) Garban Z., Garban Gabriela, Velcirov Ariana-Bianca Ghibu G.D. - Interaction of deoxyribonucleic acid with cis-platinum : structure-activity relationship, **Rev. Roum. Chimie**, 2007, 52 (1-2), 207-213

<sup>[65]</sup> (a) Garban Z., Aumüller Corina, Daranyi Gabriela, Riviș A., Precob V. - Implications of the chronobiochemistry-metabolism relationship in the induction of



homeostasis changes. IV. The action of  $\text{Cu}^{2+}$  and  $\text{Mn}^{2+}$  on hepatic DNA biosynthesis and on serum proteins, pp. 164-167, in Proceedings of the 8<sup>th</sup> International Symposium "Trace Elements in Man and Animals", Dresden-Germany, May 16-21, 1993 (Eds. Anke M., Meissner D., Mills C.F.), Verlag Media Touristik, Gersdorf, 1993, (b) Garban Z. – Influence of metals on deoxyribonucleic acid, Chap. 7. pp.401-414 in "Elements and their Compounds in the Environment: Occurrence, Analysis and Biological Relevance" Vol. I General Aspects, (Editors Merian E., Anke M., Ihnat M., Stoeppeler M.), 2<sup>nd</sup> edition, Wiley-VCH Verlag GmbH & Co KGaA, Weinheim, 2004 (ISBN 3-527-30459-2), (c) Garban Z. – Aspects of molecular biology and biochemical pathology specific for the interaction of DNA with some chemical xenobiotics of nutritive and therapeutic interest, pp. 215-226, in **Metal Elements in Environment, Medicine and Biology**, Tome VI (Eds. Garban Z., Dragan P.), Publishing House Eurobit, Timișoara, 2004.

<sup>[66]</sup> (a) Dragan P., Garban Z., Daranyi Gabriela, Bucuraș V., Botoca M., Minciu R., Urzică A., Patroi-Popescu Delia, Ehling Șt. - Investigations on the composition and etiopathogeny of lithiasis. XXVIII. Status of blood and urinary metallogram in relationship with ESWL treatment in urolithiasis, pp.173-176, **Metal Elements in Environment, Medicine and Biology**, Tome II (Eds. Garban Z., Dragan P.), Publishing House Eurobit Timișoara, 1996, (b) Garban Z., Dragan P., Daranyi Gabriela, Nemeș Gisela, Șișu E., Șoșdean Doina, Martoni Livia – Etudes biochimiques sur quelques métaux présents dans les urolithiases. II. Détermination de quelques métaux transitionnelles, **Rev. roum. Biochim.**, 1990, 27 (3), 209-216, (c) Garban Z., Dragan P., Nemeș Gisela, Daranyi Gabriela, Sigartău Gr., Șișu E., Mureșan Codruța - Biochemical studies on some metals present in urolithiasis. II. Determination of the main alkaline and alkaline-earth metals, **Rev. roum. Biochim.**, 1987, 24 (3), 221-227.

<sup>[67]</sup> (a) I. Haiduc, Metals in medicine – a promising area of interdisciplinary research (2004, p. 27-34), Organometallic compounds in the environment, medicine and biology (1998, p. 19), Metals in medicine: past, present, future (1997, p. 35), in vol. **Metal Elements in Environment, Medicine and Biology**, Edited by Z. Garban and P. Dragan, Publishing House Eurobit, Timisoara, (b) C. Socaciu, A. Bara, C. Silvestru and I. Haiduc, Antitumor organometallics. II. Inhibitory effects of two diphenylantimony(III) dithiophosphorus derivatives on in vitro and in vivo Ehrlich ascites tumor, **In Vivo**, 1991, 425; (c) A. Bara, C. Socaciu, C. Silvestru and I. Haiduc, Antitumor organometallics. I. Activity of some diphenyltin(IV) and diphenylantimony(III) derivatives on in vitro and in vivo Ehrlich ascites tumor, **Anticancer Research**, 1991, 11, 1651; (d) I. Haiduc and C. Silvestru, Metal compounds in cancer chemotherapy, **Coord. Chem. Revs.**, 1990, 99, 253; (f) I. Haiduc, C. Silvestru and M. Gielen, Organotin compounds: new organometallic derivatives exhibiting antitumor activity, **Bull. Soc. Chim. Belge**, 1983, 92, 187-189.

<sup>[68]</sup> (a) N. Moldovan, P. Lönnecke, I. Silaghi-Dumitrescu, L. Silaghi-Dumitrescu E. Hey-Hawkins, Palladium(II) and Platinum(II) Complexes with Heteroditopic 10-(Aryl)phenoxarsine (Aryl = 2-C<sub>6</sub>H<sub>4</sub>-OR, R = H, Me, Pr) Ligands: Solvent-Oriented Crystallization of cis Isomers, **Inorg. Chem.**, 2008, 47, 1524-1531; (b) E. Fischer Fodor, O. Soritau, N. Moldovan, P. Virag, I. Brie, I. D. Postescu, L. Silaghi Dumitrescu, The CellScan Technology for *In Vitro* Prediction of Platinum-Based Chemotherapeutic Agents, **Int. Symposium on Metallomics**, 2007, Nov 28-Dec 1, Nagoya, Japan, Abstract Book:30. (c) E. Fischer Fodor, N. Moldovan, P. Virag, O. Soritau, I. Brie, P. Lönnecke, E. Hay-Hawkins, L. Silaghi Dumitrescu, The CellScan Technology for *In Vitro* Studies on Novel Platinum Complexes with Organoarsenic

Ligands, ***Dalton Trans.***, **2008**, , 6393 - 6400; (d) J. Castillo, S. Gáspár, S. Leth, M. Niculescu, A. Mortari, I. Bontidean, V. Soukharev, S. A. Dorneanu, A. D. Ryabov and E. Csöregi, Biosensors for life quality. Design, development and applications, ***Sensors and Actuators B: Chemical***, **2004**, B102/2, 179.

<sup>[69]</sup> (a) R. Silaghi-Dumitrescu; B. Reeder, P. Nicholls, C. E. Cooper, M. T. Wilson, Ferryl haem protonation gates peroxidatic reactivity in globins. ***Biochemical Journal***, **2007**, 403, 391–395, (b) R. Silaghi-Dumitrescu, K. Y. Ng, R. Viswanathan, D. M. Jr Kurtz, A Flavo-diiron protein from *Desulfovibrio vulgaris* with oxidase and nitric oxide reductase activities. Evidence for an In vivo nitric oxide scavenging function. ***Biochemistry***, **2005**, 44(9), 3572-9; (c) R. Silaghi-Dumitrescu; Linkage isomerism in nitrite reduction by cytochrome cd1 nitrite reductase. ***Inorg. Chem.***, **2004**, 43(12), 3715-3718.

# EFFECTS OF FOLIAR SELENIUM APPLICATION ON PERFORMANCES IN PLANTS/ANIMALS/MAN

Ivana S. DJUJIC

Department of Chemistry, Institute of Chemistry, Technology and Metalurgy, University of Belgrade, Njegoseva 12, RS-11000 Belgrade, Serbia

E-mail : ankivana@eunet.rs

## ABSTRACT

*Effects of foliar selenium (Se) application to plant food nutritional quality and the nutrient output of agriculture systems in selenium deficient areas and its influence to human needs for nutrients satisfaction are presented. Investigations results obtained during over 10 years of research showing that Se, when added in proper, for each crops specific, amount exerts beneficial effects through multiple mechanisms. Such produced crops have beside Se concentration in the optimal interval, improved quality and, due to higher tolerance to unfavorable external conditions, better yield. Researches conducted on animals that consumed mix feed diet prepared of such produced crops reflected beneficially to animal production and quality parameters of animal products. Examinations of the health benefits that offer consumption of mentioned plant and animal products showed that its contribution to daily Se intake was enough high that Se deficient population assure enough Se in natural form, as well as that, due to higher intake of many interrelated nutrients the benefit that foliar biofortification with Se offer to the Se deficient human population overcomes the benefit of any known Se supplement.*

**Key words:** selenium, foliar supplementation, plants, animal, man.

## INTRODUCTION

Selenium (Se) is of fundamental importance to man and animal health. It is an essential component of several major metabolic pathways, including thyroid hormone metabolism, antioxidant defence systems, and immune function. Its availability and content in food, due to irregularly distribution in the environment, may vary significantly depending on the region. Se protects against many disease problems in man (endemic cardiomyopathy - Keshan disease, bone and joint disease - Keshan-Beck-Syndrome) and animals (growth depression, white muscle disease, exudative diseases, liver necrosis, pancreatic degeneration, reproductive failure) (Tan and Huang, 1991; CAS, 1990; Oldfield, 2002).

Recent research indicates that Se is a versatile protecting agent against a variety of biologic and abiologic pathogenic agents, including certain viruses, toxic heavy metals, ozone, ionizing radiation, mycotoxins and industrial cancer causing compounds. It also positively affects the functions of the cellular or humoral immune system (Fig. 1) (Djujic, 2007)




**Fig. 1.** The health benefits of Se

In order to prevent diseases resulting from Se deficiencies and assure safe and adequate Se intake, the production of high-quality food is essential. Equally important is the knowledge of maintaining food quality through the selection and the preservation of its quality components during food processing and preparation.

Production of food adjusted to consumer demands and environmental issues depends heavily on the availability of mineral and other nutrients to plants. This is why plants, animals and man have several essential nutrients in common.

According to existing knowledge Se is not essential plant nutrient. Its beneficial functions in plants is questionable. Additional nutrients to those required by plants, due to essentiality for man and animals, are Se, I, Cr and Co. Therefore they should be considered in food or feed production, and their deficiencies corrected by appropriate inputs.

Most plant nutrients are taken up from the soil solution. Leaves are able to take up only some forms of nutrient efficient. It is not easy to provide plants with exactly adequate amounts of all nutrients. Innumerable inter-related components that constantly interact causing negative or positive effects influence the extend and magnitude of Se and other nutrients flow through the soil – water – plants – animals – human chain in geographical area, its content and bioavailability in diet.

It is obvious that only multidisciplinary approach to the problem can offer a long-term solution for rapidly growing nutrients deficiency.

On the basis of field trials that we have conducted in Serbia the most feasible method to load enough Se in crops from Se deficient areas is foliar Se application. We invented procedure (*patent No, WO 99/31242*) and formulation (*patent No, WO 99/29639*) for foliar Se application that may assure in crops optimal level and distribution of Se in natural, bioavailable form, and improve negative impacts of Se deficiency to its yield, quality and tolerance to external and internal stressors.

The adopted procedure and formulations for foliar biofortification of food crops with Se is, in comparison with the biofortification through soil, more efficient and environmentally friendly, influences of soil and leaching factors are minimized, acceptability of added Se is fast (*over 50% of absorbed Se plant take in during the first 30-45 min of treatment*), quantity of Se needed for foliar treating is smaller.

Analyses of essential elements, proteins, amino and fatty acids, vitamins, tolerance to oxidative damage and yield of crops showed that foliary enriched plants with adequate Se level are source of high quality food. As such they may considerably improve Se and other essential nutrient intakes by man and animals. (Djujić, 2006).

The purpose of the present study was to presents improvements obtained in nutritional quality and agronomic properties of crops foliary treated with Se and impact of obtained food products use on performances of animals and man as and consumer in the food chain.

## MATERIALS AND METHODS

Assays procedures and methods of investigations conducted in plants, animals and humans are described in our previous publications (Djujic et al, 2000a,2000b, 2002, 2003, 2004, 2005, 2006).

## RESULTS AND DISCUSSIONS

Studies conducted on **crops** in period 1996-2008 showed that over 80% of total Se content is in L- (+) selenomethionine (SeMet) form, the favorable Se forms for man and animals. Obtained data indicates that proper Se range for each crop is different. For soybean it is 10 - 45 times higher than in corresponding control samples, while for wheat it is 10 - 15 times higher.

When Se concentrations in plants are under or above the proper ranges, antioxidant defense and seed quality decreases. Tolerance to higher Se concentrations in investigated plants, as a rule, was better than to lower Se concentrations. Data presented for TBARS, Se-dependent antioxidant enzyme, GSHPx activity and vitamin E contents in wheat and soybean seeds with different Se concentrations shows which Se interval was optimal for obtaining desired improvement in crop performances. (Fig. 2 ).

On the basis of repetitive data it was possible to obtain functional dependence between Se concentration and parameters important for antioxidative defence improvement and/or susceptibility to oxidative stress (Fig. 3 and Fig. 4).

Investigations showed that foliar Se application exerts beneficial effects on crops through multiple mechanisms. Such crops have increased contents of  $\alpha$  - tocopherol (Fig. 5) and some of essential elements which optimal level is slightly lower (Zn, Cu, Mg) (Fig. 6), higher tolerance against detrimental effects of heat (Fig. 7), improved protein (about 10%), amino (Tab. 1) and fatty acids (Tab. 2) contents, yield, as well as tolerance to unfavorable external stressors (high temperature, UV-B radiation, drought).

Presented data, as well as over 15 years of the experience with the low Se soils in Serbia, shows that higher plants, similar as human and animals, require certain for each plant specific Se amounts for plant growth. It is evident that Se deficient crops that on average have 10 to 20 times less Se than are amounts that exert a beneficial effect on them have disrupted growth via several mechanisms.




**Fig 2.** Effects of the various Se concentrations in wheat and soybean seeds on TBARS and vitamin E level and GSH-Px activity




**Fig 3.** Effects of the various Se concentrations in soybean seeds on GSH-Px activity




**Fig. 4.** Effects of the various Se concentrations in soybean seeds on TBARS and vitamin E concentration




**Fig. 5.** Influence of various Se concentrations in soybean seeds on  $\alpha$  - and  $\gamma$  - tocopherol content




**Fig 6.** Effects of foliar Se supplementation on Se, Zn, Cu and Mg in soybean




**Fig. 7.** Changes in TBARS concentration in soybean seeds foliary enriched with different Se concentrations obtained during the exposure to temperature of 45<sup>0</sup>C



**Table 1.** Essential amino acids in Se-rich soybean seeds, Se rich wheat grain and seeds/grains from control groups

| Amino acid concentration, mg/g protein | Soybean no added Se | Soybean added Se | Wheat no added Se | Wheat added Se |
|----------------------------------------|---------------------|------------------|-------------------|----------------|
| Histidine                              | 30                  | 24               | 21                | 23             |
| Isoleucine                             | 51                  | 63               | 34                | 39             |
| Leucine                                | 82                  | 78               | 69                | 72             |
| Lysine                                 | 68                  | 64               | 23                | 29             |
| Met+Cys                                | 33                  | 32               | 36                | 39             |
| Phe+Tyr                                | 95                  | 42               | 77                | 75             |
| Threonine                              | 41                  | 41               | 28                | 30             |
| Tryptophane                            | 14                  | 13               | 10                | 11             |
| Valine                                 | 52                  | 53               | 38                | 42             |
| Total                                  | 466                 | 410              | 336               | 360            |
| Protein, %                             | 35                  | 38               | 13                | 14             |

**Table 2.** Essential fatty acids in Se-rich soybean seed, Se rich wheat grain and seeds/grains from control groups

| of total lipids   | Soybean no added Se | Soybean added Se | Wheat no added Se | Wheat added Se |
|-------------------|---------------------|------------------|-------------------|----------------|
| C <sub>14:0</sub> | -                   | -                | 3                 | 3              |
| C <sub>16:0</sub> | 11                  | 9                | 14                | 12             |
| C <sub>18:0</sub> | 2                   | 3                | 5                 | 4              |
| C <sub>18:1</sub> | 20                  | 19               | 25                | 20             |
| C <sub>18:2</sub> | 64                  | 74               | 49                | 56             |
| C <sub>18:3</sub> | 3                   | 4                | 4                 | 5              |

Obviously, compensational mechanisms that each living organism has are insufficient to overcome stress caused by insufficient Se supplementation. Se deficient plants have significantly lower defense capacity to oxidative stress caused by oxygen radicals produced by internal metabolic and/or external factors. High Se additions are in turn toxic and induce pro-oxidative reactions that enhance lipid peroxidation, and disrupt metabolic harmony in plants. In Se deficiency as in excess oxidative capacity, growth, fertility, quality and nutritive value of obtained crops is reduced. Senescence of crops is faster, what reflects to the decreasing of its nutritive value.

By assuring proper Se supply nutritional value and tolerance to oxidative damage in plants may be significantly increased. This is of exceptional importance for the fast growing magnitude of damages induced by various complex and interrelated stressors in our living environment.

Investigations conducted on **domestic animals** (Japanese quails, chicken, pigs and caws) nourished with: mix feed diet fortified with inorganic Se (Na<sub>2</sub>SeO<sub>3</sub>), mix feed diet fortified with organic Se (brewery SeY) and mix feed diet composed of Se rich ingredients (crops biofortified with Se foliar) showed that the most significant improvement in animal health and nutritional quality of animal products were

obtained at animal that received a mix feed diet composed of crops foliary biofortified with Se.

First investigations were conducted on Japanese quails and the benefits obtained with crops foliary biofortified with Se in mix feed diet surpassed our expectations (Djujic, 2004, 2006, 2007). Essays conducted on chicken showed that such nutrition also act beneficially to its resistance to oxidative stress and significantly improve meat and egg quality and mass.

Analysis of the Se content in laying quail and chicken eggs collected after 1, 3 and 6 months of receiving a feed mix composed of crops bioforified with Se showed that their eggs contained in average 6 times more Se than control groups. The protein analysis showed that Se rich nutrition improve more significant protein content in quail eggs (about 15%) than in chicken eggs (7%).

The n-3 fatty acids analysis in mixed eggs samples collected at 6<sup>th</sup> month of the experiments showed that, in comparison with the control group, eggs contains significantly more n-3 fatty acids (70% in quail eggs and 36% in chicken eggs).

Monitoring of the lipid peroxidation marker (TBARS), an important determinant of egg quality and nutritive value, the deterioration in fresh eggs and during storage, showed that eggs that originates from poultry nourished with feed mixture biofortified foliar with Se have much more delayed development of lipid peroxides and thus their nutritional value remains unchanged much longer (in chicken eggs 3 weeks and in quail eggs 12 weeks of storage) .

Furthermore, such produced poultry eggs have lower cholesterol content (32-50%), much less toxic elements in eggshell and inside part of egg and are richer in many in nutrition often deficient nutrient as are choline, vitamin A, riboflavin, vitamin B12, pantothenic acid, minerals - J, Cu, Cr, Si, Ca, Fe, Zn and essential amino acids than eggs obtained by ordinary farming procedure.

In our experiments on caws only mix feed diet composed of Se rich ingredients (crops biofortified with Se foliar) induced increasing in milk production and significant improvement in its nutritional value.

Analyses conducted on meat of animals that were feed 6 months with mix feed diet composed of crops foliary biofortified with Se showed that such meat has improved Se and other essential elements concentration, as well as amino and fatty acid composition. Antioxidative enyzmes (GSHPx, SOD, CAT) and lipid peroxidation marker monitoring in them showed that tissues of animals nourished with feed mix composed of ingredients biofortified with Se have lower autoxidation of unsaturated fatty acids in which poultry meat is particularly rich and therefore longer shelf life.

Obtained data indicates that proper level of natural Se form in feed mix plays a key role in maintaining effective protection in various tissues against the damaging effects of free radicals and toxic products of their metabolism. Post-slaughter biochemical changes involved in the conversion of muscle to meat that are accompanied by a loss of cellular antioxidant defences and an increased propensity of meat lipids to undergo oxidation. Proper level and quantity of Se in nutrition of poultry contributes that undesirable changes in a number of quality parameters, including loss of water-holding capacity, texture and flavour became lower. As the lipids in poultry exhibit a higher degree of unsaturation than red meats due to a relatively high content of phospholipids, important factor for the oxidative stability of meats represent the degree of unsaturation of the phospholipids of the subcellular membrane. To improve oxidative stability, sensory quality, shelf life and acceptability of poultry now adding higher level of dietary antioxidants what, our experimental data shows, is not necessary when animal diet contains the proper level and form of Se.

On the basis of experimental data presented for animals, it can be concluded that foliar biofortification with Se offers following benefits to consumers and animals and its products producers:

- Eggs, milk and meat of such farmed animals represent an extremely rich source of natural Se forms and have significantly improved contents of many other, commonly deficient nutrients as are highly valuable proteins, omega-3 fatty acids, minerals, antioxidants).
- Content of undesirable toxic substances and its metabolic products was reduced.
- Body and egg mass, as well as eggs and milk production was higher in such feed animals.
- Animal products (eggs, milk and meat) have significantly longer shelf life and other quality parameters (loss of water binding capacity, texture, sensory quality).
- The consumption of products of such farmed animals may assure to man a daily needs for Se, highly nutritionally scored proteins, fat with the optimal omega-6/omega-3 ratio.

Comparison of data obtained on **population** groups that consumed supplements with 100 µg Se in the form of bakery SeY in the course of 8 weeks and products made from wheat foliar biofortified with Se over 6 weeks showed that efficiency of daily intake on average about 18 µg Se (12 - 35 µg) from Se enriched wheat influence in a shorter time effects comparable with the SeY supplement use. Increase of Se concentration in both groups were about 53% in plasma, 37% in erythrocytes, 44% in hair and 54% in urine (Djujić et al, 2000a). Data for oxidative stress reduction monitoring in blood plasma showed that although contribution of wheat biofortified with Se to daily selenium intake was over 5 times lower and in both cases the dominant Se-form was L – (+) Se-methionine, consumption of product obtained from Se enriched wheat reduced more efficiently oxidative stress (Djujić et al, 2000b).

In a Norwegian study it was demonstrated that high bioavailability of Se from wheat by feeding trial participants with Se-rich bread providing 100, 200 or 300 µg Se daily for 6 weeks. Serum selenium increased in a dose-response manner by 20, 37 and 53 µg/l, respectively, in the three groups ( $p < 0.001$ ) (Lyons et al, 2003).

Inclusion in diet soybean and other crops foliary biofortified with Se contributed that Se intake by Serbian residents increased by 170 %.. Investigations conducted on participants in this study showed that inclusion of crops biofortified with Se in diet did not only influence the increase of Se concentration and GSHPx activity in blood plasma and erythrocytes but optimized many other, for antioxidant defense and anti-inflammatory response important parameters (Djujić et al, 2002a, 2002b). Similar correlations between Se intake and Se and GSHPx levels in plasma and erythrocytes were found only in expressive Se deficient population groups from China and New Zealand (Xia et al, 1992; Whanger, 1998; Whanger and Butler, 1988; Schrauzer, 2000) confirming that the degree of deficiency influences Se absorption.

Monitoring of values obtained for lipid peroxidation in healthy subjects showed that good negative correlation exists between Se and TBARS ( $r \geq -0.546$ ). Investigations of lipid metabolism parameters in serum of investigated subjects showed that increased natural Se intake improves the level of these parameters.

Examinations of the health benefits that consumption of products prepared with crops foliary biofortified with Se and eggs obtained from quails that consumed feed mix composed of biofortified crops showed that its contribution to daily Se intake was high

enough that Se deficient population of Serbia can assure even more Se in natural form than is RDA (Fig. 8). Investigations of Se, concentration and GSHPx activity in blood plasma before the beginning with Se rich food consumption and after 6 weeks of consuming Se rich food showed that increased Se intake reflected significantly to GSHPx activities, but that volunteers did not in the mentioned period reach GSHPx activities accepted as optimal for healthy individuals. Obviously, time of increased Se consumption was insufficient to fill up all Se pools in the body. Lipid peroxidation marker monitoring in plasma of the mentioned groups of volunteers showed that increase of Se intake acts beneficial to oxidative stress in Se deficient humans (TBARS values in plasma decrease significantly in all groups). The health-related quality of life (HRQoL) surveys obtained on the basis of the respondent's reports to five domains of function at start and the end of investigation indicate that in all investigated groups, food rich in natural Se improved more or less all five domains of HRQoL functions (mobility, self-care, usual activities, pain/discomfort and anxiety/depression) (Djujic at al., 2005).




**Fig. 8.** Influence of food rich in natural Se consumption by Se deficient volunteers on its average daily intake, concentration in plasma and lipid peroxidation

Recent examinations, conducted on equipment that for diagnostic functioning of organism uses energy-information inter-exchange confirmed that consumption of food prepared from biofortified crops and/or animal products (eggs, meat, milk) obtained of animals that consumed crops biofortified with Se offer us possibility to prepare right combination for each in nutrient deficient individual and, at the same, time shows that none supplement can be so efficient as natural sources of essential and beneficial nutrients (Djujic and Lalic, 2008).

## CONCLUSIONS

1. Monitoring the magnitude and nature of oxidative damage induced by various complex and interrelated environmental stressors, showed that Se deficiency is responsible for a higher susceptibility of plants, animals and man to oxidative damage.

2. Foliar Se supplementation may assure optimal Se interval in plants and optimize many for plants beneficial nutrient content what significantly increase tolerance to oxidative damage in plants.
3. Foliar Se supplementation effectively alleviate primary Se deficiency consequences in agriculture and at the same time offer solutions for the secondary problems in crops and animal breeding, assuring production of highly valuable nutritious food and formulation of nutritionally well balanced diets for increasingly malnourished consumers.
4. Health benefits that in synergistic action assure adequate Se intake and optimized intake of many essential and beneficial nutrients through the food chain were incomparable with effect that offer supplementation with chemical forms provided in artificial powders, supplements or pills as well as food fortification, and some of procedures for biofortification.
5. And results of such interventions were that living organisms in food chain have better developed antioxidative defence, and other compensational mechanisms of importance for defence against internal metabolic or external factors. This is of exceptional importance for the fast growing magnitude of oxidative damage induced by various complex and interrelated stressors in our living environment.
6. Examinations of the health benefits that offer consumption of mentioned plant and animal products showed that its contribution to daily Se intake was enough high that Se deficient population of Serbia can assure even more Se in natural form then is RDA, as well as that, due to higher intake of many interrelated nutrients, the benefit that foliar biofortification with Se can offer to the deficient human population overcomes the benefit of any known Se supplement.

## REFERENCES

1. Djujic I., Demajo M., Jozanov-Stankov O., Markovic Lj.: Improvement of Japanese quail (*Coturnix Coturnix Japonica*) meat and egg nutritional qualities: supplementation of livestock feed with organic selenium including foliar treatment of feed crops, Metal Elements in Environment Medicine and Biology, Eds.: Z.Garban, P.Dragan et al., 6th Int.Symp. of Roumanian Academy-Branch Timisoara, 2004, Vol. VI, 53-60.
2. Djujic I., Jozanov-Stankov O., Demajo M., Milovac M.: Tolerance to oxidative damage in plants, animals and humans in dependent on selenium concentration, In: Macro and Trace Elements, Agrucultural, Biological and Medical Importance of Macro, Trace and Ultratrace Element, 21. Workshop, Friedrich Schiller University Jena, Eds. M. Anke, . et al., Shubert-Verlag Leipzig, 2002, 886-898.
3. Djujic I., Jozanov-Stankov ON, Milovac M, Jankovic V, Djermanovic V.: Bioavailability and possible benefits of wheat intake naturally enriched with selenium and its products. *Biol Trace Elem Res.*, 2000a, 77: 273-285.
4. Djujic I., Milovac M.: Combination used for the foliar supplementation of plants with Se salts and urea, *Patents No.* WO 99/29639, 1998.
5. Djujic I., Milovac M.: Procedure for foliar supplementation of plants with Se salts, *Patents No.* WO 99/29624, 1998.
6. Djujic I.: Trace element deficiency treatment: much more than supplementation with deficient trace element, 2008, *Cell Biol Toxicol.*, 2008, 1, 114-115.
7. Djujic I, Djermanovic V, Jozanov-Stankov O.: Dietary intake of macro and trace elements in Serbia, In: M. Anke et al. (HRSG), *Mengen-und Spurenelemente*, 20. *Arbeitstagung*, Verlag Harald Schubert, Leipzig, 2000b, 787-796.

8. Djuić I., Demajo M., Jozanov Stankov O., Marković Lj.: Health benefits of consuming selenium-enriched quail eggs, 5<sup>th</sup> International Symposium on Trace Elements in Human: New Perspectives, Athens Greece, proc., 2005, 622-635.
9. Djuić I., Milovac M., Djermanović V., Jozanov-Stankov O.: Prevention of selenium deficiency in humans by selenium rich soybean, *Trace Elements in Medicine*, 2003, 1, 55-62.
10. Djuić I.S.: Benefits of raising crops and animals with adequate selenium content in low selenium areas, Trace Elements in the Food Chain, Eds Szilagyi M and Szentmihalyi K., 2006, Hungarian Academy of Sciences, 15-20.
11. Lalic T. , Djuić I., Elements in the hair of population with different grade of hypothyroidism, 2008, *Cell Biol Toxicol.*, 2008, 1, 42.
12. Lions G.H., Ortiz-Monasterio I., Strangoulis JCR., Graham R.D.: Selenium concentration in wheat grain: Is there sufficient genotypic variation to use in breeding? *Plant Soil*, 2004, **269**, 369-380.
13. Oldfield J. E.: Selenium world atlas (Updated edition), *Selenium-Tellurium Development Association* (STDA), 2002, [www.stda.org](http://www.stda.org).
14. Schrauzer G.N. Selenomethionine: A Review of Its Nutritional Significance, Metabolism and Toxicity, *Am. Soc Nutr. Sci.*, 2000, **130**, 1653-1656.
15. Tan J.A., Huang Y.: Selenium in geo-ecosystems and its relation to endemic diseases in China. *Water, Air, and Soil Pollution.*, 1991, **57-58**: 59-68.
16. Whanger P.D.: Metabolism of selenium in humans. *J of Trace Elements in Experimental Medicine.*, 1998, **11**, 227-240.
17. Whanger, P. D., and Butler J.: Effects of various dietary levels of selenium as selenite or selenomethionine on tissue selenium levels and glutathione peroxidase activity in rats. *J. Nutr.*, 1988, **11**, 846-852.
18. Xia Y., Zhao X., Zhu L., Whanger P.D.: Metabolism of selenium and selenomethionine by a selenium deficient population of men in China., 1992, *J. Nutr. Biochem.* **3**, 202-210.
19. \*\*\*CAS: The group of Environmental and Endemic Diseases, Institute of Geography - Kashin-Beck disease in China. Geographical epidemiology and its environmental pathogenicity. *The Journal of Chinese Geography.*, 1990, 1: 71-83.

# NORMAL AND PATHOLOGICAL INVOLVEMENT OF BIVALENT CATIONS IN ORO-MAXILO-FACIAL AREA

Mihai NECHIFOR \*

Department of Pharmacology, University of Medicine and Pharmacy "Gr. T. Popa" Iași, Str. Universității 16, RO-700 115 Iași, Roumania

\* E-mail : mnechif@yahoo.com

## ABSTRACT.

*Bivalent cations have multiple actions in the oro-maxilar area. In case of parotide gland neoplasma, our data revealed significant increase in salivary magnesium concentration and a decrease in plasma zinc concentration. In maxilar sinusitis, there are no changes in calcium and magnesium concentration, whereas plasmatic zinc level is low. Zn, Cu, Ca, Mg and Mn also play important roles in the synthesis of dental enamel, and misbalances are involved in the pathogeny of tooth decay and other maxillary diseases.*

**Key words:** Ca, Mg, Zn, parotide cancer, sinusitis, teeth.

Bivalent cations have multiple roles in normal functioning of all systems in the human body. They are involved in regulating the activity of lots of enzymes and influence cell functioning in complex ways.

Misbalances in the ionic concentrations of these cations (inside and outside the cell) deeply influence normal functioning of the human body. The oro-maxilo-facial area exposes a series of important particularities and complex functional roles. Some of the characteristics particular to this area are:

- the contact of the oral mucosa with blood and saliva
- the interaction between airways and digestive pathway
- phonation and deglutition headquarters
- teeth presence
- others.

Bivalent cations play important role is in the functioning of all the cells and salivary secretion.

Misbalances in intra and extra cell concentration of these cations are implicated in a series of pathogenic conditions.

## Bivalent cations in saliva

Salivary concentrations of bivalent cations are a reflection of the saliva-secreting processes. This salivary secretion is realised from blood plasma.

Zn, Mg, Cu, Ca as well as other cations are found in the composition of sublingual calculi. Zinc and zinc-containing salts seem to be very important in this calculi formation (Knuntilla M et al, 1981).

There are differences in the composition of bivalent cations in supra and subgingival calculi.  $Mn^{2+}$  concentration is significantly higher in supragingival calculus.  $Zn^{2+}$  concentration was 5.4 times higher subgingival calculus than in supragingival calculus (Knuntilla M et al, 1979 and 1981).

In case of acetyl-choline (Ach)-evoked amylase secretion of the parotid gland in a medium containing 1.1mM  $Mg^{2+}$ , Ach significantly increase the amylase secretion. Absence of magnesium or elevated magnesium concentration (5-10 mM) reduced Ach-evoked amylase secretion.

Ach stimulation of the parotid gland elevate the saliva  $Mg^{2+}$  concentration. The increase of is an active process, since  $Mg^{2+}$  has to move against the gradient.

Some medicines such as lidocaine (frequently used as local anaesthetic in dental practice), quinidine, amylorid, etc determine modification in the oral concentration of this cation.

In patients with chronic renal failure, changes in salivary ionic composition are frequently identified (Martins C. et al, 2006) Hemodialysis also determine changes in saliva concentration and saliva flux.

The level of  $Ca^{2+}$  in saliva of diabetes mellitus patients was significantly elevated compared to healthy-matched controls. The concentration of  $Zn^{2+}$  and  $Mg^{2+}$  were significantly reduced (Mato AD, et al, 2004).

Una et al., 2004 showed that on chronic rheumatoid arthritis children the plasma level of zinc and copper is significantly lower than in the control group, but plasma magnesium level did not differ.

### **Bivalent cations in teeth pathology**

Research performed by X-ray florescent techniques showed increased copper and zinc concentrations in canins regions of dentine compared with normal part of the teeth (Haris HH et al, 2008). Increased zinc concentration in the hydroxyl apatite was noticed in tooth decay regions.

We consider that the  $Cu^{2+}/Zn^{2+}$  ratio in tooth level and saliva might be correlated with the evolution and occurrence of tooth decay.

On the other hand, zinc citrate from tooth paste reduce by 17-49% the anaerobic bacteria and streptococci at the level of the oral cavity.

Additionally, in situ microbial biofilm formation was significantly inhibited in the zinc citrate group ( $p < 0.05$ ) (Srenivoisan et al., 2008).

The dental concentrations of some cations may suffer small changes with aging and they are not the same in enamel as well as in dentine.

Derise and Ritchey date (1974) show that manganese concentrations in the enamel is  $7.41 \pm 0.13$  ppm for 10-12 years and  $6.67 \pm 0.12$  ppm for 25 years. Zinc concentration in the same age groups was  $190 \pm 4.1$  ppm vs  $180.3 \pm 3.8$  ppm ( $p < 0.05$ ). Manganese concentration in dentine decreases from  $6.20 \pm 0.16$  ppm in 10-12 years group of age to  $5.52 \pm 0.15$  ppm after 25 years. The evaluation of zinc concentration differences between sexes indicated that zinc was significantly higher in both enamel and dentine in females ( $p < 0.01$ ).

Manganese concentration was significantly higher in the enamel than in dentine. Maximum manganese concentration both in enamel and dentine can be found in children under 12 years old. After this age, it decreases, and this fact can be linked to dentine evolution in puberty. Zinc enamel concentrations differences between sexes are highly significant (higher values for female). No significant differences were found in Fe concentrations in enamel and dentine between sexes (Knychalska-Karwan et al., 1985).

Copper concentration is important for the dental osteoclasts activity stimulation.

On the contrary, increased zinc concentration in the osteoclasts environment do not modify tooth resorption process (Li BB, Yu S.F., 2007). Mg ions have no effect on osteoclasts apoptosis and active ring formation (Li X et al., 2008).



Hashemi et al., 2005 showed that the Ca/P ratio of was caries-affected dentine structures. After analysis of human molars extracted with coronal carians lesions was founded that CA/P ratio of caries was less than caries-free dentin surface. Milk carietic teeth contain less Mg, Mn and Cu and much more Zn as compared to normals (Niedzielska, 1990). Zinc concentration was higher both in the crown and the root of milk teeth than permanent teeth. Zinc level is higher in the teeth of the children than in the adult teeth (Gierat-Kucharzewska, 2003).

### **Cancers of oro-maxilar area**

Cancer represents one of the most important medical problems. The malignant tumours of the oro-maxilar area represent 1-1.5% of human neoplasm. Salivary gland malignant tumours are a heterogenous group of cancers. The most important malignant parotide tumours anatopathological aspects are: adenocarcinoma, mucoepidermoid carcinoma and acinic cell carcinoma.

The frequency of parotide gland tumors is increasing now. The permanent contact of parotid gland tissue with saliva can be important for normal cell activity and some modifications of saliva composition can influence the malignant tumour evolution.

In our studies (Nechifor et al., 2005; Gradinaru et al., 2007), we found that plasma and saliva total-Mg concentrations was significantly higher in parotid malignant tumours patients vs. healthy controls. Plasma copper concentration of patients with parotid neoplasma was significantly higher compared to normal control and plasma zinc concentration was lower in parotid malignant tumour ( $0.017 \pm 0.01 \text{ mmol/L}$  vs.  $0.024 \pm 0.01 \text{ mmol/L}$  in healthy controls,  $p < 0.01$ ).

Zinc plasma concentrations reduction can enhance the neoplasma development by reduction of the immunity (Fong et al., 2006). There are data regarding the role played by zinc in the reduction or prevention of oncogenesis. Bitar et al., 2007 showed that zinc might be an adjustment therapy for juvenile onset recurrent papillomatosis.

Changes in plasma and tissue concentration of some cations have been also identified in case of different forms of cancer (Tanaka et al., 2000).

In whole saliva collected from oral squamos carcinoma patients, magnesium concentration was significantly higher by 28%, compared to saliva people (Shpitzer T. et al., 2007). The albumin and K was lower in saliva collected from squamos in the most common head and neck malignant and is generally induced by tobacco exposure. Saliva annalasis is a reliable indicator for early diagnosis (Reznick, 2004).

Saliva play an important role in the pathogenesis of oro-pharyngeal neoplasma. In patients with xerostomia induced by cancer radiotherapy the flow rate of stimulated whole saliva decreased 83% and this volume reduction was accompanied by a significant increase in Ca and Mg concentrations (Dreizen et al., 1976).

### **The taste**

The exact mechanisms of gustative sensations development and influencing by concentrations of the bivalent cations are not yet very clear.

There are data that show calcium and magnesium air may be involved in taste perception. T1R3 receptor phenotype expression of TAs 1r3 taste receptor Tass1r3 taste receptor gene function as a gustatory calcium-magnesium receptor (Tordoff MG et al, 2008).

Oral calcium elicited less electrophysiological effect in the chorda tympani nerve in Tas1r knockout mice.

### The oro maxillar infections

We consider that ratio between some bivalent cations concentration in plasma and saliva are very important for the evolution of some infections in the oro-maxillar area.

We surveyed the plasmatic and salivary concentrations of different bivalent cations in patients with oro-maxillo-facial area bacterial infections.

We noticed the  $\text{Ca}^{2+}$ ,  $\text{Mg}^{2+}$ ,  $\text{Cu}^{2+}$ ,  $\text{Zn}^{2+}$  plasmatic concentrations, the  $\text{Ca}^{2+}$ ,  $\text{Mg}^{2+}$  salivary concentrations for 71 adults patients, both sex, hospitalized for soft tissues suppurations in the oro-maxillo-facial area (study group) and for 60 healthy volunteers, with a similar distribution upon age and sex (the control group). The suppurations were produced by different types of bacterial species: Gram-positive anaerobic cocci, facultative anaerobes (*Enterococcus spp*, *Streptococcus pneumoniae*, etc). The determinations were done by absorption spectrometry, before the beginning of the drug or surgical therapy, during the morning 12 hours after the last food ingestion. In this study we didn't include patients with other infections, pregnancy, liver cirrhosis, renal insufficiency, malabsorption syndrome, diarrhea, patients treated by diuretics/ cardiac glycosides/ antihypertensive drugs or other drugs containing cations.

In the oro-maxillo-facial bacterial infections the salivary magnesium concentration is statistically significant increased in comparison to the control group ( $4,76 \pm 0,78$  mg/l vs  $3,37 \pm 0,65$  mg/l,  $p < 0,05$ ). The salivary Mg /Ca ratio in the study group is statistically significant increased in comparison to the control group. The zinc plasmatic concentration is statistically significant decreased in the study group in comparison to the control group ( $1,18 \pm 0,51$  mg/l vs  $1,53 \pm 0,20$  mg/l,  $p < 0,05$ ).

We consider that an increased Mg /Ca salivary ratio correlated to a decreased  $\text{Zn}^{2+}$  plasmatic concentration can be considered as a biological indicator of a suppurative process in the oro-maxillo-facial area and can favour the development of the bacterial infections in the mentioned area.

Our data (Nechifor et al., 2006) are in concordance other that show that Mg-dependent ATP-ase is involved in the invasion of salivary gland by some microorganisms like *Rhodnius prilius* (Fonseca FV et al., 2006).

Periodontitis are bacterial infections of both tooth suppurating tissues. The essential elements of periodontitis are inflammation and teeth loss. This disease is frequent, being some of the most problems in dental medicine and some of the most frequent infections world-wide (Xu Y et al., 1999). Patients with periodontitis have decreased total saliva Ca and decreased serum Zn and Ca (Kuraner, 1991). Meisel et al., 2005 showed that magnesium deficiency is associated with periodontal disease. A cross-sectional investigation involving 4290 subjects aged 20-80yrs showed that increased serum Mg/Ca ratio was significantly associated with reduced probing depth ( $p < 0.001$ ) and a higher number of remaining teeth ( $p < 0.005$ ) (Meisel et al., 2006). Higher serum Mg/total Ca ration can lower the periodontitis incidence (Meichant et al., 2006).

Our data show that increased calcium favours periodontitis.

Subjects with calcium antagonist treatment had significantly deeper gingival pocket and the probing depth was increased (Meisel, 2005).

Another aspect of the implication of the bivalent cations in saliva and teeth in oral cavity pathology is constituted by the possibility that some of these cations possibly influencing dental enamel characteristics and biofilm formation. The biofilm is essential for the development of bacteria in the dental plaque. Mittrakul K et al., 2004 showed that copper can influence bacterial biofilm detachment in the *Streptococcus gordonii* imitation of dental plaque development.

Situation seem to be different in case of fungal infections.

Calcium blocks fungicidal activity of salivary histatin 5 which kills the fungal pathogen *Candida albicans*.

A high level of salivary calcium can favorize the development of fungicidal oral infections. Histatins constitute a group of cationic multifunctional proteins with antifungal activity. Their activity depends on cationic concentrations in the environment. They possess high fungicidal activity against azole-resistant fungal species (Tsai H, Bobek LA, 1998).

We studied Ca, Mg, Zn and Cu plasma concentrations and t-Ca and t-Mg levels in saliva in patients with chronic maxillary sinusitis without suppuration. Our results (fig. nr. 1 and fig. nr. 2) showed that in chronic maxillary sinusitis without suppuration no changes in t-Mg and t-Ca, but plasma zinc level is lower compared to normal control group ( $10.82\text{mg/mL} \pm 0.24$  vs.  $1.46\text{ mg/mL} \pm 0.27$ ,  $p < 0.01$ ). The significant decrease in serum zinc level a reduction of lymphocytes and macrophages immunity involvement. This decreased level of plasma zinc concentration can favourise the development of chronic sinusitis by reduction of defence of the human body against microbial infections.




**Fig.1** Salivary concentrations of calcium and magnesium in patients with maxillary sinusitis




**Fig. 2.** Plasmatic concentrations of copper and zinc in patients with maxillary sinusitis

#### The influence on some locally administered drugs

Considering the use of the periodontal gel of some drugs (such as metronidazol), the bioadhesive properties of the gels was affected was affected by the variations of the  $\text{Ca}^{2+}$  concentrations in saliva (Varshosaz J et al., 2002).

Bivalent cations play important roles, unable to be disregarded, in the oro-maxilar area.

## REFERENCES

1. Bitar M, Baz R, Fuleihan N, Muallem M.: Can zinc be an adjuvant therapy for juvenile onset recurrent respiratory papillomatosis? *Int J Pediatr Otorhinolaryngol.* 2007;71(8):1163-73
2. Derise N.L., Ritchey S.J.: Mineral composition of normal human enamel and dentin and the relation of composition to dental caries. II. Microminerals. *J Dent Res.* 1974;53(4):853-858
3. Dreizen S, Brown L.R., Handler S, Levy B.M.: Radiation-induced xerostomia in cancer patients. Effect on salivary and serum electrolytes. *Cancer.* 1976;38(1):273-278
4. Fonseca F.V., Fonseca de Souza A.L., Mariano A.C., Entringer P.F., Gondim K.C., Meyer-Fernandes J.R.: Trypanosoma rangeli: characterization of a Mg-dependent ecto ATP-diphosphohydrolase activity. *Exp Parasitol.* 2006;112(2):76-84
5. Fong L.Y., Jiang Y, Farber J.L.: Zinc deficiency potentiates induction and progression of lingual and esophageal tumors in p53-deficient mice. *Carcinogenesis.* 2006;27(7):1489-1496
6. Gierat-Kucharzewska B, Braziewicz J, Majewska U, Góźdz S, Karasinski A.: Concentration of selected elements in the roots and crowns of both primary and permanent teeth with caries disease. *Biol Trace Elem Res.* 2003;96(1-3):159-167

7. Grădinaru I, Ghiciuc C.M., Popescu E, Nechifor C, Mândreci I, Nechifor M.: Blood plasma and saliva levels of magnesium and other bivalent cations in patients with parotid gland tumors. *Magnes Res.* 2007;20:254-258
8. Harris H.H., Vogt S, Eastgate H, Legnini D.G., Hornberger B, Cai Z, Lai B, Lay P.A.. Migration of mercury from dental amalgam through human teeth. *J Synchrotron Radiat.* 2008;15(Pt 2):123-128
9. Hashemi Z.K., Oshida Y, Deely J.J., Ki Y.: Ca/P mol ratio of caries-affected dentin structures. *Biomed Mater Eng.* 2005;15(4):251-260
10. Knuuttila M, Lappalainen R, Kontturi-Närhi V.: Concentrations of Ca, Mg, Mn, Sr and Zn in supra- and subgingival calculus. *Scand J Dent Res.* 1979;87(3):192-6
11. Knuuttila M, Lappalainen R, Lammi S.: Zn concentration of human subgingival calculus related to F, Mg and Cu contents. *Scand J Dent Res.* 1981;89(5):412-416
12. Knychalska-Karwan Z, Pawlicki R, Karwan T.: Quantitative, qualitative and topographic distribution of lead, zinc, iron and copper in hard teeth tissues. *Folia Histochem Cytobiol.* 1985;23(1-2):21-25
13. Kuo H.W., Chen S.F., Wu C.C., Chen D.R., Lee J.H. : Serum and tissue trace elements in patients with breast cancer in Taiwan. *Biol Trace Elem Res.* 2002;89:1-11
14. Kuraner T, Beksac M.S., Kayakirilmaz K, Çağlayan F, Onderoğlu L.S., Özgünes H. : Serum and parotid saliva testosterone, calcium, magnesium, and zinc levels in males, with and without periodontitis. *Biol Trace Elem Res.* 1991;31(1):43-49
15. Li BB, Yu SF.: In vitro study of the effects of copper ion on osteoclastic resorption in various dental mineralized tissues, *Zhonghua Kou Qiang Yi Xue Za Zhi.* 2007;42:110-113
16. Li X, Senda K, Ito A, Sogo Y, Yamazaki A.: Effect of Zn and Mg in tricalcium phosphate and in culture medium on apoptosis and actin ring formation of mature osteoclasts. *Biomed Mater.* 2008 25;3:45002
17. Martins C, Siqueira W.L., Guimarães : Primo LS. Oral and salivary flow characteristics of a group of Brazilian children and adolescents with chronic renal failure. *Pediatr Nephrol.* 2008;23:619-24
18. Mata A, Marques D, Martínez-Burgos M.A., Silveira J, Marques J, Mesquita M.F., Pariente J.A., Salido G.M., Singh J. : Magnesium-calcium signalling in rat parotid acinar cells: effects of acetylcholine. *Mol Cell Biochem.* 2008;307:193-207
19. Meisel P, Schwahn C, Luedemann J, John U, Kroemer H.K., Kocher T.: Magnesium deficiency is associated with periodontal disease. *J Dent Res.* 2005;84:937-41
20. Mitrakul K, Loo C.Y., Hughes C.V., Ganeshkumar N.: Role of a *Streptococcus gordonii* copper-transport operon, *copYAZ*, in biofilm detachment. *Oral Microbiol Immunol.* 2004;19:395-402. Erratum in: *Oral Microbiol Immunol.* 2005;20:65
21. Nechifor M, Grădinaru I., Popescu E., Gogălniceanu D, Mândreci I, Nechifor C: Variations of plasmatic and salivary concentrations of different bivalent cations in oromaxillo-facial area bacterial infections, *Metal ions in biology and Medicine*, Ed. John Libbey, Eurotext paris 2006, Eds. Maria Carmen Alpoim, Paula Vasconcellos, Maria Amelia santos, Armando J Cristovao, Jose A Centro, Philippe Colley, Vol. 9, p. 556-560
22. Nechifor M., Grădinaru Irina, Gogălniceanu D. , Mândreci I.: Blood plasma and saliva concentration of some bivalent cations in patients with ora-maxilar cancer, *Trace Elements and Electrolytes*, 22, 2005, p. 71
23. Nechifor M. Grădinaru Irina, Mândreci I., Gogălniceanu D.: Magnesium Plasma and saliva levels of magnesium in sinusitis, 11th International Magnesium

- Symposium, Kashikojima, 23-26 Octombrie 2006, Abstract Book, J Jap Soc for Magn Res, Vol. 25, No. 2, 2006, p. 101
24. Niedzielska K, Struzak-Wysokińska M, Wujec Z. : Analysis of correlations between the content of various elements in hard tissues of milk teeth with and without caries, *Czas Stomatol.* 1990;43:316-322
  25. Pussinen P.J., Paju S, Mäntylä P, Sorsa T.: Serum microbial- and host-derived markers of periodontal diseases: a review. *Curr Med Chem.* 2007;14:2402-2412
  26. Reznick A.Z., Hershkovich O, Nagler R.M. : Saliva--a pivotal player in the pathogenesis of oropharyngeal cancer. *Br J Cancer.* 2004;91:111-118
  27. Shpitzer T, Bahar G, Feinmesser R, Nagler R.M.: A comprehensive salivary analysis for oral cancer diagnosis. *J Cancer Res Clin Oncol.* 2007;133:613-617
  28. Sreenivasan P.K., Furgang D, Markowitz K, McKiernan M, Tischio-Bereski D, Devizio W, Fine D.: Clinical anti-microbial efficacy of a new zinc citrate dentifrice. *Clin Oral Investig.* 2008 11
  29. Tanaka M, Matsunaga K, Kadoma Y.: Correlation in inorganic ion concentration between saliva and plaque fluid. *J Med Dent Sci.* 2000;47:55-59
  30. Teraki Y, Ishiyama M. Studies on the distribution of mineral elements in the tooth of zinc-deficient rats. *Shigaku.* 1990;78:251-257
  31. Tordoff M.G., Shao H, Alarcón L.K., Margolskee R.F., Mosinger B, Bachmanov A.A., Reed D.R., McCaughey S.: Involvement of T1R3 in calcium-magnesium taste. *Physiol Genomics.* 2008;34:338-348
  32. Tsai H, Bobek LA. Human salivary histatins: promising anti-fungal therapeutic agents. *Crit Rev Oral Biol Med.* 1998;9(4):480-497
  33. Una M, Tamer L, Pata Y.S., Kilies S, Dec Germanic U, Akbas Y. : Serum levels of antioxidant vitamins copper and magnesium in patients with chronic rheumatoid arthritis, *J. Trace Elem and Electrol Med Biol*, 2004, 18, 189-192
  34. Varshosaz J, Tabbakhian M, Zahrooni M. : Development and characterization of floating microballoons for oral delivery of cinnarizine by a factorial design. *J Microencapsul.* 2007;24:253-623
  35. Xu Y, Ambudkar I, Yamagishi H, Swaim W, Walsh T.J., O'Connell B.C.: Histatin 3-mediated killing of *Candida albicans*: effect of extracellular salt concentration on binding and internalization. *Antimicrob Agents Chemother.* 1999;43:2256-2262

# POTENTIALLY METALLOENZYME INHIBITOR HYDROXAMIC ACIDS: EFFECTS OF AN IMIDAZOLE DONOR ON THE METAL BINDING CAPABILITY

Etelka FARKAS\*, Edit CSAPÓ, Péter BUGLYÓ

*\*Department of Inorganic and Analytical Chemistry, University of Debrecen, H-4010 Debrecen, Hungary*

\* E-mail : efarkas@delfin.unideb.hu

## ABSTRACT

*The role of  $R_C$  substituents, involving imidazole moiety, in complexation of various potentially metalloenzyme inhibitor hydroxamic acids,  $R_C$ -CO-N( $R_N$ )-OH, (derivatives of amino acids and small peptides) with Cu(II), Ni(II) and Zn(II) ions has been investigated using pH-potentiometry and different spectroscopic methods. The main factors found to affect the competition of the imidazole with the hydroxamate moiety for the metal ions (e.g. the distance between these two functions, presence/absence of additional donor in the  $R_C$  chain, sort of metal ion) are planned to evaluate in the presentation.*

**Keywords:** hydroxamic acid, imidazole, metal complexes, metalloenzyme inhibitors









## INTRODUCTION

Although the coordination properties might be different and affected by numerous factors, but if an imidazole is present in a molecule, e.g. either in amino acids or in derivatives [1], in small peptides [2-6] or even in large protein molecules [7], very often this group is one of the most effective metal binding sites (often the most effective one) of that molecule. For example, in a large number of natural biomolecules, e.g. metalloproteins, this type of group plays a predominant role in the metal binding. Moreover, there is also a great affinity of an imidazole group for different non-covalent interactions (hydrogen bond, stacking interaction).

On the other hand, if a hydroxamic function is situating in a molecule, this can also play a determinant role in the metal binding (a five-membered chelate, via the two oxygen donors, is most frequently formed). The biological role of the hydroxamate based siderophores [8] and also the known metalloenzyme inhibitory effect of hydroxamic acids are highly based on the strong metal binding activity of the hydroxamate function [9,10]. Great effort has been made during the past years to develop selective inhibitors to various metalloenzymes, e.g. to Matrix Metallo-Proteinases (because large variety of serious diseases can be correlated to the disfunction of MMPs). One part of the substrate-based inhibitor molecules, developed up to present, consists of a hydroxamic metal-binding group and various  $R_C$  and  $R_N$  substituents. These substituents are planned for selective non-covalent interactions with the sub-sites of the enzyme [11, 12]. However, if any potential donor atom is situating in the substituents, this might modify (often significantly) the metal-binding ability of that hydroxamate-based compound [13, 14]. Evidently, if the mode of the

metal binding is changing, this results in the change of the inhibitory effect too. That is the reason, why our main aim is to clarify the effects of the various donor atoms on the metal (first of all Zn(II), Ni(II) and Cu(II), because these metal ions are often included in the active centers of metalloenzymes) binding properties of hydroxamic derivatives. Since the imidazole is a common donor site in the biological systems, in the present study all molecules involve imidazole moiety at the  $R_C$  substituent. The effects of the distance between the imidazole and hydroxamic functions and also the effects of additional donor(s) situating in different positions compared to the above mentioned two functions on the metal complexation have been investigated by involving a series of imidazole-containing compounds (the formulae of them together with their dissociation constants, taken from ref 15, are listed in the Table 1).

**Table 1:** Formulae of the studied hydroxamic acids and their corresponding dissociation constants ( $pK_a$ )

| <br>(Z) <span style="margin-left: 100px;">(E)</span>   |                                                                                       |                  |       |
|------------------------------------------------------------------------------------------------------------------------------------------|---------------------------------------------------------------------------------------|------------------|-------|
| <i>Completely protonated forms of the ligands (L)</i>                                                                                    |                                                                                       | $R_C$            | $R_N$ |
| Imidazole-4-carboxhydroxamic acid ( $H_2L^+$ )<br><b>(Im-4-Cha)</b><br>$pK_{a1} = 3.76$ ; $pK_{a2} = 8.82$                               |   | -H               |       |
| <i>N</i> -Methyl-imidazole-4-carboxhydroxamic acid ( $H_2L^+$ )<br><b>(N-Me-Im-4-Cha)</b><br>$pK_{a1} = 5.06$ ; $pK_{a2} = 8.57$         |  | -CH <sub>3</sub> |       |
| Imidazole-4-acetohydroxamic acid ( $H_2L^+$ )<br><b>(Im-4-Aha)</b><br>$pK_{a1} = 6.17$ ; $pK_{a2} = 9.05$                                |   | -H               |       |
| Histidinehydroxamic acid ( $H_3L^{2+}$ )<br><b>(Hisha)</b><br>$pK_{a1} = 5.38$ ; $pK_{a2} = 7.11$ ; $pK_{a3} = 9.06$                     |   | -H               |       |
| Alanyl-histidinehydroxamic acid ( $H_3L^{2+}$ )<br><b>(AlaHisha)</b><br>$pK_{a1} = 6.11$ ; $pK_{a2} = 7.72$ ; $pK_{a3} = 8.82$           |   | -H               |       |
| Histidyl-Alaninehydroxamic acid ( $H_3L^{2+}$ )<br><b>(HisAlaha)</b><br>$pK_{a1} = 5.56$ ; $pK_{a2} = 7.55$ ; $pK_{a3} = 9.02$           |   | -H               |       |
| Alanyl-alanyl-histidinehydroxamic acid ( $H_3L^{2+}$ )<br><b>(AlaAlaHisha)</b><br>$pK_{a1} = 6.31$ ; $pK_{a2} = 7.82$ ; $pK_{a3} = 8.84$ |   | -H               |       |



## MATERIALS AND METHODS

### Reagents, metal ion and ligand stock solutions

The ligands are either commercially available (histidine hydroxamic acid, HisHa), or synthesized as previously published [15]. The purity of the ligands and the concentrations of the ligand stock solutions were determined by *Gran's* method. [16] The metal ion stock solutions were prepared from  $\text{CuCl}_2 \cdot 2\text{H}_2\text{O}$  and  $\text{NiCl}_2 \cdot 6\text{H}_2\text{O}$  (Reanal) dissolved in doubly distilled water.  $\text{ZnO}$  (Reanal) was dissolved in a known amount of HCl solution (0.1 M). The concentration of the metal ion stock solutions was determined gravimetrically via precipitation of quinolin-8-olates. The HCl concentration of the  $\text{Zn(II)}$  solution and the exact concentration of the carbonate-free KOH titrant were determined by pH-potentiometry.

### Potentiometric and spectroscopic studies

The pH-potentiometric titrations were performed in the pH range 2.0-10.5 or until precipitation occurred on samples of 4.00 or 5.00  $\text{cm}^3$  at an ionic strength of 0.2 M (KCl) and at  $25.0 \pm 0.1$  °C. During the titrations purified, strictly oxygen-free argon was continuously bubbled through the samples. Samples were always freshly prepared and titrations were completed within 3 hours. The ligand concentrations were varied within the range 0.001-0.0025  $\text{mol}\cdot\text{dm}^{-3}$  and the metal to ligand ratios were varied between the ranges of 1:4 - 1.5:1. The experimental data were collected by titrating minimum 4-5 different samples. The titrations were performed by using a Radiometer pHM84 instrument equipped with a Metrohm combined electrode (type 6.0234.100). Titrant KOH was added from a Metrohm 715 Dosimat auto burette. The electrode system was calibrated according to Irving et al. [17] so that the pH-meter readings could be directly converted into hydrogen ion concentration. The water ionization constant ( $\text{pK}_w$ ) obtained was  $13.76 \pm 0.01$ . The pH-metric results were utilized to find the stoichiometry of species and to calculate the stability constants. The calculations were made with the aid of the computer program PSEQUAD [18]. UV-Vis measurements on systems containing  $\text{Cu(II)}$  and  $\text{Ni(II)}$  were performed. The ligand concentrations were varied within the range 0.0015-0.0025  $\text{mol}\cdot\text{dm}^{-3}$  and the metal to ligand ratios were 1:1, 1:2 and in the case of some  $\text{Cu(II)}$  containing systems were also 1.5:1. A Perkin Elmer Lambda 25 spectrophotometer was used to record the spectra in the region of 300-800 nm. Path length was 1 cm in all cases.

CD spectra for copper(II)-containing systems with all ligands at 1:1, 1:2 and 1.5:1 metal to ligand ratios were also obtained at 25 °C under a constant flow of nitrogen on a Jasco J-810 spectropolarimeter, which was calibrated with an aqueous solution of the ammonium salt of (1R)-(-)-10-camphorsulfonate. The concentrations of the ligands in the samples were 0.001 M in all cases. Measurements were carried out in water at different pH values, using 1 cm path length cuvettes in the 290-800 nm wavelength regions.

The  $^1\text{H}$  NMR measurements were recorded on a Bruker Avance AM 360 by using  $\text{D}_2\text{O}$  as solvent and TSP ((trimethylsilyl)-propionic acid- $\text{D}_4$ , sodium salt) as standard under the following conditions: metal to ligand ratio was 1:1 in all samples and the analytical concentrations were 0.002 M. 0.1 M NaOD and DCl solutions were used to adjust the pD within the range of pD 2 – 11.

Anisotropic X-band EPR spectra of frozen solutions were recorded at 120 K, using a Bruker EMX spectrometer after addition of ethylene glycol to ensure good glass formation. Copper(II) stock solution for EPR measurements were prepared from  $\text{CuSO}_4 \cdot 5\text{H}_2\text{O}$  enriched with  $^{63}\text{Cu}$  to get better resolution of EPR spectra. Metallic

copper (99.3%  $^{63}\text{Cu}$  and 0.7%  $^{65}\text{Cu}$ ) was purchased from JV Isoflex, Moscow, Russia for this purpose and converted into the sulfate. The concentration of Cu(II) was 4 mM in all cases, and the metal to ligand ratios were 1:1 and 1.5:1.

On a few samples, electrospray ionization time-of-flight mass spectrometric (ESI-TOF MS) analysis was carried out using a Bruker Biotof II ESI-TOF instrument. The solutions were introduced directly into the ESI source by a syringe pump (Cole-Parmer Ins. Comp. type 74900) at a flow rate of 2  $\mu\text{L}/\text{min}$ . The temperature of drying gas ( $\text{N}_2$ ) was 100°C. The pressure of the nebulizing gas ( $\text{N}_2$ ) was 30 psi. Voltages applied at the capillary entrance, capillary exit and the first and the second skimmers were -4500, 120, 40 and 30 V, respectively. The spectra were accumulated and recorded by a digitalizer at a sampling rate of 2 GHz. The spectrometer was operated at unit mass resolution and was calibrated to sodium trifluoroacetate.

## RESULTS AND DISCUSSIONS

### Imidazole analogues of $\alpha$ - and $\beta$ -amino hydroxamic acids (Im-4-Cha, N-Me-Im-4-Cha, Im-4-Aha)

The relatively low basicity imidazole-N(3) was found to behave as anchor donor not only to  $\text{Cu}^{2+}$  and  $\text{Ni}^{2+}$ , but even to  $\text{Zn}^{2+}$ , although this latter metal ion prefers oxygen donors over nitrogen ones. By increasing the pH water insoluble polynuclear species have been found to form with Im-4-Cha. A pentanuclear metallacrown (Scheme 1) is exclusively formed above pH 4 between Cu(II) and Im-4-Aha. The stability of this species with Ni(II) and Zn(II) is significantly lower, and as a consequence, it appears together with various mononuclear bonding isomers in the Ni(II)-Im-4-Aha and Zn(II)-Im-4-Aha systems.




Fig. 1

Beside the mononuclear species a dinuclear complex,  $[\text{M}_2\text{L}_3]$  (Scheme 2), was found in the Cu(II)- and Ni(II)-N-Me-Im-4-Cha. A possible bonding mode is that one of the three ligands bridges two metal ions, while each of the other two ligands binds to one metal. Stacking interaction between the imidazole rings of the latter two ligands might stabilize this species.




Fig. 2

Based on the results, this molecule, which is capable for the interaction with two metals, might be effective inhibitor of a metalloenzyme involving two divalent metal ions in the catalytic unit.

### **Hydroxamic derivative of histidine (Hisha)**

In this molecule the  $R_C$  involves an imidazole in  $\gamma$ -position to the hydroxamic and to  $\beta$ -position to the terminal amino group. As it is known from previous results [1], the imidazole takes part in the coordination first of all in the complexes with Cu(II), but also with Ni(II), where octahedral complexes with tridentate coordination of the ligand ( $N_{\text{amino}}, N_{\text{imidazole}}, N_{\text{hydroxamate}}^-$ ) dominate below pH 7. Above this pH, however, the hydroxamate-coordinated complexes become dominant. With Zn(II) ion, the coordination starts similarly to the simple  $\alpha$ -amino hydroxamic acid derivatives, but NMR results indicate also imidazole coordination.

### **Hydroxamic derivatives of histidine-containing di- and tripeptides (AlaHisha, HisAlaha, AlaAlaHisha)**

The lower basicity of the imidazole, compared to that of the hydroxamate gives a benefit to the former donor especially at lower pH in the studied peptide derivatives and the role of the hydroxamate generally increases upon increasing the pH. It is also a common feature that, upon increasing the pH, when the imidazole-N is displaced by the hydroxamate in the coordination sphere, the former moiety does not remain uncoordinated, but imidazole-bridged di- and polynuclear complexes are formed. In contrast with this, in the Cu(II)-AlaAlaHisha system, the hydroxamate can take part in the coordination only at lower pH, where the hydroxamate chelate can successfully compete with the amino – imidazole macrochelate, but can not compete with the very high stability 4N-coordinated (amino-, amide-, amide- and imidazole-N donors) bonding mode. As a consequence, in this system, the hydroxamate does not play measurable role in the coordination at high pH. With Ni(II), the formation of planar complexes is generally more favoured with hydroxamic derivatives of the histidine-containing di- and tripeptides compared to molecules containing carboxylic instead of hydroxamic moiety. This demonstrates the exceptional coordinating ability of the imidazole ring. Any indication for deprotonation and coordination of pyrrole-nitrogen was not found in the systems. With Zn(II), formation of various bonding isomers with rather low stability was found. The complex formation does not start below pH ca. 6 and precipitation occurs at pH. 7.5-9 depending on the ligand and Zn(II) to ligand ratio.

## **CONCLUSIONS**

Increased selectivity of the hydroxamate-based metalloenzyme inhibitors can be expected, if, in addition to the interaction of the hydroxamate group with the catalytic metal, non-covalent interactions also occur between the substituents of the inhibitor molecule,  $R_C$  and  $R_N$ , and the enzyme. To achieve this, favoured coordination of the hydroxamate chelate over the imidazole to the metal ion is necessary. Because of the above mentioned facts, the role of  $R_C$  substituents, involving imidazole moiety, in Cu(II)-, Ni(II)- and Zn(II)-complexation of various hydroxamic acids (derivatives of amino acids and small peptides) has been investigated. A competition between the hydroxamate and imidazole functions was found in all studied systems, but the extent of that competition was found to differ from metal to metal, from ligand to ligand and to depend very much on the pH. The

imidazole played the most determinant role in the Cu(II) complexes, somewhat less in the Ni(II)-containing ones, while the most negligible role occurred in the Zn(II)-complexes.

**Acknowledgements:** This work was supported by the Hungarian Scientific Research Funds (OTKA T 046366 and T 049612).

## REFERENCES

1. B. Kurzak, H. Kozłowski, E. Farkas, *Coord. Chem. Rev.* 114 (1992) 169
2. H. Sigel, R. B. Martin, *Chem. Rev.* 82 (1982) 385
3. I. Sóvágó, K. Ősz, J. Chem. Soc. Dalton. Trans., (2006) 3841
4. V. Józai, Z. Nagy, K. Ősz, D. Sanna, G. Di Natale, D. La Mendola, G. Pappalardo, E. Rizzarelli, I. Sóvágó, *J. Inorg. Biochem.* 100 (2006) 1399
5. Cs. Kállay, K. Ősz, A. Dávid, Z. Valastyán, G. Malandrinos, N. Hadjiliadis, Sóvágó, *Dalton Trans.* (2007) 404
6. I. Török, T. Gajda, B. Gyurcsik, G. K. Tóth, A. Péter, *J. Chem. Soc. Dalton Trans* (1998) 1205
7. H. Kozłowski, W. Bal, M. Dyba, T. Kowalik-Jankowska *Coord. Chem. Rev.* 184 (1999) 319
8. A-M. Albrecht-Gary, A. L. Crumbliss, in *Metal Ions in Biological Systems*, vol. 35 (Eds.: A. Sigel, H. Sigel), Marcel Dekker, New York, 1998
9. E. M. F. Muri, M. J. Nieto, R. D. Sindelar, J. S. Williamson, *Current Med. Chem.* 9 (2002) 1631
10. C. J. Marmion, D. Griffith, K. B. Nolan, *Eur. J. Inorg. Chem.* (2004) 3003
11. D. T. Puerta, S. M. Cohen, *Current Topics in Med. Chem.* 4 (2004) 1551
12. M. Jani, H. Tordai, M. Trexler, L. Bányai, L. Patthy, *Biochimie* 87 (2005) 385
13. E. Farkas, I. Kiss, *J. Chem. Soc. Dalton Trans*, (1990) 749
14. P. Buglyó, E. M. Nagy, E. Farkas, I. Sóvágó, D. Sanna, G. Micera, *Polyhedron* 26 (2007) 1625
15. E. Farkas, D. Bátka, E. Csapó, P. Buglyó, W. Haase and D. Sanna, *Polyhedron*, 26, (2007), 543;
16. E. Farkas, E. Csapó, P. Buglyó, C.A. Damante, G. Di Natale, *Inorg. Chim. Acta* (submitted)
17. Gran, *Acta Chem. Scand.*, 4 (1950) 559
18. H.M. Irving, M.G. Miles, L.D. Pettit, *Anal. Chim. Acta.*, 38 (1967) 475
19. L. Zékány, I. Nagypál, in *Computational Methods for the Determination of Stability Constants*, ed. D.L. Leggett, Plenum Press, New York, 1985, p. 291.

# INTEGRATED MODELING OF GEOCHEMICAL AND BIOGEOCHEMICAL PROCESSES ACROSS SPACE-TIME SCALES

Radu LĂCĂTUȘU<sup>1\*</sup>, Virgil IORDACHE<sup>2</sup>

1.\* Research Institute for Soil Science and Agrochemistry, RO-011 464 Bucharest, Roumania;

2 Research Centre for Ecological Services – CESEC, University of Bucharest, RO-050 095 Bucharest, Roumania

\* E-mail: virgil.iordache@g.unibuc.ro

## ABSTRACT

*In order to find out how the environment is organized one investigates structural elements and processes. Geochemical, soil and biogeochemical processes are amongst the most frequently used for characterizing the so called “hierarchical levels” of the environment’s structure. The goal of the project we present in this paper is to test in five catchments hypotheses 1: the budgets of microelements assessed in large catchments don’t close with the budgets of their sub-catchments and hypotheses 2: the bioaccumulation in mobile organisms (top consumers) cannot be predicted base on concentrations in organisms which are their trophic base, on their local abundances, and on the trophic preferences of the top consumers. The activities are organized under three objectives: one dedicate to the ecosystem level mechanisms supporting the processes at landscape level, one for testing the first hypotheses, and one for testing the second hypotheses. The activities are grouped in seven workpackages (wps): in objective 1 a wp for investigating the role of microorganisms (with a sub-wp for symbioses with plants), and wp for organo-mineral complexes, with a sub-wp dedicated to chemical speciation of metals, in objective 2 a wp for fluxes of metals by groundwater, and one for runoff and erosion (with a sub-wp for sediment transport in rivers), including math modeling, in objective 3: a wp for bioaccumulation and ecotoxicology in terrestrial systems (with sub-wps for plants, carabids, frogs and small mammals) and a wp for bioaccumulation and ecotoxicology in aquatic lotic systems. The seventh wp supports all objectives and is dedicated to math modeling with four sub wps: gis database, stochastic modeling, cellular automata modeling and neural networks modeling*

**Key words:** biogeochemistry, soil science, ecology, geology, hydrology, metals, mathematical modeling

## INTRODUCTION

In this paper we present a research project for complex exploratory research (acronym BIOGEOST) submitted under the call 2008, which passed the first selection phase and now is in the final evaluation phase. Whatever the result of this evaluation will be, we believe that the ideas and the approach specific to this project are valuable for discussion in the scientific community.

Consequently the structure of the paper will be informal, following the structure of the project. We have not presented the very technical part of the project, which is not of general interest. The paper can be seen also as a short review in the area of trace elements biogeochemistry.

---

„Metal Elements in Environment, Medicine and Biology”, Tome VIII, pp.41-54, Publishing House “Eurobit” Timișoara, 2008

Proc. of 8<sup>th</sup> Int. Symp. of Roumanian Academy - Branch Cluj-Napoca and Branch Timișoara, 2008, December 5-6, Timișoara, Roumania

## 1. IMPORTANCE AND RELEVANCE OF SCIENTIFIC CONTENT

After the failure of the heavily subsidized attempts to do produce holistic models of ecological systems in the 1970' in the frame of the International Biological Program, the financing of such research program went down to reasonable dimensions, despite the widespread recognition of the existence of a hierarchy of ecological systems, from local to global level, whose functional modelling is a desirable research objective (Loijen, 1998). The idea to do such costly modelling revived recently under the heading of "modelling the socio-ecological systems", but still the possibility of successful implementation of this research agenda is debated: the holistic approaches have failed in the past, and the reductionistic ones are said to not reflect the true characteristics of the systems. At this general level, the idea of this project is to implement a new strategy for modelling hierarchically organized ecological systems, coupling holistic and mechanistic approaches.

The complex systems such as the ecological ones are characterised by and intrinsic principle of incertitude (Jorgensen and Svirezhev, 2004; Stanciulescu, 2005). Because there are no precise solutions for complexly organized systems, there is no single method for their study. It isn't quite "anything goes", but we cannot tell what might work without trying it. Taking into consideration also the non-linearity of the complex systems and their real, non-stochastic, indeterminacy (due to biological evolution, for instance), one can say, from a philosophy of science standpoint, that there cannot be a final mathematical model of such systems: models of the functioning of ecosystems will evolve (in an evolutionary epistemology sense). Our strategy is to not look for a single comprehensive model, whose epistemic "mutations" and evolution are very costly, and consequently difficult to accept by the scientific community, but to produce a maximally coherent portfolio of models with a modular structure, most appropriate for a high rate of epistemic evolution occurring by piecemeal changes at the level of the modules and of the relations between them. By now the models belonging to the potential "modules" have evolved independently, disciplinary, without explicit concern about their relationships with other disciplinary developed models.

To give substance to this strategy, the project deals with modelling biogeochemical cycling of metals occurring in hierarchically organized ecological systems. That this is a basic research priority is proved by a comprehensive analyses of the bioaccumulation models coupled with ecotoxicological models done in the frame of a previous ESF project. In order to do describe the metals' circulation there is a need to develop food-web models (modifying and extending their structure), especially models describing natural, complete terrestrial ecosystems ([www.esf.org/ecolmat](http://www.esf.org/ecolmat), 2003). Starting from the conclusions of this ESF project we performed a review of the state of the art concerning the mathematical modelling of the cycling and effect of metals at landscape level (Iordache and Stelian, 2007). The general conclusions were:

1. Current models applicable at landscape scale describe mainly the abiotic side of the processes, the best example being reactive transport models. However, there are many opportunities for interdisciplinary transfer of ideas profitable for the field of mathematical modelling in biogeochemistry and ecotoxicology.
2. The limits related to modelling the involvement of the organisms cannot be surpassed at the current state of biological knowledge, despite the efforts in the systems biology field.
3. With respect to the development of the scientific knowledge in biogeochemistry and ecotoxicology the potential of modelling is underused, because the modelling efforts are focused mainly on the side rapidly relevant for socio-economic systems.
4. There is an important opportunity to use populations of similar contaminated and, eventually, bioremediated sites (seen as in situ experimental areas) for testing biogeochemistry hypotheses

with modelling tools (i.e. to build some kind of “long term *contaminated* ecological research” network).

The difficulty which blocked the refinement of this kind of models for *in situ* cases is related to the large heterogeneity of the control parameters, the distribution in space, as well as to the large number of relationships between the control parameters for each metal. In these conditions, to set up mathematical relations between the control parameters of metals mobility and their mobility proved to be not yet feasible *in situ*. It may be that the development of new mathematical tools, more adapted to biological phenomena, would contribute to changing the landscape of models in this area. For instance, Schryver et al. (2006) use non-linear analyses with artificial neural networks with good results to describe the relationships between groundwater contamination with metals and the structure of bacterial communities as measured by molecular markers. Beside neural networks, another new tool is the Multivariate Adaptative Regression Splines (MARS). It has been used for instance for predicting the denitrification rate based on data sets at European scale (Pinay et al, 2007).

These mathematical tools could be applied in order to describe relationships between parameters describing metals biogeochemistry at the *explanandum* (higher hierarchical) level, on the one side, and its control parameters at the *explanans* (lower hierarchical) level. Once the analyses are performed, one would have a holistic phenomenological description of the relationships in mathematical forms at site scale. One limitation of this approach is to uncover mathematical laws with 1 member of the reference class (the studied site). This limitation can be surpassed only by studying a population of sites across gradients. Another limitation of the holistic approach is that the discovered mathematical laws will give no indication about the mechanisms supporting them. This indication can be provided only by a mechanistic (bottom-up) approach. By now reductionism has created unwarranted biases in certain mathematical models and experimental designs. Mechanistic models may incorporate unnecessary lower level details that compromise verisimilitude and predictive success. However, without this approach one cannot have an image on the involved mechanisms, and thus on the possibilities of controlling the metals' mobility. Therefore one can proceed this way, having in mind that the mechanistic frame has to be re-verified after each step of upgrading and prediction. Our idea is to develop (bottom-up) mechanistic models of metals' behaviour under the (top-down) constraint of the results of the holistic modelling (figure 1).




**Fig 1.** Overall structure of the modelling approach.

The main focus from the mechanistic perspective will be on the difficulty of

allocating values to the parameters under high field heterogeneity. The heterogeneity of the real systems at a certain hierarchical level cannot be predicted based on principles (Dungan et al., 2002). The heterogeneity problem can be dealt with by empirical research for delineating in situ the elementary units of models application (systems identification) and using a portfolio of models covering the diversity of the elementary units. Different processes require different space-time scale of the elementary units and models' development. Programming in GIS can then be used for upscaling the models' results from the elementary units of application to the site and for linking the results of the models with different space-time scale (Iordache and Bodescu, 2005). The upscaling problem is linked to some extent to the reduction of the ontological levels problem, and can be solved partially by recognizing the irreducibility of ecological levels of organization, such as elementary ecosystems and landscapes (contaminated lands are usually at these levels). Then each level can be approached by specific model(s). The remaining part of the upscaling problem, within a certain level, directly reflects the heterogeneity problem. In this context, the general objectives of a research program in this area can be:

1. To uncover mathematical laws describing the mobility and effects of metals at contaminated landscape level.
2. To develop coupled top-down and bottom-up models of key processes of metals biogeochemistry and effects in a population of contaminated landscapes.

This project should be seen as one step in such a long-term research program.

## **2. RESEARCH METHODOLOGY (AIM, OBJECTIVES, AND WAYS OF REACHING THEME)**

The goal of the project is to test the following hypotheses:

- Hypotheses 1: The budgets of microelements assessed in large catchments don't close with the budgets of their sub-catchments. Rationale: the boundary conditions of the sub-catchments have their own dynamic, specific to the functioning of the system of larger scale (for instance the change in the landscape structure can be unpredictable at small scale, but predictable at larger scale) 2) buffering and remobilization areas can occur on the flow path of underground and surface waters (so called hot spots and hot moments). Testing method: coupling models for the budgets in small catchments and comparing the results with the results of models developed for the integrating catchment. Math modeling of the microelements budget will be performed as integrated modeling (separate sub-models for the involved mechanisms coupled in between).
- Hypotheses 2: The bioaccumulation in mobile organisms (top consumers) cannot be predicted base on concentrations in organisms which are their trophic base, on their local abundances, and on the trophic preferences of the top consumers. Rationale: The emergent properties at large scale would interfere with the simple accumulation in trophic chains by the way it controls the availability of trophic resources at large scale and indirectly the feeding behaviour of mobile organisms. Testing method: the performances of deterministic models of bioaccumulation (cellular automata type) will be compared with neural networks based models to predict the accumulation in plants, carabids, rodents and birds. We expect that the differences in predictions will become larger as the mobility of the organisms increase, and that the neural networks based models will be more accurate in the validation phase.

These two hypotheses have an attractor role for the whole project.



Complementarily, in each WP the scientific groups involved will test specific, local hypotheses. In this way the data sets obtained will be useful both for the disciplines involved in the project, and for the systemic biogeochemical approach providing the identity of the project.

The experimental and theoretical activities, having as final end to produce new elements of knowledge and developing principles and concepts, will be organized, from a management perspective under three separate objectives dedicated to 1) Biogeochemical mechanisms at ecosystem level 2) Testing the hypotheses concerning the budget of trace elements at landscape scale, 3) Testing the hypotheses about trophic-dynamic modules emergent at landscape scale.

The activities are grouped in seven workpackages (wps): a wp for investigating the role of microorganisms (with a sub-wp for symbioses with plants), and wp for organo-mineral complexes, with a sub-wp dedicated to chemical speciation of metals, a wp for fluxes of metals by groundwater, and one for runoff and erosion (with a sub-wp for sediment transport in rivers), including math modeling, a wp for bioaccumulation and ecotoxicology in terrestrial systems (with sub-wps for plants, carabids, frogs and small mammals) and a wp for bioaccumulation and ecotoxicology in aquatic lotic systems; the seventh wp supports all objectives and is dedicated to math modeling with four sub wps: gis database, stochastic modeling, cellular automata modeling and neural networks modeling.

Figure 2 presents the Pert diagram of the project and the relations between WPs and objectives. Table 1 presents the Gant diagram of the project.




**Fig 2.** Pert diagram of the project and the relationships between workpackages and objectives.

**Table 1.** Gant diagram of the project.

| Objective                                                                            | Workpackage                                       | 2008 | 2009 | 2010 | 2011 |
|--------------------------------------------------------------------------------------|---------------------------------------------------|------|------|------|------|
| To characterize biogeochemical mechanisms at ecosystem level                         | WP1 Organic-mineral complexes                     |      |      |      |      |
|                                                                                      | WP1.1 Metals speciation                           |      |      |      |      |
|                                                                                      | WP2 Role of microorganisms                        |      |      |      |      |
|                                                                                      | WP2.1 Mechanisms of symbioses                     |      |      |      |      |
|                                                                                      | WP7.1 Data base                                   |      |      |      |      |
| To test the hypotheses about the budget of trace elements at catchment level         | WP7.2-4 Math modelling Ob1                        |      |      |      |      |
|                                                                                      | WP4 Fluxes mediated by groundwater                |      |      |      |      |
|                                                                                      | WP5 Fluxes mediated by runoff and erosion         |      |      |      |      |
|                                                                                      | WP5.1 Sediment transport in rivers                |      |      |      |      |
|                                                                                      | WP7.1 Data base                                   |      |      |      |      |
| To test the hypotheses about the emergent trophic-dynamic modules at landscape scale | WP7.2-4 Math modelling Ob2                        |      |      |      |      |
|                                                                                      | WP6 Bioaccumulation and terrestrial ecotoxicology |      |      |      |      |
|                                                                                      | WP6.1 Plante                                      |      |      |      |      |
|                                                                                      | WP6.2 Carabidae                                   |      |      |      |      |
|                                                                                      | WP6.3 Amfibieni si mamifere mici                  |      |      |      |      |
| Scientific coordination, management                                                  | WP3 Bioaccumulation and aquatic ecotoxicology     |      |      |      |      |
|                                                                                      | WP7.1 Data base                                   |      |      |      |      |
|                                                                                      | WP7.2-4 Math modelling Ob3                        |      |      |      |      |
|                                                                                      | Tasks distribution and monitoring                 |      |      |      |      |

The field sites include five catchments with contaminated lands, spread over the climatic gradients of Romania (E-W, N-S, table 2).

**Table 2.** Catchments studied by the partners. Acronyms explained in figure 5.

| Studied catchments         | UAIC-Ggrf | UAIC-Geol | UBB-Geo | UBB-Bio | UB-Eco | UB-Ghydro | UB-Gch | ICPA | USAM VB | UBM |
|----------------------------|-----------|-----------|---------|---------|--------|-----------|--------|------|---------|-----|
| <i>Săsar</i>               | x         | x         | x       | x       | x      | x         | x      | x    | x       | x   |
| <i>Abrud-InferiorArieș</i> | x         | x         | x       | x       | x      | x         | x      | x    | x       |     |
| <i>Ampoi-Geoagiu</i>       | x         |           | x       | x       | x      | x         | x      | x    | x       |     |
| <i>LeșuUrsului</i>         | x         | x         |         | x       | x      | x         | x      | x    | x       |     |
| <i>Pantelimon</i>          | x         |           |         | x       | x      | x         | x      | x    | x       |     |

**Table 3.** Types of approaches in the studied systems.

| Type of system            | Type of approach |           |                          |                 |
|---------------------------|------------------|-----------|--------------------------|-----------------|
|                           | Field study      |           | Experimental             | Math simulation |
|                           | extensive        | intensive |                          |                 |
| Low order catchment       | x                |           |                          | x               |
| Highly contaminated lands |                  | x         | micro, mezo, field scale | x               |

Each catchment will be approached by several research methods at three scales of analyses (table 3): contaminated ecosystem level, small order catchment, and medium order catchment.

For the symbiotic part in WP2 we will use ectomycorrhiza (EM) as study model (figure 2), hypothesizing a peculiar organization of the trophic-dynamic modules of this group. From this perspective we are able to identify the following components of EM structural diversity in a tree island: number of TDMs, species richness inside each TDM ( $\alpha$ ), at tree level ( $\beta$ ) and forest level ( $\gamma$ ), evenness inside each TDM.




**Fig. 3.** Model of EMs community structure in a forest.

This way of conceptualizing the structural diversity allows a direct connection with functional interpretation. For instance, changes in microbial diversity reported in the literature did not always correspond to changes in functional redundancy. The reason for this is that diversity is usually characterized unstructured, at tree island level, which mixes the diversity of different TDMs. As functional redundancy of species occurs only at infra-TDM level, an increase of overall diversity (across pooled TDMs) does not necessarily imply an overall increase in functional redundancy. One can have for instance decrease of redundancy in a TDM coupled with an increase in another TDM, or simply the appearance of new TDMs in the system's structure. The functional diversity, excepting for the redundancy component, will change only with the change in the number of TDMs. The research done will allow us to test this concept model.

The main objective of the groundwater workpackage is a transient coupled model at different scale of space and time for geochemical and biological processes as. Five experimental sites (Sasar, Abrud – Inf. Aries, Ampoi- Geoagiu, Leșul Ursului, Pantelimon) have to be characterized with respect to flow paths, and in subsequent years the waters sampled from soil, surface and ground water wells. The result of monitoring for formation of groundwater, chemistry of soil and runoff water will be used in fitting operation of the transient coupled model. The support of coupled models will be the spatial distribution of metals (WP1) and an important factor of stability of the environmental conditions is the sediment transport in the rivers (WP5) and the elements of aquatic ecotoxicology (WP3). In the final stage the fitted transient coupled model will be connected with sthochastic models for bioaccumulation and terrestrial ecotoxicology (WP6).

In workpackage 6 we will provide the population level data needed for developing bioaccumulation models for four model organisms characterized by different space-time scale of activity. A comprehensive analyses of the bioaccumulation models coupled with ecotoxicological models was done in the frame of an ESF project ([www.esf.org/ecolmat](http://www.esf.org/ecolmat), 2003). They state that ecosystem modelling and, more specifically, modeling of food-web processes, provides a proper vehicle to study the ecotoxicological impacts and to make the risk assessment. They conclude that an important obstacle to the use of food-web models in ecotoxicological risk assessment is the lack of hard data, or more precisely a database containing these data. This includes ecophysiological and life history parameters (data on consumption, mortality and the like, as well as data on concentration-effect

relationships for those parameters) for the various functional groups in an ecosystem. Supplementary research in order to use food-web models for ecotoxicological risk assessment involves the further development of food-web models (modifying and extending their structure), especially models describing natural, complete terrestrial ecosystems. It also involves collecting the required input data and parameter values either in the laboratory or in the field. Furthermore, a comprehensive set of field data is required for validation purposes ([www.esf.org/ecolmat](http://www.esf.org/ecolmat), 2003). Andretta et al. (2006) discuss the most important theoretical aspects of polluted soil Risk Assessment Methodologies, which have been developed in order to evaluate the risk, for the exposed people, connected with the residual contaminant concentration in polluted soil, and make a short presentation of the major different kinds of risk assessment methodologies. They also underline the relevant role played, in this kind of analysis, by the pollutant transport models and describe a new and innovative model, based on the general framework of the so-called Cellular Automata (CA), initially developed in the UE-Esprit Project COLOMBO for the simulation of bioremediation processes. This line of research will be followed in the last protocol, modeling.

The data used for modeling (WP7) will have three sources: comprehensive critical analyses of the (experimental and field) literature, field investigations and experimental activity.

Figure 4 shows the relationships between the models developed based on experimental data and those developed based on field data. As suggested in figure 4, the dataset produced as a result of the experiments will be characterized by a higher number of parameters, but a lower variability of parameters values, while the dataset produced from the field study will consist in a lower number of parameters but with much higher variability of values due to the natural space-time heterogeneity. These data sets with different characteristics will be used for the production of mathematical models following the scheme pictured in figure 4. In order to set up the nonlinear correlation model we will use the software MATLAB with the needed modules, and the software MARS.

The models developed based on the experimental data will be extrapolated at field scale by a soft programmed in GIS and using maps of the basic control parameters distribution and eventually modular submodels of derived control parameters such as the redox potential in soil. The results obtained after the extrapolation will allow us to check the nonlinear correlation obtained from field data, and, to the extent that the fit between the two approaches will be good, to point out some of the mechanisms underlying the field correlations.

A variety of other inter-comparisons between the models will be performed, as described in figure 4. The overall result will be a coherentised portfolio of models used finally for the simulation of metals mobility in different ecosystems types and degrees of metals contamination, at different scales.




**Fig. 4.** Relationship between spatially explicit and un explicit models in this project.

### 3. NOVELTY AND ORIGINALITY DEGREE

After the accumulation of knowledge concerning local and global biogeochemical processes, in the last 10-15 year the international interest moved towards the study of systems of intermediary scale, between local and global (as demonstrated by the increased number of article in ISI journals). The applicative stake is huge, being linked to understanding how local actions are connected with global effects. The researches lead to the proposal of concepts like hot spots (HS) and hot moments (HM), meaning zones in space and periods of time in which the rates of the geochemical and biogeochemical processes are much larger than in other zones and areas. A question not yet answered is at what scale should these concepts be applied. From an applied perspective the scale is imposed by the management problem to be solved, and then the question vanishes (actually this is the answer of the socio-ecological approach). From a naturalistic perspective, adopted in this project, the question can be reformulated as follows: are there objectively certain space-time scales of analyses at which HS and HM can be identified, or rather HS and HM can be identified at any scale of analyses ? O positive answer to the first question would support the theory that the hierarchical theory of the environment (according to which there is a number of hierarchical level and each level is characterized by certain emergent properties which can be identified; the presence of HS and HM could be such emergent properties). If the answer to the second question would be positive, then the notion of hierarchical level would become inappropriate and we should re-conceptualize the relation between the scale of analyses and the occurrence of emergent properties (eventually by using

and quantitative indicator for the appearance of emergent properties as the space-time scale of analyses increases). In this context, the project contributes to testing the general hypotheses that there is emergency of biogeochemical properties at certain space-time scales, not reducible to the functioning of the systems identifiable at smaller scale. In particular, the hypothesis is that HS and HM organizes relatively independently at certain space-time scales. In systems analyses terms, the hypothesis implies that large scale systems are not fully decomposable in subsystems, that there is not a homomorphic model able to describe all its relevant properties. From the perspective of math modeling the hypothesis implies that one cannot build models fully reducing the functioning of large scale systems to that of small scale ones, and then one should develop independent models at least in what concerns the some properties of the system. These consequences would be imposed by the nature of the environment, and not by pragmatic constraints (which currently lead to a *de facto* independent approach of the geochemical and biogeochemical modeling of systems characterized by different scales). One could then ask: which are those ST scales at which the properties we are interested in emerge? An next: do these scales vary in space and time ? The project, beyond its direct contribution to geochemistry and biogeochemistry, contributes to the international efforts to answer such theoretical questions. Applicative importance: Soil Directive, WFD, and ecological risk assessment, by objectives.

#### **4. INTERDISCIPLINARITY AND/OR TRANSDISCIPLINARITY**

We will provide arguments for the fundamental character of the project, and next we point out its inter- and trans-disciplinarity. Then the epistemic legitimacy of the approach proposed in this project as compared with other approaches in the research field is discussed.

##### ***4.1. The fundamental character***

We provide two kinds of arguments for the fundamental character of the research proposed: by the results of the critical analyses of the existing scientific knowledge, and by referring external authorities (National Science Foundation, European Research Council, European Science Foundation, and Directorate General XII – FP7).

##### ***4.1.1. Arguments by the critical analyses of the knowledge***

In order to find out how the environment is organized one investigates structural elements and processes. Geochemical, soil and biogeochemical processes are amongst the most frequently used for characterizing the so called “hierarchical levels” of the environment’s structure. Implicitly, geochemical and biogeochemical parameters are conceived to emerge only at certain levels of environment’s organization. On the other hand, obtaining general as general as possible knowledge (eventually applicable in all systems of a certain hierarchical level) requires the study of a statistical population of similar sites, which is a difficult task in the case of large systems like catchments. Such research projects are relatively rare (for instance those from the Hubbard Brook experimental area, <http://www.hubbardbrook.org/>), and have concentrated mostly on macroelements. The idea to intentionally contaminate ecosystem was considered un-ethical (despite the fact that the method was applied in the Experimental Lakes Area, [http://www.dfo-mpo.gc.ca/regions/central/science/enviro/ela-rle\\_e.htm](http://www.dfo-mpo.gc.ca/regions/central/science/enviro/ela-rle_e.htm)). Alternatively, recently aroused and consolidated the idea to use sites already contaminated for fundamental research of geochemical and biogeochemical processes (Kozlov and

Zvereva, 2007). Their advantage for fundamental research is the high number and the low socio-economic interest in using these sites for other purposes. As Romania's potential in this respect is, unfortunately, large, the institutions of our consortium oriented their research in this direction and several projects of smaller dimension. Two members of the project team performed a comprehensive critical analyses of literature about mathematical modelling of the biogeochemical cycles of microelements (Iordache and Stelian, 2008), and its results together with the experience accumulated in previous projects will be used in this proposal.

#### *4.1.2. Argument by the authority of prestigious institutions*

In the last decade the National Science Foundation had an integrated program dedicated to geochemical and biogeochemical research (<http://www.nsf.gov/pubs/1999/nsf999/nsf999.htm>), which is currently continued by the U. S. Department of Energy Office of Science in direct correlation with contaminated areas ([http://www.lbl.gov/ERSP/generalinfo/geochem\\_biogeo.html](http://www.lbl.gov/ERSP/generalinfo/geochem_biogeo.html)). At the level of European Science Foundation the biogeochemistry of trace elements was promoted in the context of polluted systems ([http://www.esf.org/fileadmin/be\\_user/research\\_areas/LESC/Documents/GPollGoteborg.pdf](http://www.esf.org/fileadmin/be_user/research_areas/LESC/Documents/GPollGoteborg.pdf)), of the use of lanthanides for inferring some characteristics of the biogeochemical cycles, of the limiting factors for the development of organisms, and of the modeling of bioaccumulation in food chains ([www.esf.org/ecolmat](http://www.esf.org/ecolmat)). The European Research Council recently funded (by FP7) 8 Ideas starting grants in area PE8 (Earth system sciences), and 13 are on the reserve list. The FP6 program financed in the area of our proposal an ERA-NET project (SNOW-MAN, <http://www.snowman-era.net/index.php>). Members of our consortium are in course of involvement in this ERA-NET's research project with a la carte funding, and they are also contributing to the scientific aspects of the extension of group of funding agencies with a Romanian one. A priority of FP7 is "Soil processes modeling".

#### **4.2. Interdisciplinarity and transdisciplinarity of the proposed fundamental research**

The interdisciplinary character is obviously underlined by the name of the scientific field (geo-chemistry, bio-geo-chemistry). As for the transdisciplinarity, one can envisage by this term either the fact that the project supports the creation of a new discipline, or that the project supports the creation of a new way of science transcending the disciplinary approach. In the first interpretation, one can argue that biogeochemistry is a new emerging discipline, as suggested by the increasing number of journals dedicated to the field. If one envisage a transdisciplinary character in the second sense, one could argue (from the perspective of the philosophy of science – Stăicu and Iordache, 2007) that transdisciplinary approaches by intention function as cultural mutations, and will be subject to selective pressure. Consequently one needs as many such mutations as possible, in order to ensure the selection of the most effective one. The integration of all scientific efforts in a single intentionally transdisciplinary approach (e.g. of the „socio-ecological systems” type) is ineffective from an evolutionary perspective, despite its efficiency on the short term by centralizing the resources.

### **5. CONSORTIUM FUNCTIONING SCHEME**

The places of the partners of this project in the Consortium for the biogeochemistry of trace elements and the complementarities with major



infrastructure projects are presented in figure 5. The management structure of the project is depicted in figure 6. The steering committee of each WP (SCWP) will include on representative from each research unit with expertise in the WP area. Each SCWP will have on WP leader and will send two delegates to the Coordination committee (CC) of the project in such a way to ensure equilibrium of organizations in its structure.

## 6. MEASURES FOR FULFILLING THE DEONTOLOGICAL NORMS/STANDARDS OF RESEARCH

The large number of feedback loops and the external connection in the structure of the project management (figure 6) will allow 1) the internal control of the quality of results, 2) the external control of the quality of results 3) the early mitigation of any disputes between the institutions or member, including those dealing with ethical aspects. On could remark also the democratic organization, but with clear responsibilities, of the management structure.




**Fig.5.** Current structure (open to other potential partners) of the national consortium for the biogeochemistry and ecotoxicology of trace elements.

With orange are indicated the partners involved in the BIOGEOST project. The institutions on pink are holders of the infrastructure for trace elements biogeochemistry purchased in the running INFRABIM Capacities project. The institution on yellow is holder of the infrastructure for metals in food chain purchased in the running HEVMETFOOD project. *Legend:* Biotechnol = research center externalized from USAMV Bucuresti, FLPR= National Institute for the Physics of lasers, plasma and radiations, IBB=Biology Institute of Bucharest, Romanian Academy, ICPA = National Institute for Pedology and Agrochemistry, IMNR= National Institute for rare and non-ferrous metals, ISMMA = National Institute for statistics and applied mathematics, Romanian Academy, ROSA = Romanian Space Agency, UPB= University Politehnica of Bucharest.



## CONCLUSIONS

We developed a large integrated project in order to catalyze the cooperation between soil scientist, geologists, hydrologist, biologists and ecologists in the frame of the interdisciplinary field of trace elements biogeochemistry. We believe that the above elements and approach can function as an attractor also for other Romanian institutions interested in interdisciplinary environmental research. In particular, for operational reasons we have not included research directions dealing with human and animal health in the project, but this is a necessity and could be done in complementary projects. The value of this approach is confirmed at European level by the success of similarly organized project including Romanian partners ([www.umbrella.uni-jena.de](http://www.umbrella.uni-jena.de)).




**Fig. 6.** Structure of BIOGEOST project  
(adapted from FP6 IP ALARM, Settele, J., et al., 2005, GAIA, 14, 69-72).

## REFERENCES

1. Dungan J. L., Perry J.N., Dale M.R.T., Legendre P., Citron-Pousty S., Fortin M.-J., Jakomulska A., Miriti M., Rosenberg M.S.- A balanced view of scale in spatial statistical analysis, *Ecography*, 2002, 25, 626-640

2. Iordache V., Bodescu F.- Emergent properties of the Lower Danube River System: consequences for the integrated monitoring system, Arch. Fur Hydrobiologie. 2005, Suppl. 158, (Large Rivers 16), 95-128
3. Iordache V., Stelian I., Bodescu F., Pohoata A.- Mathematical modeling in bioremediation: potential and limitations, Chemie der Erde, 2009, accepted for publication
4. Jorgensen S.E., Svirezhev Y.M.- Towards a thermodynamic theory of ecological systems, Elsevier, Netherlands, 2004.
5. Loijen R.C.- Holism and reductionism in biology and ecology – The mutual dependence of higher and lower research programmes, Stichting Drukkerij, Regenborg, 1998.
6. Pahl Vostl - The dynamic nature of ecosystems, J. Willey and Sons, New York, 1995.
7. Pinay G., Gumiero B., Tabacchi E., Gimenez O., Planty-Tabacchi A. M., Hefting M., Burt T., Black V., Nilsson C., Iordache V., Bureau F., Vought L., Petts G., Décamps H.- Patterns of denitrification rates in European alluvial soils under various hydrological regimes, Freshw. Biol., 2007, 52, 252-266
8. Schryver J.C., Craig C., Susan Brandt, Pfiffner M., Palumbo A.V., Peacock A.D., White D.C., McKinley J.P., Long P.E.- Application of Nonlinear Analysis Methods for Identifying Relationships Between Microbial Community Structure and Groundwater Geochemistry, Microbial ecology, 2006, 51, 177–188
9. Stanciulescu F.- Modelling of high complexity systems with applications, WIT Press, 2005.
- 10.\*\*\* - Food-web modeling for ecological assessment of terrestrial pollution (EcolMAT), ESF publication, [www.esf.org/ecolmat](http://www.esf.org/ecolmat), 2003,

# MINERAL ELEMENT CONTENT OF LEAVES OF TABLE BEET VARIETIES AND THEIR ROLE IN NUTRITION

Maria TAKACS-HAJOS<sup>1\*</sup>, Rudolf KASTORI<sup>2</sup>, Mira PUCAREVIC<sup>2</sup>,  
Tijana Zeremski SKORIC<sup>2</sup>

1\*. Tessedik Sámuel College, Faculty of Agricultural Water- and Environment Management, 1-3. Szabadság str., H-5540 Szarvas, Hungary; 2. Institute of Field and Vegetable Crops, Maksima Gorkog 30, RS-21 000 Novi Sad, Serbia

\* E-mail : hajos.maria@mvk.tsf.hu

## ABSTRACT

In our experiment the production of baby beetroot, grown by a method different from the traditional one, was examined with shorter maturation time (70 days) and different genotypes on alluvial meadow soil. In this growing phase the tender leaves are edible so the examination of mineral element content of lamina and petiole was considered important. It was found that the Mg, Ca, Mn, Fe, Cu and Zn content of the lamina was the multiple of the values measured in the petiole. The nutrition-physiologically important ion ratio –  $(Ca^{2+}+Na^{+})/(Mg^{2+}+K^{+})$ , the optimal value of which in the human organism is ~1.0 – showed better, i.e. lower values in the petiole in all varieties. So, it is worth consuming not only the lamina of beetroot leaves but also their petioles because their mineral element content and composition are very favourable. There is difference between the varieties especially in the Fe content. In this regard the best genotype was the Boro F<sub>1</sub>.

**Key words:** table beet, mineral element, lamina, petiole, ion ratio, genotypes

## INTRODUCTION

The mineral element content of root of tablebeet and its nutrition physiological effects have been demonstrated in several studies (Stefanovits Bányai et al., 2002, Blázovits et al, 2007). It is well-known from the technical literature that the leaves of tablebeet were used by the people of the ancient times mainly for medicinal purposes. As a result several scientific papers were published on the bioactive materials of tablebeet leaves. The folic acid content (vitamin B derivative) is very high, thus the consumption of leaves of tablebeet is very useful from point of view of nutrition physiology (Maison, 1994). Based on the above mentioned arguments not only the root of the table beet but also its foliage can enrich our vegetable selection. Thus the leaves of 6-8 cm size gained at thinning can be used for making salads.

The modern alimentation prefers the consuming of more vegetables preferably in fresh form. The special salads have become very popular. In gastronomy they are called mesclun. The word comes from the Latin word *misculare* that means mixing. Nowadays this word is used for naming the multicoloured mixed salads. For making

these salads the very young, tender leafy parts of all vegetables are used. They should be very fresh and crunchy and possibly different in colour (claret, yellow, as well as light and dark green). If we prepare a mesclun salad ourselves the leafy parts of the following vegetables should be mixed into it: green lettuce, leaves of baby spinach, tender spinach beet and table beet leaves as well as purslane, lamb's lettuce, dandelion, chervil and parsley.

The favourable nutrition physiological effects of tablebeet are proven by its mineral element content, which is shown in Table 1 compared to the root of tablebeet.

**Table1.** Mineral element content in different parts of beetroot

| Elements       |    | Element content (mg/100g raw material) |              |             |
|----------------|----|----------------------------------------|--------------|-------------|
|                |    | in leaves                              | in the flesh | in the rind |
| Macro-elements | Ca | 156.0                                  | 10.04        | 21.19       |
|                | Mg | 170.3                                  | 22.23        | 44.33       |
|                | K  | 687.7                                  | 296.4        | 375.70      |
|                | Na | 540,8                                  | 121,4        | 158,60      |
|                | P  | 37,38                                  | 49,4         | 65,65       |
| Micro-elements | Zn | 0,88                                   | 0,40         | 0,54        |
|                | Fe | 2,91                                   | 0,85         | 2,09        |
|                | Cu | 0,20                                   | 0,11         | 0,13        |
|                | Mn | 1,15                                   | 0,22         | 0,28        |
|                | Al | 5,68                                   | 0,50         | 0,58        |

Based on our measurements (Takácsné Hájos et al., 2000) it is visible that the Ca, Mg and K content of rinds of tablebeet leaves is several times higher than that the values measured in the root.

In growing of this plant the thinning of tablebeet individuals at the 2-4 leaves stage is a part of the production technology when the surplus individuals are removed. These plants are suggested to be used for making fresh vegetable salads as at this stage the leaves can be used together with the petioles. The folic acid content of tablebeet is also very important, however its amount is highly dependent on the varieties. In the varieties characterized by intensively green leaves this compound has been found in higher proportion (Wang - Goldman, 1996).

Taking into consideration the above mentioned reasons and the measurement data, the leaves of tablebeet could be utilized for human consumption offering a bigger choice of vegetables.

## MATERIALS AND METHODS

The experiments were conducted in the Horticultural Experimental Farm of Tessedik Sámuel College, Faculty of Agricultural Water and Environmental Management, on alluvial meadow soil. Six beetroot varieties (*Beta vulgaris* L. ssp. *esculenta* var. *rubra*) were examined. Among them there were five Dutch varieties (Taunus, Boro F<sub>1</sub>, Pablo F<sub>1</sub>, Bikores, Little Ball) and one Hungarian variety (Rubin). The sowing was carried out on 26 April, 2006 where the row space was 15 cm and the size of the plot was 2 m x 0,3 m. Harvesting was carried out on 6 July, 2006 and after it the following measurements were taken: Mineral element values were determined by an atomic absorption method using ICP OES IRIS Thermo Jarrel ASH

device. It was conducted in the Institute of Field and Vegetable Crops, Novi Sad. The characterization of soil of our experimental field is shown in Table 2.

**Table 2** Soil composition

| pH  | EC                 | NO <sub>3</sub> -N | P                  | K                  | Ca                 | Mg                 | Na                 | Cl                 | HCO <sub>3</sub>   |
|-----|--------------------|--------------------|--------------------|--------------------|--------------------|--------------------|--------------------|--------------------|--------------------|
|     | mScm <sup>-1</sup> | mgkg <sup>-1</sup> | mgkg <sup>-1</sup> | mgkg <sup>-1</sup> | mgkg <sup>-1</sup> | Mgkg <sup>-1</sup> | mgkg <sup>-1</sup> | mgkg <sup>-1</sup> | mgkg <sup>-1</sup> |
| 6.9 | 0.15               | 8.23               | 1.26               | 21.13              | 12.33              | 6.96               | 16.86              | 11.43              | 32.10              |

pH and EC values include distilled water extract in a ratio of 1:2.

## RESULTS AND DISCUSSIONS

It can be concluded that the Mg, Ca, content of the lamina is multiple of the values measured in the petiole (Tables 3-4, and Fig.1), whereas considering the differences in K and Na supply the differences can be neglected in most varieties. The nutrition-physiologically important ion ratio –  $(Ca^{2+}+Na^{+})/(Mg^{2+}+K^{+})$ , the optimal value of which in the human organism is ~1.0 – showed better, i.e. lower values in the petiole in all varieties. So, it is worth consuming not only the lamina of beetroot leaves but also their petioles because their mineral element content and composition are very favourable.

**Table 3** Element content (g kg<sup>-1</sup>) and ion ratio in the petiole of table beet varieties

| varieties / elements | K     | Mg   | Na    | Ca   | $(Ca^{2+}+Na^{+})/(Mg^{2+}+K^{+})$ |
|----------------------|-------|------|-------|------|------------------------------------|
| Taunus               | 61,74 | 2,75 | 39,63 | 4,31 | 0,681                              |
| Boro F1              | 54,15 | 3,41 | 26,73 | 4,14 | 0,536                              |
| Pablo F1             | 54,36 | 3,53 | 31,09 | 4,30 | 0,611                              |
| Bikores              | 48,72 | 2,33 | 26,46 | 3,55 | 0,588                              |
| Rubin                | 54,47 | 2,69 | 34,75 | 4,83 | 0,692                              |
| Little Ball          | 53,62 | 2,83 | 30,29 | 5,15 | 0,628                              |
| mean                 | 54,51 | 2,92 | 31,49 | 4,38 | 0,62                               |
| St. dev.             | 4,16  | 0,46 | 5,03  | 0,56 | 0,06                               |

**Table 4** Element content (g kg<sup>-1</sup>) and ion ratio in the lamina of table beet varieties

| varieties / elements | K     | Mg    | Na    | Ca    | $(Ca^{2+}+Na^{+})/(Mg^{2+}+K^{+})$ |
|----------------------|-------|-------|-------|-------|------------------------------------|
| Taunus               | 50,81 | 17,31 | 36,97 | 15,14 | 0,765                              |
| Boro F1              | 43,15 | 20,00 | 35,66 | 15,97 | 0,818                              |
| Pablo F1             | 41,67 | 19,98 | 33,39 | 17,60 | 0,827                              |
| Bikores              | 45,69 | 20,22 | 37,26 | 16,35 | 0,813                              |
| Rubin                | 44,73 | 15,66 | 33,08 | 16,45 | 0,820                              |
| Little Ball          | 46,46 | 19,26 | 39,71 | 18,75 | 0,889                              |
| mean                 | 45,42 | 18,74 | 36,01 | 16,71 | 0,82                               |
| St. dev.             | 3,16  | 1,85  | 2,52  | 1,28  | 0,04                               |




**Fig. 1.** Mezo-element content in the different part of table beet leaves

Evaluating the mezo-elements (Table 5-6, Fig.2) we considered, that higher amount of them is the Fe in the table beet leaves. The most important statement, that the Mn and Fe content of lamina is multiple as those in petiole.

**Table 5** Mezo-element content ( $\text{mg kg}^{-1}$ ) in the petiole of table beet varieties

| varieties / elements | Cu    | Zn    | Mn    | Fe     | Al     |
|----------------------|-------|-------|-------|--------|--------|
| Taurus               | 13,05 | 15,43 | 54,86 | 209,53 | 305,43 |
| Boro F1              | 18,99 | 20,97 | 51,51 | 523,53 | 513,65 |
| Pablo F1             | 25,31 | 20,75 | 52,31 | 456,40 | 588,10 |
| Bikores              | 15,79 | 20,94 | 39,82 | 445,63 | 582,73 |
| Rubin                | 17,60 | 17,46 | 60,84 | 369,70 | 414,55 |
| Little Ball          | 11,09 | 20,10 | 45,10 | 384,40 | 388,90 |
| mean                 | 16,97 | 19,28 | 50,74 | 398,20 | 465,56 |
| St. dev.             | 5,01  | 2,31  | 7,39  | 107,65 | 114,17 |

**Table 6** Mezo-element content ( $\text{mg kg}^{-1}$ ) in the lamina of table beet varieties

| varieties / elements | Cu    | Zn    | Mn     | Fe      | Al      |
|----------------------|-------|-------|--------|---------|---------|
| Taurus               | 42,60 | 27,93 | 357,83 | 743,43  | 689,15  |
| Boro F1              | 52,55 | 50,93 | 238,55 | 1144,08 | 1189,90 |
| Pablo F1             | 61,83 | 49,28 | 334,47 | 876,18  | 1075,70 |
| Bikores              | 65,09 | 49,41 | 238,02 | 1035,00 | 903,78  |
| Rubin                | 66,02 | 39,55 | 354,42 | 983,00  | 809,75  |
| Little Ball          | 47,71 | 37,10 | 333,70 | 1102,88 | 1195,48 |
| mean                 | 55,97 | 42,37 | 309,50 | 980,76  | 977,29  |
| St. dev.             | 9,77  | 9,11  | 56,05  | 149,53  | 209,28  |




**Fig. 2.** Mezo-element content in the different part of table beet leaves

We measured great difference in Fe content between the varieties (Fig.3) which is may be connected with the assimilation ability of lamina. The Boro F1 exceeded in several important mineral element content. It is affirmed, that choice of varieties is very important in this roil too.




**Fig. 3.** Fe-content in the table beet varieties

## CONCLUSIONS

It can be concluded that the Mg, Ca, Mn, Fe, Cu and Zn content of the lamina is multiple of the values measured in the petiole, whereas considering the differences in K and Na supply, the differences can be neglected in most varieties. The nutrition-physiologically important ion ratio –  $(Ca^{2+} + Na^+) / (Mg^{2+} + K^+)$ , the optimal value of which in the human organism is  $\sim 1.0$  – showed better, i.e. lower values in the petiole in all

varieties. So, it is worth consuming not only the lamina of beetroot leaves but also their petioles because their mineral element content and composition are very favourable. By the way it seems very important the choice of varieties too, because there is great difference in mineral element content.

## REFERENCES

1. Blázovics A., Sárdi É., Szentmihályi K., Váli L., Takács-Hajos M., Stefanovits Bányai É. - Extreme consumption of *Beta vulgaris* L. var. *rubra* can cause metal ion accumulation in the liver, *Acta Biologica Hungarica*, 2007, 58(3), 281-286.
2. Maisin J.B. - Folate and colonic carcinogenesis: search for a mechanistic understanding, *J. Nutr. Biochem.*, 1994, 5, 170-175
3. Stefanovits Bányai É., Kiss A. S., Csikkel Szolnoki A., Sz. Varga I., Takacs Hajos M. - Phytochemical and macroelement study of *Beta vulgaris* L. ssp. *esculenta* var. *rubra*., *Central European Journal of Occupational and Environmental Medicine*, 2002, (8), 2-3. 167-171.
4. Takácsné Hájos M., Csikkel-Szolnoki A., Kiss A.S. - Mineral content of table beet roots as depending on varieties, *Magnesium Res.*, 2000, 12, 326-327.
5. Wang M., Goldman I. L. - Phenotypic variation in free folic acid content among F1 hybrids and open pollinated cultivars of red beet, *J. Amer. Hort. Sci.*, 1996, 121 (6), 1040-1042.



## CHARACTERISTICS OF THE METALLOGRAME IN CYSTINE UROLITHIASIS – INVESTIGATION MEANS

Adina-Elena AVACOVICI <sup>1\*</sup>, Zeno GARBAN<sup>2</sup>

1. Faculty of Chemistry and Environmental Engineering, University Politehnica Timișoara, Pta Victoriei, Nr.2, RO-300 006 Timișoara, Roumania; 2. Department of Biochemistry-Molecular Biology, University of Agricultural Sciences and Veterinary Medicine Timișoara, Calea Aradului Nr. 119, RO-300 457 Timișoara, Roumania

\* E-mail : avacovici@yahoo.com

### ABSTRACT

*Urinary stone disease reveals a multifactorial ethiogenesis. All types of renal stones originate on excessive supersaturation of the urine with a poorly soluble material. This depends not only on solute concentration but also on pH, ionic strength and the existence of soluble complexes.*

*Cystine crystals are found in urine of almost exclusively patients with a genetic disease giving an impairment in renal and intestinal cystine transport with decreased proximal tubular reabsorption resulting an increased cystine excretion.*

*Our study used infrared spectroscopy (IRS) for the determination of urolithiasis' types and atomic absorption spectroscopy (AAS) to draw the metallograms.*

**Keywords:** cystine urolithiasis , metallograms

### INTRODUCTION

Cystine is an amino acid dimer formed from two molecules of cysteine joined by a disulphidic bond. Cystinuria is an autosomal recessive disorder of the amino acid transport in the kidney leading to an abundance of cystine in the urine. This is a poorly soluble molecule that crystallizes at concentration of above 300 mg/L. This value may be regarded as the "clinical saturation point" for cysteine. Over this concentration, cystine calculi may form or enlarge. There have been described more types of cystinuria: due to two mutations of the SLC3A1 gene encoding a subunit of the transport mechanism; two mutations of the SLC7A9 gene encoding the transport channel itself. There also appear mixed types from the combination of recessive and dominant alleles (Matouschek, 1981; Heining, 1986; Goodyer, 1998). Cystinuria is the most common defect in the transport of an amino acid.

Although the exact molecular basis is not yet elucidated, future investigations are based on molecular and gene therapy.

### MATERIALS AND METHODS

Cystine crystals form hexagonal-shaped crystals which can be viewed upon microscopic analysis of the urine. Cystine stones are visible on standard abdominal

radiographs because of the relative density of the sulphur constituent of cystine (Robertson and Peacock, 1984; Watts, 1984; Coe and Parks, 1988; Martins et al., 2002; Ahmed et al., 2006; Evan et al., 2006).

*Sample prelevation.* Investigations have been performed on 92 calculi surgically removed or spontaneously eliminated by patients from the University Clinic of Urology Timișoara.

*Analytical investigations.* In previous investigations we obtained by IRS the standard spectra of the pure compounds frequently found in kidney stones. In the first step we recorded the spectra of each calculus and established, by comparison with the data base, the composition of uroconcrements. Subsequently, by AAS, was determined the metal composition, i.e. the content of alkaline, alkaline-earth and transitional metals.

*Statistical evaluation.* By the use of a computerized method, the mean value (X) and standard deviation (SD) were obtained.

## RESULTS AND DISCUSSIONS

Although cystine is not the only overly excreted amino acid in cystinuria, it is the least soluble of all naturally occurring amino acids. The key process in the development of kidney stones is supersaturation. As to frequency among other types of urolithiasis, cystinic calculi are about 1% in adults and up to 8% in children. The stones are usually multiple, they tend to recur after surgical removal, and bacterial infections often complicate the clinical signs (Williams and Chisholm, 1976; Ala-Opas et al., 1990; Galka et al., 1990; Sakhaee, 1994)

The IRS identification of unknown compounds makes use of standard spectra that offer data on characteristic absorption domains and vibrational states. Fig. 1 presents the standard spectra of cystine.




**Fig. 1.** Standard spectra of cystine

The cystine transporter also promotes the reabsorption of the other dibasic amino acids such as: cystine, lysine, ornithine, arginine, homoarginine. The structural formula of the amino acids excreted in high amounts are presented in figure 2.




**Fig. 2.** Amino acids in excess decelated in cystinuria

Metal ions present in the blood and urine play an important role in the organism. Some of these metals can prevent the concrements formation at the level of urinary tract, gall bladder or salivary gland while the others inhibit the crystallization.

Metal ions compete in the process of coprecipitation of the calculi' ingredients by the bioinorganic mechanism of lithogenesis.

A synopsis of the types of studied lithiasis is given in table 1 and table 2. One can observe that out of all 90 calculi only 3 contained cystine.

**Table 1.** Simple urolithiasis revealed by IRS

| Type of urolithiasis        | Composition         |                                | Number of cases |     |       |
|-----------------------------|---------------------|--------------------------------|-----------------|-----|-------|
|                             |                     |                                | Total           | Men | Women |
| SIMPLE                      | Oxalates (O)        |                                | 21              | 12  | 9     |
|                             | Purinic derivatives | Urates (U)                     | 15              | 7   | 8     |
|                             |                     | Xanthine (X)                   | --              | --  | --    |
|                             |                     | 2,8-dihydroxyadenine (2,8-DHA) | --              | --  | --    |
|                             | Phosphates (F)      |                                | 14              | 5   | 9     |
|                             | Cystine (C)         |                                | 3               | 1   | 2     |
| Total – simple urolithiasis |                     |                                | 53              | 25  | 28    |

Table 2. Mixed urolithiasis revealed by IRS

| Type of urolithiasis      | Composition |                                               | Number of cases |     |       |
|---------------------------|-------------|-----------------------------------------------|-----------------|-----|-------|
|                           |             |                                               | Total           | Men | Women |
| MIXTE                     | Binary      | Oxalates- Phosphates (O-P)                    | 13              | 6   | 7     |
|                           |             | Oxalates - Urates (O-U)                       | 5               | 1   | 4     |
|                           |             | Oxalates - Cholesterol (O-COL)                | 3               | 2   | 1     |
|                           |             | Urates -Xanthine (U-X)                        | 3               | 1   | 2     |
|                           |             | Phosphates - Oxalates (P-O)                   | 6               | 2   | 4     |
|                           |             | Phosphates - Carbonates (P-CARB)              | 3               | 1   | 2     |
|                           | Ternary     | Phosphates – Oxalates - Carbonates (P-O-CARB) | 1               | 1   | --    |
|                           |             | Oxalates - Urates - Phosphates (O-U-P)        | 3               | 1   | 2     |
| Total –mixte urolithiasis |             |                                               | 37              | 15  | 22    |

The presence of metals in the organisms, their homeostasis, their involvement in metabolic processes as well as the perturbation of these processes, e.g. urolithiasis, represent problems of special interest (Daranyi et al., 1994; Garban et al., 1996; Dragan et al., 1998; Worcester et al., 2006). Determination of the metals present in uroconcrements can contribute, indirectly, to the explanation of the mechanism of heterogenous nucleation to which organic and inorganic metabolites compete.

Determination of concentration of the main alkaline, alkaline-earth is presented in table 3.

**Table 3.** Metallogram of cystinic uroconcrements (alkaline and alkaline-earth metals)

| Metals         |    | UM                       | No. of cases | Concentration of metals<br>$\bar{X} \pm SD$ |
|----------------|----|--------------------------|--------------|---------------------------------------------|
| Alkaline       | Na | $\mu\text{g/g calculus}$ | 11           | $242.53 \pm 58.10$                          |
|                | K  |                          | 11           | $601.77 \pm 183.15$                         |
| Alkaline-earth | Ca |                          | 11           | $3050.86 \pm 1416.78$                       |
|                | Mg |                          | 11           | $320.18 \pm 121.37$                         |

Determinations regarding transitional metals, that exist in more reduced quantities than the alkaline and alkaline-earth ones, are presented in table 4. There are given results for a greater number of calculi, i.e. there were added 8 more from a previous period of analytical determinations.

**Table 4.** Metallogram of cystinic uroconcrements (microelements)

| Metals | UM                          | No. of cases | Concentration of metals<br>$\bar{X} \pm SD$ |
|--------|-----------------------------|--------------|---------------------------------------------|
| Zn     | $\mu\text{g/g}$<br>calculus | 11           | $41.20 \pm 14.45$                           |
| Fe     |                             | 11           | $14.51 \pm 4.05$                            |
| Cu     |                             | 11           | $15.06 \pm 4.16$                            |
| Mn     |                             | 11           | $4.36 \pm 1.83$                             |

Researches performed at the University Clinic of Urology Timișoara revealed the role of metals in the etiopathogeny of lithiasis and led to the hypothesis of bioinorganic mechanism of urolithogenesis (Garban and Daranyi, 1994; Dragan et al., 1998). Obviously, this mechanism does not facilitate the formation of starters and thereafter, of uroconcrements (Williams și Chisholm, 1976).

## CONCLUSIONS

1. The appearance of metals in urolithiasis is the consequence of their presence in urine;
2. By means of infrared spectroscopy the types of lithiasis (simple and mixed ones) have been established;
3. Metallograms were drawn using atomic absorption spectroscopy, revealing: alkaline metals:  $\text{Na} < \text{K}$ ; alkaline earth metals :  $\text{Mg} < \text{Ca}$ ; microelements:  $\text{Mn} < \text{Fe} < \text{Cu} < \text{Zn}$
4. Metals play a predictive role on the bioinorganic composition of cystinic uroconcrements.

*Acknowledgements.* The authors express their thanks to the doctors from the Clinic of Urology Timișoara who cooperated at the collection of calculi from surgical interventions. It is mentioned that the University Clinic of Urology, University of Medicine and Pharmacy "Victor Babeș" Timișoara, was led by Prof. dr. Petru Dragan (1932-2007).

## REFERENCES

1. Ahmed K, Dasgupta P, Khan MS : Cystine calculi: challenging group of stones, *Postgraduate medical journal*, 2006, 82 (974): 799–801
2. Ala-Opas M., Sipila I., Kivinen S., Lethonen T. : Diagnostic problems associated with cystinuria, *Scand. J. Urol. Nephrol.*, 1990, 24, 133-135
3. Coe F.L., Parks J.H. : Nephrolithiasis: Pathogenesis and Treatment. Year Book Medical Publishers Inc., Chicago-London-Boca Raton, 1988.
4. Daranyi Gabriela, Drăgan P., Bucuraș V., Matusz Anca, Papoe Gabriela, Martoni Livia, Gârban Z. : Investigations on the composition and etiopathogeny of lithiasis. XVI. Metallic composition of cystinic uroconcrements, pp.77-80, *Metal Elements in Environment, Medicine and Biology*, Tome I (Eds. Dragan P., Garban Z.), Publishing House Mirton Timișoara, 1994.
5. Dragan P., Garban Z., Holban Șt. : Clinical aspects and biochemical pathology in cystinic urolithiasis - peculiarities of the metallic composition, pp. 27-34, in *Metal*

- Elements in Environment, Medicine and Biology*, Tome III (Eds. Garban Z., Dragan P.), Publishing House Eurobit Timișoara, 1998
6. Evan A.P., Coe F.L., Lingeman J.E., et al.: Renal crystal deposits and histopathology in patients with cystine stones. *Kidney Int* 2006; 69:2227.
  7. Galka M., Paluszkiwicz C., Czaja M., Scieslinski J., Kwiatek W.: Analysis of surgically removed renal concretions in recurrent cystine nephrolithiasis, *Urologia Polska*, 1990, 43 (3), 185-188.
  8. Garban Z., Daranyi Gabriela: Peculiarities of the metal composition of cystinic urolithiasis, pp.662-667, in *"Defizite und Überschuss an Mengen- und Spurenelemente in der Ernährung"*, 14. Arbeitstagung Mengen- und Spurenelemente, 1994, Friedrich-Schiller-Universität Jena, (Hrsg. Anke M., Meissner D. et al.), Verlag Harald Schubert, Leipzig, 1994.
  9. Garban Z., Daranyi Gabriela, Dragan P., Sigărtău Gr., Avacovici Adina, Urzică A.: Investigations on the composition and etiopathogeny of lithiasis. XXVII. Thermal analysis and atomic absorption spectroscopy in the identification of urolithiasis, pp. 307-312, in *"New Perspectives in the Research of Hardly Known Trace Elements"* (ed. Pais I.) University Press of U.H.F.S. Budapest, 1996
  10. Goodyer, P, Saadi, I, Ong, P, et al.: Cystinuria subtype and the risk of nephrolithiasis. *Kidney Int* 1998; 54:56.
  11. Heininger J.A., Verjee Z.H., McNeely M.D.D.: Disorders of amino acids and protein metabolism. pp.420-43, in *"Applied Biochemistry of Clinical Disorders"* 2nd ed. (Gornall A.G., ed.), J.B. Lippincott Company, Philadelphia, 1986.
  12. Martins M.C., Meyers A.A., Whalley N.A., Rodgers A.L.: Cystine: a promoter of the growth and aggregation of calcium oxalate crystals in normal undiluted human urine., *J Urol.* 2002; 167(1): 317-21
  13. Mathoushek E., Huber R.D.: *Urolithiasis: Pathogenese, Diagnostik, Therapie*, Schattauer Verlag, Stuttgart-New York, 1981
  14. Robertson W.G., Peacock M.: Risk factors in the formation of urinary stones, pp. 267-278, in *Scientific Foundations of Urology*, Second ed. (Eds. Chisholm G.D., Williams D.I.), William Heinemann Medical Books Ltd., London, 1984
  15. Sakhaee K. - Cystinuria: pathogenesis and treatment, *Miner. Electrolyte Metab.*, 1994, 20(6), 414-423
  16. Watts R. W. E.: Cystinuria and cystine stone disease. pp. 302-309, in *Scientific Foundations of Urology*, Vol.I, (Williams D., Chisholm G.D., eds.), W. Heinemann Medical Books Ltd., London 1984.
  17. Williams D.I., Chisholm G.D.: *Scientific Foundations of Urology*, Second ed., William Heinemann Medical Books Ltd., London, 1976
  18. Worcester E.M., Coe F.L., Evan A.P., Parks J.H.: Reduced renal function and benefits of treatment in cystinuria vs other forms of nephrolithiasis. *BJU Int* 2006; 97:1285.

# CONSEQUENCES OF SUBACUTE INTRATRACHEAL EXPOSURE OF RATS TO CADMIUM OXIDE NANOPARTICLES: ELECTROPHYSIOLOGICAL AND TOXICOLOGICAL EFFECTS

Andras PAPP\*, Leila SARKÖZI

Dept. of Public Health, University of Szeged, Faculty of Medicine, Dom Tér 10, H-6720 Szeged, Hungary

\* Email: ppp@puhe.szote.u-szeged.hu

## ABSTRACT

*Cadmium is a metal used in various industrial applications, whereby emission of Cd-containing fumes is likely. The submicron sized particles in the fumes represent an extra risk due to their high mobility within the organism and high surface area. Toxicity of Cd on the liver, kidney and bones is well known, but there are less data on its neurotoxicity. Here, male Wistar rats were treated for 3 and 6 weeks by intratracheal instillation of CdO<sub>2</sub> nanosuspension. The treated rats' body weight gain was significantly decreased, and in the high dose (0.4 mg/kg Cd daily) rats the weight of lungs and thymus was significantly increased. In this group, the spectrum of spontaneous cortical electrical activity was shifted to higher frequencies, the latency of the somatosensory evoked potential was lengthened and its frequency following ability, impaired. The visual evoked potential also had increased latency. Cadmium had access from the airways to the brain and exerted neurotoxic effects.*

**Key words:** cadmium, nanoparticles, lungs, body weight, organ weight, electrophysiology

## INTRODUCTION

Cadmium has various industrial applications in steel and other alloys, pigments, in batteries and in electroplating. Occupational exposure is mainly due to inhalation of metal fumes (WHO, 1992), and the exposure of smokers due to Cd content of tobacco leaves is similar (Elinder et al., 1983). Cd is a heavy metal with considerable general toxicity. Its main target organs include the liver, kidneys and bones, and the respiratory and cardiovascular systems (ATSDR, 1999), but Cd exposure can also cause behavioral and neurological disorders. In occupational Cd exposure, reduced visuomotor performance and difficulties of concentration and postural balance were observed (Viaene et al., 2000). In children, a straight relationship between hair Cd and visual or auditory evoked potential parameters was found (Thatcher et al., 1984), and school behavioural problems were reported (Marlowe et al., 1985). Similar effects were observed in rats (Agar et al., 1990; Yargicoglu et al., 1996). Our previous works on Cd neurotoxicity revealed altered electrocorticogram (ECoG) power spectrum, and effects on cortical evoked potentials and peripheral nerve action potentials (Papp et al., 2003; Institóris et al., 2002).

---

„Metal Elements in Environment, Medicine and Biology”, Tome VIII, pp.67-72, Publishing House “Eurobit” Timișoara, 2008

Proc. of 8<sup>th</sup> Int. Symp. of Roumanian Academy - Branch Cluj-Napoca and Branch Timișoara, 2008, December 5-6, Timișoara, Roumania

Cd containing metal fumes (from smelting, casting, welding etc.) contain both microscopic and submicroscopic particles, and the two show dissimilar behavior after inhalation (Elder et al., 2006). The health importance of submicron particles has been recognized newly, and attributed to their high number and large surface area (Oberdörster et al., 2005). Nanoparticles (NPs) easily cross barriers within the organism, such as the respiratory or olfactory epithel and the blood-brain barrier, not penetrable for larger (microscopic) particles. Transport from the olfactory epithel directly into the CNS via then olfactory nerve has also been described.

Within the framework of the *Regional University Knowledge Centre for Environmental and Nanotechnology, Szeged, Hungary* a series of animal experiments has been carried out to see the nervous system effects of toxicologically relevant nanoparticles. In the present work, CdO<sub>2</sub> nanoparticles were given to adult rats, and general and nervous system effects were observed.

## MATERIALS AND METHODS

Adult male Wistar rats of 320-350 g body weight were housed under standard conditions (22 - 24°C, 12-hour light/dark cycle with light on at 06:00) with free access to tap water and standard pellet. The rats were divided into 4 groups of 20 animals each at start. Nanoparticles of CdO<sub>2</sub> (20±5.7 nm diameter) were synthesized at the Department of Applied Chemistry, University of Szeged, and were suspended in distilled water. The suspension was instilled into the rats' trachea, in the daily doses shown in Table 1., 5 days a week (Monday to Friday). The volume instilled was 1 ml/kg b.w. Treatment was continued for 3 and 6 weeks, so that 10 rats from each group were sacrificed after 3 weeks treatment, and the remaining 10 after 6 weeks. There was an untreated control group (Con), and a vehicle control group (W), as shown in Table 1.

**Table 1.** Treatment groups and doses.

| Group             | Code | Treatment and dose                                                   | Duration      |
|-------------------|------|----------------------------------------------------------------------|---------------|
| Untreated control | Con  | ---                                                                  |               |
| Vehicle control   | W    | Distilled water<br>1 ml/kg b.w.                                      |               |
| Low dose          | LD   | CdO <sub>2</sub> nanosuspension,<br>0.04 mg Cd/kg b.w.; 1 ml/kg b.w. | 3 and 6 weeks |
| High dose         | HD   | CdO <sub>2</sub> nanosuspension,<br>0.4 mg Cd/kg b.w.; 1 ml/kg b.w.  |               |

Intratracheal instillation was done in diethyl ether anesthesia. The rat was suspended on a 60° inclined board so that its upper incisors were held by a wire loop to keep the animal's mouth open. The trachea was illuminated transdermally. The tongue was pulled forward with a pair of non-traumatic forceps, and a custom-made laryngoscope was used to visualize the glottis. The nanosuspension (or distilled water for group W) was instilled into the trachea by means of a syringe and 1.2 mm OD plastic tubing, inserted between the vocal chords. The untreated control group (Con) had neither ether anesthesia nor instillation, while the water control (W) group was anesthetized and instilled with distilled water. The nanosuspension was vigorously sonicated before, and repeatedly during, administration to prevent agglomeration.



Body and organ weights were the endpoints for general toxic effect of the nano-CdO<sub>2</sub>. The rats' body weight was measured each Monday during treatment, and once more on the day of sacrifice, and the mean body weight of the groups was plotted against time to see the course of weight gain. Following electrophysiology (see below), the rats were sacrificed by an overdose of urethane, dissected, and the organ weight of the brain, liver, lungs, heart, kidneys, spleen, thymus and adrenals was measured. Relative weights were calculated by relating organ weights to 100 g body weight.

Electrophysiological recording was done 3-4 days after the last instillation. In urethane anesthesia the animal's head was fixed and the sensory areas of left hemisphere were exposed. The wounds were sprayed with 10% lidocaine, and a thin layer of petroleum jelly was applied on the dura to prevent drying. After 30 minutes recovery, silver electrodes were placed on the primary somatosensory (SS), visual (VIS) and auditory (AUD) areas. Electrocorticogram (ECoG) was recorded from these areas for 6 minutes, and the relative spectral power of the frequency bands (delta, theta, alpha, beta1, beta2, gamma; standard human EEG bands) was determined.

Then, sensory evoked potentials (EPs) were recorded by the same electrodes. For somatosensory stimulation, 2 needles were inserted into the contralateral whiskery skin to deliver square electric pulses (3-4 V, 0.05 ms, 1-10 Hz). Visual stimulation was produced by a high-luminance white LED aimed directly at the rat's right eye, driven by 0.2 ms pulses at 1 Hz. The acoustic stimuli were clicks (1 Hz, 40 dB) guided from a miniature earphone into the animal's right ear via the hollow ear bar. Fifty stimuli of each modality per rat were applied and the evoked activity was recorded. After averaging, latency and duration of the evoked responses was measured manually (for details, see Lukács and Szabó, 2007).

The change of latency of the somatosensory EP with increasing stimulation frequency was also investigated as a possible indicator of the action of the treatment on the state of the cortex. All electrophysiological recording and analysis was done by means of the Neurosys 1.11 software (Experimetria Ltd, Budapest, Hungary). From the general toxicological, behavioral and electrophysiological data, group means ( $\pm$ SD) were calculated. The results were tested for significance with one-way ANOVA and the post hoc analysis was done by Scheffe's test.

## RESULTS AND DISCUSSIONS

Intratracheal exposure by the nanoparticulate CdO<sub>2</sub> had marked effect on the rats' body weight gain. As shown by Table 2, the untreated controls' (Con) weight

**Table 2.** Body weight gain (grams) of the rats during the 6 weeks of treatment

| Groups | Con          | W             | LD              | HD                               |
|--------|--------------|---------------|-----------------|----------------------------------|
| Weeks  |              |               |                 |                                  |
| 0      | 360 $\pm$ 18 | 360 $\pm$ 18  | 353 $\pm$ 26    | 353 $\pm$ 24                     |
| 1      | 382 $\pm$ 20 | 375 $\pm$ 21  | 364 $\pm$ 27    | 351 $\pm$ 31                     |
| 2      | 416 $\pm$ 24 | 400 $\pm$ 24  | 388 $\pm$ 32*   | 359 $\pm$ 31***                  |
| 3      | 434 $\pm$ 23 | 409 $\pm$ 24* | 400 $\pm$ 31**  | 378 $\pm$ 35***### <sup>oo</sup> |
| 4      | 462 $\pm$ 49 | 426 $\pm$ 28* | 409 $\pm$ 38*** | 411 $\pm$ 37***##                |
| 5      | 495 $\pm$ 27 | 445 $\pm$ 26  | 430 $\pm$ 37**  | 438 $\pm$ 36**                   |
| 6      | 510 $\pm$ 27 | 444 $\pm$ 32  | 439 $\pm$ 34*** | 450 $\pm$ 33***                  |

\*, \*\*, \*\*\*:  $p < 0.05$ ,  $0.01$ ,  $0.001$  vs. Con; ###, ####:  $p < 0.01$ ,  $0.001$  vs. W; <sup>oo</sup>:  $p < 0.01$  vs. LD.

The somatosensory EP showed significant latency increase in the HD group vs. Con (Fig. 3). The slight dependence of the latency on the inter-stimulus interval (inverse frequency) seen in Con was more expressed in the treated groups, up to the significant difference between the latencies obtained with 1000 and 100 ms interval in the HD group. The changes of the EP duration were less clear-cut but the increase in HD vs. Con was significant. The latency of the visual EP changed similarly to the

somatosensory EP in the treated animals (Fig. 4). The changes of the auditory EP were inconclusive.




**Fig. 3.** Latency (left) and duration (right) of the somatosensory evoked potentials in the control and treated rats at varied inter-stimulus intervals (in ms, see insert).

\* \*\*, \*\*\*:  $p < 0.05$ ,  $0.01$ ,  $0.001$  vs. control; °:  $p < 0.05$  vs. lowest frequency stimulation in the same group.




**Fig. 4.** Latency of the visual evoked potential in the control and treated rats. 1 s inter-stimulus interval only. Displayed as in Fig. 3.

The general and electrophysiological changes indicated that cadmium instilled into the rats' trachea in form of CdO<sub>2</sub> nanoparticles was in fact absorbed and unfolded its toxic effects. The ECoG shift was similar to that found with oral application of dissolved Cd for 12 weeks, alone (Papp et al., 2003) and combined with the insecticide propoxur (Institóris et al., 2002). The body and organ weight effects were also in agreement with our previous results. The time trend of body weights and the increase of alterations after 6 vs. 3 weeks indicated a gradual build-up of Cd and its effects in the treated rats' body. Absorption of Cd following inhalation (or similar exposure) has been described repeatedly. Takenaga et al. (2004) found Cd in the blood, liver and kidney of rats after inhalation of a much smaller amount than our dose. In exposed workers, elevated urine Cd level was associated with reduced visuomotor performance and difficulties of concentration and stance (Viaene et al., 2000). First of all the latter effect indicated that Cd had access from the lungs to the brain, despite its low permeability across the blood-brain barrier (ATSDR, 1999). In rats exposed by Cd orally for two months, deposition in the cortex, hippocampus and other brain regions was seen (Clark et al., 1985). EEG alterations were seen in rats after only 9 days of Cd exposure (Vataev et al., 1994).

Literature data and our own results sufficiently prove that Cd exposure bears a considerable risk of nervous system damage and that Cd, entering the organism via the airways, has access to the CNS. This means that workplace airborne Cd has to

be strictly regulated, Cd containing waste be treated accordingly, and Cd replaced with less hazardous substances where technically possible.

## REFERENCES

1. Agar A., Yargicoglu P., Edremitlioglu M., Kara C., Oguz Y.: The effect of cadmium on somatosensory evoked potentials (SEP) and conduction velocity in alloxane-induced diabetic rats: relation to lipid peroxidation. *J. Bas. Clin. Physiol. Pharmacol.* 1999, 10, 41-56
2. Clark D.E., Nation J.R., Bourgeois A.J., Hare M.F., Baker D.M., Hindeberger E.J.: The regional distribution of cadmium in the brains of orally exposed adult rats. *NeuroToxicology* 1985, 6, 109-114
3. Elder A., Gelein R., Silva V., Feikert T., Opanashuk L., Carter J., Potter R., Maynard A., Ito Y., Finkelstein J., Oberdörster G.: Translocation of ultrafine manganese oxide particles to the central nervous system. *Environ. Health Persp.* 2006, 114, 1172-1178
4. Elinder C.G., Kjellström T., Lind B., Linnman L., Piscator M., Sundstedt K.: Cadmium exposure from smoking cigarettes: variations with time and country where purchased. *Environ. Res.* 1982, 32, 220-227
5. Institóris L., Papp A., Siroki O., Banerjee B.D., Dési I.: Immuno- and neurotoxicological investigation of combined subacute exposure with the carbamate pesticide propoxur and cadmium in rats. *Toxicology*, 2002, 178, 161-173
6. Lukács A., Szabó A.: Functional neurotoxic effects of heavy metal combinations given to rats during pre- and postnatal development. *Centr. Eur. J. Occup. Environ. Med.* 2007, 13, 299-304
7. Marlowe M., Cossairt A., Moon C., Errera J., MacNeal A., Peak R., Ray J., Schroeder C.: Main and interaction effects of metallic toxins on classroom behavior. *J. Abnorm. Child. Psychol.* 1985, 13, 185-198
8. Oberdörster G., Oberdörster E., Oberdörster J.: Nanotoxicology: An Emerging discipline evolving from studies of ultrafine particles. *Environ. Health Persp.* 2005, 7, 823-839
9. Papp A., Nagymajtényi L., Dési I.: A study on electrophysiological effects of subchronic cadmium treatment in rats. *Env. Toxicol. Pharmacol.* 2003, 13, 181-186
10. Takenaga D.S., Karg E., Kreyling W.G., Lentner B., Schulz H., Ziesenis A., Schramel P., Heyder J.: Fate and toxic effects of inhaled ultrafine cadmium oxide particles in the rat lung. *Inhal. Toxicol.* 2004, 16 Suppl1, 83-92
11. Thatcher R.W., McAlaster R., Lester M.L.: Evoked potentials related to hair cadmium and lead in chiló dren. *Ann. N.Y. Acad. Sci.* 1984, 425, 384-390
12. Vataev S.I., Malgina N.A., Oganessian G.A.: The effect of cadmium on the structure of the circadian cycle of waking-sleep and on the EEG in Wistar Rats. [in Russian] *Zh. Evol. Biokhim. Fiziol.* 1994, 30, 408-419
13. Viaene M.K., Masschelein R., Leenders J., De Groof M., Swerts L.J., Roels H.A.: Neurobehavioural effects of occupational exposure to cadmium: a cross sectional epidemiological study. *Occup. Environ. Med.* 2000, 57, 19-27
14. Yargicoglu P., Agar A., Sentürk U.K., Izgut N., Oguz Y.: Effect of chronic cadmium exposure on VEP and EEG spectral components. *Int. J. Neurosci.* 1996, 85, 173-184
15. \*\*\* ATSDR: Toxicological Profile for Mercury. US Department of Health and Human Services, Atlanta, 1999
16. \*\*\*WHO: Cadmium. Environmental Health Criteria N° 134, Geneva, 1992

# BIOACCUMULATION OF METALS IN FOOD CHAINS OF WETLANDS

Aurora NEAGOE <sup>1\*</sup>, Virgil IORDACHE <sup>2</sup>, Florian BODESCU<sup>1</sup>

*1 Research Center for Ecological Services – CESEC, University of Bucharest, Soseaua Panduri No. 90-92, RO-050657 Bucuresti, Roumania; 2 Department of Systems Ecology, University of Bucharest, Roumania*

\* E-mail : aurora.neagoe@unibuc.ro

## ABSTRACT

*We studied the accumulation of metals in food chains of Danube floodplain, from soil and detritus towards vegetation, Carabidae and frogs. We found no simple pattern of the bioaccumulation of metals in the food chain, but most of the bio concentration occurred at the level of Carabidae, and not at the level of Rana sp. Mobile groups seem to not be a good indicator of pollution with metals at local level in large flood plains.*

**Key words:** heavy metals; flood plains; bioaccumulation;

## INTRODUCTION

Floodplains are involved in the retention of the metals transported by rivers (Finlayson, 1994; Wallach et al., 2001) and are a usual study model for the biogeochemistry of metals (Hobbelen et al., 2004; Griethuysen et al., 2005; Boivin et al., 2007). The flood plain of the lower Danube River has a length of hundreds of km and consists of riparian and insular micro landscapes (Vădineanu et al., 2001), 80% of which are diked. However, the lower Danube flood plain preserves several natural areas under natural flooding regime which are actively involved in the retention of nutrients and metals from the Danube basin (Iordache and Ruță, 2006).

The issue of the retention time of metals in flood plains is an important and still unresolved one. Remobilization processes and transfer through food chain are possible, with subsequent ecotoxicological risks. In particular, the metals can affect planktonic systems (Baca and Threlkeld, 2000), birds (Levengood and Skowron, 2001), amphibians (Bank et al., 2005), and detritus consuming invertebrates (Weis et al., 2002).

Estimating bioaccumulation of metals along the food chains is subject to the challenge of how to integrate the space-time heterogeneity of the distribution of organisms. This is an important issue especially where the fluxes between ecosystems are intense, as in flood plains. Mobile species would presumably intercept metals from larger areas than sessile species, but biogeochemical data to support this statement are scarce. Ground beetles (Carabidae) and frogs (Rana sp.) are mobile species commonly occurring in flood plains, usable as overall indicators of wetlands functionality.

Carabidae have been subject of previous studies in areas contaminated with metals, with focus on the distribution (Read et al., 1987, Andrews et al., 1989,

Rabitsch, 1995; Heikens et al., 2001) or the effects of metals (Lock et al., 2001; Stone et al., 2001; Lindqvist et al., 2005). Frogs have also been subject of such studies (Birdsall et al., 1986; Roe et al., 2005; Loumbourdis et al., 2007). By their diet, frogs integrate fluxes from various sources, ranging from phytofagous, to detritivorous and to raptor invertebrates (Cogălniceanu, 1997; Cogălniceanu et al., 1999). Carabidae species are an appropriate model for the diet of frogs because they belong to all these trophic groups (Ciubuc, 1997). However, the study of metals accumulation in Carabidae and frogs has seldom been put in the larger context of a food network.

In this context, our purpose in this article is to characterize the accumulation of metals in food chains of Danube floodplain, from soil and detritus towards vegetation, Carabidae and frogs. The results will show to what extent these compartments are useful as direct indicators of the local contamination with metals.

## MATERIALS AND METHODS

### *Site description*

We studied three micro landscapes located in the Small Island of Brăila, Danube floodplain (Figure 1). Two landscapes are located in islands (H and I), and one in the riparian area (G). “H” landscape is a late succesional one, with mature forests and marshes in its structure, and with low sediment dynamic during floods. “I” landscape is an early succesional one, with young forests and former channels in its structure, and with high sediment dynamic during floods. While H and I are in a natural state, “G” landscape is impacted by human activities: part of it is disconnected from floods by a dike, it includes a planted forest and it is grazed non-intensively by livestock.




**Fig. 1** The location of transects in the Danube floodplain. The vertical dimension is exaggerated in order to point out the floodplain

In each micro landscape we had four stations, from the Danube to the internal part of the landscape. Transect H started from the shore (H1), crossed a mature forest located on a high levee (H2), a natural forest from the nearby depression (H3), and a marsh (H4). Transect I crossed the shore (I1), a low forested levee (I2), a silted former Danube channel (I3), and a high forested levee (I4). Transect G included a shore (G1), a planted forest on high levee (G2), a depression (G3), and a diked depression (G4).

### ***Sampling program***

Top soil was sampled monthly during one year in five replicates. Detritus was sampled in five replicates using a 0.25 x 0.25m quadrat at several moments, depending on the state of the ecosystem (flooded or not). We had detritus of terrestrial origin (decomposed litter from previous years), and detritus of aquatic origin (snails and mussels at I1 and I3, snails at G1, snails and aquatic macrophytes at H3 and H4; details and species in lordache and Ignat in preparation). In some ecosystems both types of detritus were present (H3, H4). We sampled previously known dominant herbaceous plant species (Cristofor et al., 1997) at three moments during the growing season in nine replicates. The dominant plant species differed from one ecosystem to another (details and species in lordache in preparation). Terrestrial invertebrates were sampled fortnightly with nine traps in each ecosystems (method described in Ciubuc, 1999), in the frame of a project dealing with biodiversity. Dominant species of Carabidae were provided for analyses of metals concentrations, as follows: *Pseudophonus rufipes* (phytofagous), *Carabus cancellatus* (raptor), *Chlaenius festivus* (raptor), *Chlaenius spoliatus* (mostly detritivorous, sometimes raptor) and *Poecilus cupreus* (detritivorous). Frogs (*Rana* sp.,) were sampled on a transect base during several campaigns (using the methods described in Cogalniceanu, 1997), in the frame of a project dealing with the amphibians community. From transects G and I we analyzed adults (15 replicates), and from transect H adults, juveniles and tadpoles (10-15 replicates).

### ***Samples preparation and chemical analyses***

The soil samples were sieved at < 0.5 mm. The detritus of aquatic origin was separated in organic material and shells, in order to analyze only the organic material, accessible to detritivorous Carabidae. After harvesting, the plants were weighed (fresh weight) and partitioned for analysis into aboveground and underground part of plants. In this article we use the data characterizing the aboveground parts. Because Carabidae had been stored in ethanol before analyses (a method used also by Rabbitsch in 1995) we checked the effect of the preservation method by analyzing *Pseudophonus rufipes* individuals (the most frequent species) captured alive and stored by freezing or in ethanol (for 3 months). The t test using ten sample units of each type indicated no statistical differences between the concentrations of the analyzed metals. Adult frogs had been sub sampled for tissues: muscle, liver, gonad (in case of females) and "rest of body".

Soil, terrestrial detritus, plants and amphibians were dried in an oven at 105°, aquatic detritus and carabids were dried at 60°, then ground in an uncontaminated mill and stored at -20 °C until processing. All samples were analysed for Fe, Mn, Zn, Cu, Cr, Pb and Cd. For soil samples it was used the pseudo-total extraction with aqua regia (Hoffmann, 1991); for detritus, plant samples, Carabidae and amphibians the digestion was performed using nitric acid. The analyses were performed by F-AAS and GF-AAS (Perkin Elmer). The quality of the analyses was checked using BCR reference materials (river sediment, plant, and mussel tissue).

### **Data processing**

A weighted average of the concentration of metals in adult frogs was computed using the concentrations in each sub sample (tissue), and the weight of each sub sample. Such an average concentration in whole body is of interest not only from bioaccumulation point of view, but also when computing the stock of metals in the population of frogs.

Bioaccumulation in plants was assessed for each species and ecosystem based on the average concentration of metals in the soil of the ecosystem. Bioaccumulation in detritivorous Carabidae (including *Chlaenius spoliatus*) was assessed at landscape level based on a weighted average of metals in the detritus of the landscape. The concentration of detritus at each station was multiplied with the number of months in which the detritus was available to Carabidae (i.e. not flooded during the growing season, 2-6 months depending of the ecosystem), and the sum of concentration-months was divided by the total number of months-ecosystems. A similar approach was used for the bioaccumulation in phytofagous Carabidae. Bioaccumulation in raptor invertebrates (excluding *Chlaenius spoliatus*) was assessed based on the annual average of metals concentrations in the source compartments (detritivores or phytofags). Bioaccumulation in frogs was assessed based on the whole body concentration of adults and the annual average of metals concentrations in the source compartments.

## **RESULTS AND DISCUSSIONS**

The average bioaccumulation factors in all three micro landscapes are presented in Figure 2. Factors larger than 1 (bioconcentration) occurred most often for Pb and Cd, and in some cases for Fe, Zn, Cu. The metals were concentrated especially from plants to phytofagous Carabidae, and from phytofagous Carabidae to raptor Carabidae. The plants did not concentrate the metals from soil (although there are exceptions in case of some species which concentrate Mn in frequently flooded areas). Detritivorous Carabidae did not concentrate the metals from detritus, excepting in case of Cd. Frogs did not concentrate the metals from Carabidae, excepting for Pb from phytofagous species.


In Table 1 are presented differences of the concentration of metals average values between the stations of each transect and between transects (Mann-Whitney test). We do not present here the underlying data for space reason.

The mechanisms of metals retention in very large rivers are comparable to those in constructed wetlands), but the scale is different (Iordache et al., 2000; Iordache, 2003a, b). The percent of metals retained in the lower Danube floodplain from the upstream fluxes varied between 2.2% in case of Cd and 20.79% in case of Fe, but there are important differences in the pattern of metals' retention between the micro landscapes and ecosystems, depending on their succession stage and human impact (Iordache and Ruța, 2006). In this context one might expect to find consequent differences in the distribution of metals in the biotic compartments of the ecosystems.

Our results (further detailed in Iordache and Neagoe, in preparation) show that the number of significant differences between the ecosystems of a micro landscape decreases as one goes up in the food chain (from soil to detritus, plants, and Carabidae). Also the usefulness of the biological compartments for comparing the micro landscapes tends to decrease as we go to more mobile species. This is due probably to the fact that more mobile species can integrate fluxes from large areas (by having access to food in many ecosystems). A final answer to this issue can be



provided only by population ecology studies. It is also due to the heterogeneous pattern of metals transfer from one trophic level to another, impinging on the existence of a simple correlation between the metals distribution at the bottom and at the top of the food chain.



Grey cells indicate values of the bioaccumulation factor bigger than 1. Legend: Phyt. Car. = phytofagous Carabidea, Det. Car. = detritivorous Carabidae, Rap. Car. = raptor Carabidae.

**Fig. 2** Bioaccumulation of metals in the food chains of the investigated transects (average and SD).

Studying the metals biogeochemistry of flood plains is relevant for the design of decision support tools (Iordache et al., 2001), in particular for the monitoring system (Iordache and Bodescu, 2005). Floodplains management by restoration would lead to an important increase in the retention of metals and to a reduction of the pressure on the current wetlands (Iordache, 2002).

**Table 1** Differences of the concentration of metals average values between the stations of each transect and between transects (Mann-Whitney test).

| Transect |   | Fe                            | Mn                                   | Zn                                | Cu                 | Cr            | Pb                 | Cd                                    |
|----------|---|-------------------------------|--------------------------------------|-----------------------------------|--------------------|---------------|--------------------|---------------------------------------|
| G        | S | 14*, 24*, 34**                | 12**, 13**, 14***, 23**, 24**, 34*** | 12**, 14**, 23**                  | 12*, 13*           | NS            | NS                 | NS                                    |
|          | D | 12*, 13*, 14*                 | 12*, 13*, 14*                        | 12*, 13*, 14*                     | 12*, 13*, 14*, 23* | 12*, 13*, 14* | 12*, 13*, 14*, 23* | 12*, 13*, 14*                         |
|          | P | NS                            | NS                                   | NS                                | NS                 | NS            | NS                 | NS                                    |
|          | C | NS                            | NS                                   | NS                                | NS                 | NS            | NS                 | NS                                    |
| H        | S | 12*, 13***, 14***, 23**, 24** | 12***, 14*, 23***, 24***, 34**       | 12***, 13***, 14***, 24***, 34*** | 12**, 24*          | NS            | NS                 | 12*, 13***, 24**, 34*                 |
|          | D | NS                            | 23*                                  | NS                                | NS                 | NS            | NS                 | NS                                    |
|          | P | 34*                           | 23*                                  | NS                                | NS                 | NS            | NS                 | NS                                    |
|          | C | NS                            | NS                                   | NS                                | NS                 | NS            | NS                 | NS                                    |
| I        | S | 12**, 13*, 14**               | 12**, 13**, 14**, 23*, 34*           | 12**, 13*, 14**, 34*, 24*         | 12*                | NS            | NS                 | NS                                    |
|          | D | 12*, 14*, 23*, 34*            | 12*, 14*, 23*, 34*                   | 12*, 13*, 14*, 23*, 34*           | 12*, 14*, 23*, 34* | 12*, 13*, 23* | 12*                | 12*, 14*, 23*, 34*                    |
|          | P | NS                            | 12*, 13*, 23*, 34*                   | NS                                | NS                 | NS            | 12**               | NS                                    |
|          | C | NS                            | 14*(pr)                              | 14*(cc)                           | 12*(cc), 14*(cc)   | NS            | NS                 | NS                                    |
| All      | S | GH**, GI**, HI**              | GH***, GI**                          | GH***, GI***, HI***               | GH*, GI***         | GH*           | HI*                | HI*                                   |
|          | D | NS                            | GH*                                  | GH*, HI*                          | GH*, GI*           | GH*           | GH*, GI*           | NS                                    |
|          | P | NS                            | NS                                   | GI*                               | NS                 | NS            | NS                 | NS                                    |
|          | C | NS                            | HI*(cc)                              | GH*(c sp.), HI*(c sp.)            | GH*(pr)            | NS            | NS                 | GI*(pr), GI*** (c sp.), HI*** (c sp.) |
|          | F | NS                            | NS                                   | NS                                | NS                 | NS            | NS                 | HI*                                   |

Legend: S = soil, D = detritus, P = plants, C = Carabidae, F = frogs, cc = Carabus cancellatus, c sp. = Chlaenius sp., pr = Pseudophonus rufipes. G, H, I = transect name, 1-4 = station, NS not significant, \*, \*\*, \*\*\* significant at  $p < 0.05$ ,  $< 0.01$ , and  $< 0.001$

Our results suggest that the monitoring of the pollution state of large floodplains slightly contaminated with metals, current or restored in the future, would find no much use of mobile groups of organisms as indicators of local, ecosystem or micro landscape level, pollution. This may not be the case in the flood plains of smaller rivers the lower Danube.

## CONCLUSIONS

1. The number of significant differences between the concentration of metals in ecosystems of a micro landscape decreases as one goes with the analyses up in the food chain (from soil to detritus, plants, and Carabidae).
2. There is not a simple pattern of the bioaccumulation of metals in the food chain, but most of the bioconcentration occurred at the level of Carabidae, and not at the level of Rana specie.
3. Mobile groups seem to not be a good indicator of pollution with metals at local level in large flood plains.

**Acknowledgements.** This research was supported by several research projects of the National University Research Council (Romania). We thank to Dr. Dan Cogalniceanu and Dr. Constantin Ciubuc for providing the samples of frogs and beetles.

## REFERENCES

1. Andrews S. M., Johnson M. S., Cooke J. A. - Distribution of trace element pollutants in a contaminated grassland ecosystem established on metalliferous fluorspar tailings. 2: Zinc. *Environmental Pollution*, 1989, 59, 241-252
2. Baca R. M., Threlkeld S.T. - The effects of toxicants on planktonic systems: an integrated approach using the analysis of size distributions. *Journal of Aquatic Ecosystem Stress and Recovery*, 2000. 8, 95-105
3. Bank M. S., Loftin C.S., Jung R.E. - Mercury Bioaccumulation in Northern Two-lined Salamanders from Streams in the Northeastern United States. *Ecotoxicology*, 2005, 14, 181-191
4. Birdsall C. W., Grue C.E., Anderson A. - Lead Concentrations in Bullfrog *Rana Catesbeiana* and Green Frog *R. clamitans* Tadpoles Inhabiting Highway Drainages. *Environmental Pollution (Series A) EPEBD7*, 1986, 40, 233-247
5. Boivin M.E.Y., Greve G.D., García-Meza J.V., Massieux B., Sprenger W., Kraak M.H.S., Breure A.M., Rutgers M. Admiraal W. - Algal-bacterial interactions in metal contaminated floodplain sediments. *Environmental Pollution*, 2007, 45, 884-894
6. Ciubuc C. - Terrestrial macroinvertebrates diversity in Danube wetlands. In „Final Report to European River Margins, ERMAS2-Ro project. IC20-CT96-0010 to ENV4-CT95-0061, EC DG XII”, (Vadineanu A., co-ordinator), 1999.
7. Cogalniceanu D. - Structure and functioning of the amphibian communities in the Small Island of Braila, PhD theses, University of Bucharest, 1997.
8. Cogalniceanu D., Aioanei F., Ciubuc C., Vadineanu A. - Food and feeding habits in a population of common spadefoot toads (*Pelobates fuscus*) from an island in the lower Danube floodplain. *Alytes* 1999, 15, 145-157
9. Cristofor S., Sarbu A., Vadineanu A., Ignat G., Iordache V., Postolache C., Dinu C., Ciubuc C. - Effects of Hydrological Regimen on Riparian Vegetation in the Lower Danube Floodplain. *Internat. Assoc. Danube Res.*, 1997, 32, 233-237
10. Finlayson M. C. - A metal budget for a Monsoonal Wetland in the Northern Australia, pp.433-451, In: *Toxic Metals in Soil Plant Systems*, (S. M. Ross, Ed.), John Willey and Sons, 1994.
11. Griethuysen C. van, Luitwieler M., Joziassse J., Koelmans A. A. - Temporal variation of trace metal geochemistry in floodplain lake sediment subject to dynamic hydrological conditions. *Environmental Pollution*, 2005, 137, 281-294
12. Heikens A., Peijnenburg W.J.G.M., Hendriks A.J. - Bioaccumulation of heavy metals in terrestrial invertebrates. *Environmental Pollution*, 2001, 113, 385-393
13. Hobbelen P.H.F., Koolhaas J.E., van Gestel C.A. M. - Risk assessment of heavy metal pollution for detritivores in floodplain soils in the Biesbosch, The Netherlands, taking bioavailability into account. *Environmental Pollution*, 2004, 129, 409-419
14. Hoffmann G. - Methodenbuch, Band 1, Die Untersuchung von Boden. VDLUFA-Verlag, Darmstadt, 1994.
15. Iordache V., Ruta F. - Wetlands use for the removal of metals from water. *Proceedings of 7<sup>th</sup> International Symposium on "Metal elements in Environment, Medicine and Biology"*, Printing House Eurobit Timișoara, 2006.
16. Iordache V. - Distribution of metals in plants and litter of the Danube floodplain, in press
17. Iordache V., Neagoe A., Cernatoni A., State M., Altorfer T., Preda E., Cristofor S., Vadineanu A. - Heavy metals in the Danube floodplain related to the Yugoslavian conflict. In: *Proceedings of the 5th Symposium on Environmental Contamination*, Prague, Czech Republic, 2000, CD-ROM format, 7 pages
18. Iordache V. - Preliminary assessment of the pollution risk associated with the restoration of the Danube floodplain, *Proceedings of the 5th International Symposium on Metal Elements in Environment, Medicine and Biology*, 211-219, Printing House Eurobit Timișoara, 2002.

19. Iordache V. - The difficulties in assessing the effects of the Yugoslavian conflict in the Lower Danube River System pointed out the need for changing the monitoring system. *J. Env. Protection and Ecology*, 2003a, 4(3), 557-563
20. Iordache V. - The ecotoxicology of heavy metals in the Small Island of Braila, Danube floodplain. PhD thseses, University of Bucharest, Department of Systems Ecology, 2003b.
21. Iordache V., Neagoe Aurora - Mobile organisms integrate fluxes of metals from large areas: consequences for the bioaccumulation in food chains, *Environmental Geochemistry and Health*, in press
22. Iordache V., Cristofor S., Vadineanu A. - Optimization pathways of the decision making process regarding the ecotoxicological effects in the Lower Danube River System. *Proceedings of the "Deltas and Wetlands" International Symposium*, Tulcea, Romania, 2001, 91-97
23. Iordache V., Ignat, G.- Bioaccumulation of metals in aquatic food chains of Danube floodplain, in press
24. Levengood J.M., Skowron L.M. - Use of a Simulated Gizzard to Measure Bioavailability of Metals and Other Elements to Waterfowl. *Ecotoxicology*, 2001, 10, 299-304
25. Lindqvist L., Block M., Tjalve H. - Distribution and excretion of Cd, Hg, methyl-Hg and Zn in the predatory beetle *Pterostichus niger* (Coleoptera: Carabidae). *Environ. Toxicol. Chem.*, 1995, 14, 1195-1201
26. Lock K., Desender K., Janssen C.R. - Effects of metal contamination on the activity and diversity of carabid beetles in an ancient Pb-Zn mining area in Plombieres (Belgium). *Entomol. Exp. Appl.*, 2001, 99, 355-360
27. Loumbourdis N.S., Kostaropoulos I., Theodoropoulou B., Kalmanti D. - Heavy metal accumulation and metallothionein concentration in the frog *Rana ridibunda* after exposure to chromium or a mixture of chromium and cadmium. *Environmental Pollution*, 2007, 145, 787-792
28. Rabitsch W. - Metal accumulation in arthropods near a lead/zinc smelter in Arnoldstein, Austria. *Environmental Pollution*, 1995, 90, 221-237
29. Read H.J., Wheeler C.P., Martin M.H. - Aspects of the ecology of carabidae (coleoptera) from woodlands polluted by heavy metals. *Environmental Pollution*, 1987, 48, 61-76
30. Roe J.H., Hopkins W.A., Jackson B.P. - Species- and stage-specific differences in trace element tissue concentrations in amphibians: implications for the disposal of coal-combustion wastes. *Environmental Pollution*, 2005, 136, 353-363
31. Stone D., Jepson P., Kramarz P., Laskowski R. - Time to death response in carabid beetles exposed to multiple stressors along a gradient of heavy metal pollution. *Environmental Pollution*, 2001, 113, 239-244
32. Vadineanu A., Cristofor, S., Iordache V. - Lower Danube River System biodiversity changes. Pp.29-63, In: B. Gopal, W. J. Junk and J. A. Davis (Eds), *Biodiversity in Wetlands: Assessment, Function and Conservation*, Backhuys Publishers, 2001
33. Wallach R., Grigorin G., Rivlin J. - A comprehensive mathematical model for transport of soil-dissolved chemicals by overland flow, *J. Hydrol.*, 2001, 247/1-2, 85-99
34. Weis J.S., Windham L., Santiago-Bass C., Weis P. - Growth, survival, and metal content of marsh invertebrates fed diets of detritus from *Spartina alterniflora* Loisel and *Phragmites australis* Cav. Trin. ex Steud. from metal-contaminated and clean sites. *Wetlands Ecology and Management*, 2002, 10, 71-84

# ALUMINIUM NEUROTOXICITY

Tibor HALASI<sup>1\*</sup>, Dusan LALOSEVIC<sup>2</sup>, Milka CULIC<sup>3</sup>, Slobodan SEKULIC<sup>2</sup>,  
Ivan CAPO<sup>2</sup>, Gordana GRBIC<sup>3</sup>

1\*.Department of Chemistry, Faculty of Sciences, Trg Dositeja Obradovica 3, RS-21 000 Novi Sad, Serbia, 2. Faculty of Medicine, University of Novi Sad, RS-21 000 Novi Sad, Serbia, 3. Institute for Biological Research, University of Belgrade, RS-11 000 Belgrade, Serbia

\* E-mail : halasi@ih.ns.ac.yu

## ABSTRACT

*In this paper is described the role and consequences of aluminium toxic effects at the nerve cells. Aluminium was investigated as an environmental factor that could trigger inflammatory events in the brain and could enhance existing age-related inflammation. The model of aluminium toxicity was induced by chronic treatment of Wistar rats with aluminum chloride, 1.5% solution instead drinking water for 6 months. Al-treated animals was hypoactive to a catatonic behavior and disrupted exploratory movements. Histological analyze revealed most degenerated neurons, especially in the cerebral cortical V layer and Purkinje cells.*

**Key words:** aluminium, neurotoxicity, rats, models of Alzheimer's disease

## INTRODUCTION

Aluminium is a trivalent cation found in its ionic form in most kinds of animal and plant tissues and in natural waters everywhere. It is the third most prevalent element and the most abundant metal in the earth's crust. Dietary aluminium is ubiquitous, but in such small quantities that it is not a significant source of concern in persons with normal elimination capacity. Aluminium produces toxic effects at the cell membrane by altering the physical properties, interfering with the function of voltage-activated ionic channels and altering the secretion of transmitters (Becaria, 2002). Within the cell, aluminum can affect many key processes in the nucleus, cytoplasm and mitochondria, including glucose metabolism, signal transduction, neurotransmitter synthesis, phosphorylation and dephosphorylation of cytoskeletal proteins, slow axonal transport of neurofilament proteins and inhibition of nucleotide activity (Walton, 1998).

It is estimated that the normal person takes in between 3 and 10 milligrams of aluminum per day. Aluminum is the most abundant metallic element in the earth. It can be absorbed into the body through the digestive tract, the lungs and the skin, and is also absorbed by and accumulates in the body tissues. Aluminum is found naturally in air, water and soil. It is also used in the process of making cooking pots and pans, utensils and foil. Other items such as over the counter pain killers, anti-inflammatory products, and douche preparations can also contain aluminum. Aluminum is also an additive in most baking powders, is used in food processing, and is present in antiperspirants, toothpaste, dental amalgams, bleached flour,

grated cheese, table salt, and beer (especially when the beer is in aluminum cans). The biggest source of aluminum, however, comes from our municipal water supplies.

No known physiologic need exists for aluminum; however, because of its atomic size and electric charge (0.051nm and 3+, respectively), it is sometimes a competitive inhibitor of several essential elements of similar characteristics, such as magnesium (0.066 nm, 2+), calcium (0.099nm, 2+), and iron (0.064nm, 3+). Approximately 95% of an aluminum load becomes bound to transferrin and albumin intravascularly and is then eliminated renally. Aluminum was investigated as an environmental factor that could trigger inflammatory events in the brain and could enhance existing age-related inflammation (Campbell, 2004).

The contribution of aluminum to neurodegenerative diseases such as Alzheimer's diseases (AD), based on the relation of aluminum concentration in drinking water, was indicated in the epidemiological study (Rondeau et al., 2000). Urban water supplies may contain a greater concentration because water is usually treated with the element before becoming part of the supply. Autopsies on many people who have died of Alzheimer's disease showed accumulations of up to four times the normal amount of aluminum in the nerve cells in the brain, especially in the hippocampus which plays a central role in memory. Aluminum causes an oxidative stress within brain tissue. Aluminum excess has a direct effect on hematopoiesis and has been shown to induce anemia. Findings on peripheral smears in patients with aluminum toxicity include *microcytic anemia* (*hypochromic*, *normochromic*), *anisocytosis*, *poikilocytosis*, *chromophilic* cells, and basophilic stippling. Note that these are the same findings observed in patients with lead poisoning.

Aluminum can also be found in bone marrow of *macrophages*. Since the elimination half life of aluminum from the human brain is 7 years, this can result in cumulative damage via the element's interference with neurofilament axonal transport and neurofilament assembly. Some experts believe it plays a role in leading to the formation of Alzheimer like *neurofibrillary* tangles. Aluminum is absorbed from the GI tract in the form of oral phosphate-binding agents (aluminum hydroxide), parenterally via immunizations, via dialysate or total parenteral nutrition (TPN) contamination, via the urinary mucosa through bladder irrigation, and transdermally in antiperspirants. Lactate, citrate, and ascorbate all facilitate GI absorption. If a significant load exceeds the body's excretory capacity, the excess is deposited in various tissues, including bone, brain, liver, heart, spleen, and muscle. This accumulation causes morbidity and mortality through various mechanisms.

Toxic effects are dependent upon the amount of metal ingested, entry rate, tissue distribution, concentration achieved, and excretion rate. Mechanisms of toxicity include inhibition of enzyme activity and protein synthesis, alterations in nucleic acid function, and changes in cell membrane permeability. Aluminum toxicity is usually found in patients with renal impairment. Acute intoxication is extremely rare; however, in persons in whom aluminum clearance is impaired, it can be a source of significant toxicity.

### **Pathophysiology of Aluminium toxicity**

Aluminum toxicity was originally described in the mid-to-late 1970's in a series of patients in Newcastle, England, through an associated *osteomalacic* dialysis *osteodystrophy* that appeared to reverse itself upon changing of the dialysate water to deionized water (ie, aluminum-depleted water). Previously, the only known dialysis-associated bone disease was *osteitis fibrosa cystica*, which was the result of abnormalities in vitamin D production that resulted in a secondary

hyperparathyroidism, increased bone turnover, and subsequent peritrabecular fibrosis. In aluminum-related bone disease, the predominant features are defective mineralization and osteomalacia that result from excessive deposits at the site of osteoid mineralization, where calcium would normally be placed.

Aluminum also has a direct effect on *hematopoiesis*. Excess aluminum has been shown to induce *microcytic anemia*. Daily injections of aluminum into rabbits produced severe anemia within 2-3 weeks. The findings were very similar to those found in patients suffering from lead poisoning. Aluminum may cause anemia through decreased heme synthesis, decreased globulin synthesis, and increased *hemolysis*. Aluminum may also have a direct effect on iron metabolism: it influences absorption of iron via the intestine, it hinders iron's transport in the serum, and it displaces iron's binding to transferrin. Patients with anemia from aluminum toxicity often have increased reticulocyte counts, decreased mean corpuscular volume, and mean corpuscular hemoglobin. Other organic manifestations of aluminum intoxication have been proposed, such as a slightly poorer immunologic response to infection, but the mechanism by which it exerts its effect is complex and multifactorial.

When there is a high level of absorption of aluminum and silicon, the combination can result in an accumulation of certain compounds in the cerebral cortex and can prevent nerve impulses being carried to and from the brain properly. Long term calcium deficiency can further aggravate the condition. Workers in aluminum smelting plants on a long term basis, have been known to experience dizziness, poor coordination, balance problems and tiredness. It has been claimed that the accumulation of aluminum in the brain could be a possible cause for these issues.

## MATERIALS AND METHODS

The model of aluminum toxicity (Culic et al., 2007) was induced by chronic treatment of *Wistar* rats with  $\text{AlCl}_3$  for four weeks (2mg Al/kg, i. p. daily). The treated animals were compared to the controls. For histological examination the brain tissue was fixed in 10% formaline immediately after euthanasia of animals. Tissue was routinely included in paraffin wax, cutting and staining by Walton procedure. Specific neurodegeneration was noted under UV light at fluorescent microscope Leica.

## RESULTS AND DISCUSSIONS

In aluminium-treated animals by parenteral administration of aluminum chloride (2mg/kg for 28 days) we noted focal neurodegeneration like dark-stained neurons, particularly in the cerebral cortex. On the Walton staining those neurons were intensive red and under UV light showed brightly fluorescence (figure 1).

Mechanisms underlying aluminum neurotoxicity are not completely resolved. For both aluminum treated rats (intracerebroventricular  $\text{AlCl}_3$  injections for 5 days) and saline-infused controls, no major signs of great histological changes were evident in cresyl violet-stained sections (Platt et al., 2001). Platt and her colleagues found by fluorescent Morin staining that Al was found in white matter of striatum, corpus callosum and cingulate bundle. Immunoreactivity of astrocytes and phagocytic microglia based on glial fibrillary acidic protein and ED1 markers, revealed a greater inflammatory response in Al-injected animals compared to controls, while damage of the cingulate bundle in Al-treated animals led to a severe

anterograde degeneration of cholinergic terminals in cortex and hippocampus, as indicating by acetylcholinesterase labeling.




**Fig 1.** Dark neurons in the lamina V of rat cerebral cortex, Al-treated rat (Walton stain, white and UV light, magnified 1000)

Al-maltolate treated rabbits developed progressive behavioral symptoms consisting of forward head tilting tremor following touch or passive movement of the extremities, hemiplegic gait and splaying of the extremities (Yokel, 1989). Al-gluconat treated rats (0.85 mg/kg i.p., three times a week, 6 months) also developed impaired spatial memory deficits, hypoactive to a catatonic behavior and disrupted exploratory movements (Miu and Andreescu, 2003). Aluminum induced learning and memory impairment model (Sun et al., 2005) was established by gavage of  $\text{AlCl}_3$  (600mg/kg for 90 days) and in the hidden platform trials, escape latencies of the naloxone-treated rats (naloxone 0.8mg/kg, i.p. daily for 7 days) and the number of entries in the target area were significantly shorter than that of the non-treated model rats. We did not observe any changes in motor control, like paresis or tremor, in the Al treated animals. There has been a certain progress in investigation of animal models of AD (Price and Sisodia, 1994) as AD is a most common dementing disorder of late life comprising complex neurobiochemical and neuropathological events characterized by extracellular deposition of  $\beta$ -Amyloid proteins, formation of intraneuronal neurofibrillary tangles and selective neuronal loss.

Aluminum is neurotoxic in both experimental animals and certain human conditions but whether aluminum presents a health hazard to humans as a contributing factor to Alzheimer's disease is still for debate (Savory et al., 1996). There are special heterozygous APP23 mice (Van Dam et al., 2003) and single, double and triple transgenic mice (Bloom et al., 2005) as models for human neurodegenerative disorders like AD. While Al still remains a mystery, Al-maltolate treated aged rabbits are suggested as a reliable and sensitive animal model for understanding AD neuropathology (Bharathi et al., 2006). Besides a great number of animal models, the techniques of computational simulation have begun to be applied to modeling neurodegenerative disease (including the model of Alzheimer dementia), focusing on attractor-based network dynamics and on linking sequences of attractor states and their role in cognition (Frinkel, 2000).



Treatment of aluminum toxicity includes elimination of aluminum from the diet, TPN, dialysate, medications, antiperspirants, and an attempt at the elimination and chelation of the element from the body's stores. Avoidance of aluminum is easily achieved once the need to do so is recognized. Educate pregnant and breastfeeding females, and any patient with compromised renal function, about the use of aluminum-containing antacids and the potential dangers of their use and overuse. A safe alternative includes calcium carbonate, such as found in Tums. Educate patients to refrain from driving or operating hazardous machinery if they develop dizziness or impaired vision or hearing during treatment. Eat a diet that is high in fiber and includes apple pectin.

Use stainless steel, glass, or iron cookware. Stainless steel is the best choice. Beware of any product containing aluminum or dihydroxyaluminum. A hair analysis can be used to determine levels of aluminum in the body. Research has shown that the longer you cook food in aluminum pots, the more they corrode, and the more aluminum is absorbed into the food and hence into the body. Aluminum is more readily dissolved by acid forming foods, such as coffee, cheese, meat, black and green tea, cabbage, cucumbers, tomatoes, turnips, spinach and radishes. Acid rain leeches aluminum out of the soil and into drinking water.

Depending upon the degree of dementia and overall medical frailty of the patient, most improve in with deferoxamine therapy. Some patients, however, succumb to their underlying disease processes before any noticeable improvement in mental status or anemia occurs. Whether aluminum toxicity itself is fatal is unknown. Typically, patients' underlying diseases and medical frailty lead to early morbidity and mortality. Treatment of aluminum toxicity includes elimination of aluminum from the diet, TPN, dialysate, medications, antiperspirants, and an attempt at the elimination and chelation of the element from the body's stores. Avoidance of aluminum is easily achieved once the need to do so is recognized. Elimination is accomplished through the administration of deferoxamine through any of several routes. Deferoxamine, the metal-free ligand of the iron-chelate isolated from the bacterium *Streptomyces pilosus*, is used for acute and chronic iron toxicity and aluminum toxicity. It has a high affinity for ferric iron and does not affect iron in hemoglobin or cytochromes. Educate pregnant and breastfeeding females, and any patient with compromised renal function, about the use of aluminum-containing antacids and the potential dangers of their use and overuse. A safe alternative includes calcium carbonate, such as found in Tums. Educate patients to refrain from driving or operating hazardous machinery if they develop dizziness or impaired vision or hearing during treatment. Eat a diet that is high in fiber and includes apple pectin.

## CONCLUSIONS

None of the models available mimics exactly cognitive, behavioural, electrophysiological, biochemical and histopathological abnormalities observed in neurological disorders characterised by cognitive impairment. Partial reproduction of neuropathology and/or cognitive deficits of Alzheimer's disease, vascular dementia and dementia occurring in Huntington's and Parkinson's diseases, or in other neurodegenerative disorders may represent a basis for understanding pathophysiological traits of these diseases and for contributing to their treatments (Tayebati, 2006). Hopefully, different methodological approaches in investigation of various models will contribute to delineate pathogenic mechanisms, early diagnostics and test new therapies for treating AD in humans (Culic et al., 2007)

**Acknowledgement.** The authors wish to thank Ministry Science of the Republic of Serbia for financial support this work with Project: European Dimension of Change of the Education System in Serbia, No. 149009 (2006-2010).

## REFERENCES

1. Becaria A., Campbell A., Bondy S.C. : Aluminum as a toxicant. *Toxicology and Industrial Health*, 2002, 18 ,309-320.
2. Bloom F.E., Reilly J.F., Redwine J.M., Wu C.C. : Mouse models of human neurogenerative disorders, *Arch. Neurol.*, 2005, 62,185-187.
3. Campbell A. : Inflammation, Neurodegenerative Diseases and Environmental Exposures. *Ann. NY Acad. Sci.*, 2004,1035,117-132.
4. Culic M., Maratac Lj., Grbic G., Kesic Sr., Spasic S.I., Sekulic S.I., Lalosevic D., Capo I. : Aluminum toxicity in rat Brain: Electrophysiological and behavioral evidence, In: *From Basic Motor Control to Functional Recovery*, Ed. N. 2007.
5. Miu A.C., Andreescu C.F. (2., Ganchev, Marin Drinov Academic Publ. House, Sofia, p. 224-230.003) : A behavioral and histological study of the effects of long term exposure of adult rats to aluminum. *Intern. J. Neuroscience*, 113:1197-1211.
6. Platt B., Fidler G., Riedel G., Henderson Z. : Aluminium toxicity in the rat brain: histochemical and immunocytochemical evidence. *Brain Res. Bull.*, 2001, 55, 257-267.
7. Price D. I., Sisodia S.S. : Celular and molecular biology of Alzheimer's disease and animal models. *Ann. Rev. Med.*, 1994, 45,435-446.
8. Rondeau V., Commenges D., Jacqmin-Gadda H., Dartigues J. : Relation between aluminum concentration in drinkong water and Alzheimer's disease: An 8 year follow-up study. *Am. J. Epidemiol.*, 2000, 152,59-66.
9. Savory, J., Exley, C., Forbes., W. F., Huang, J. G., McLachlan, D. R. C., Wakayama I. : Can the controversy of the role of aluminium in Alzheimer's disease be resolved? What are the suggested approaches to this controversy and methodological issues to be considered. *J. Toxicol. Environ. Health*, 1996, 48, 615-636.
10. Spasic S., Kalauzi A., Culic M., Grbic G., Martac Lj. : Fractal analysis of rat brain activity after injury. *Med. Biol. Eng. Comput.*, 2005, 43, 345-348.
11. Sun S., Hua L., Yun Z., Dong H. : Effect of maloxone on aluminium-induced learning and memory impairment in rats. *Neurology India*, 2005, 53, 79-82.
12. Tayebati S.: Animal models of cognitive dysfunction. *Mech. Ageing Dev.* 2006, 127, 100-108.
13. Van Dem D., D'Hooge R., Staufenbiel M., Van Ginneken C., Van Meir F., De P. P. : Agendependent cognitive decline in the APP23 model precedes amyloid deposition. *Eur. J. Neurosci.*, 2003, 17, 388-396.
14. Walton J. (1998) - Absorbtion of aluminium and its effect on brain cells. In: *Aluminium: report of an International meeting*, Brisbane, 20-21 April, 1995, Eds. P. Imray, M. R. Moore, P. W. Callan, W. Lock, Rundle Mall-Public and Environmental Health Service, p. 22-34.
15. Yokel R.A. : Aluminium produces age related behavioral toxicity in rabbit, *Neurotoxicol Teratol*, 1989, 11, 151-242.

# CORRECTION OF IODINE DEFICIENCY BY AN IODIZED VEGETABLE OIL – FLORIOL IODINE PRODUCT – (PARTIAL RESULTS)

Mihaela SIMESCU <sup>1\*</sup>, Bruno DE BENOIST <sup>2</sup>, Mariana SAVA <sup>1</sup>,  
Claudia PODIA-IGNA<sup>3</sup>, Francois DELANGE <sup>4</sup>

*1\** Institute of Endocrinology, B-dul Aviatorilor 34-36, Bucharest, RO-011 863, Roumania; *2* Bunge, 13, Rue de Florissant, Geneve, Switzerland.; *3*. Private Clinic, Sibiu, Roumania; *4*. Department of Pediatrics, University Hospital Saint Pierre, Brussels, Belgium

\* E-mail : mihaelasimescu@yahoo.com

## ABSTRACT

*An iodine deficit in a population may delay the development of the brain of fetus and children with irreversible mental retardation. It was reported that still 33% of world population is affected by iodine deficiency. The aim of the study: to estimate if oral iodized oil is a major alternative or complement to iodized salt for correcting iodine deficiency in a population; to follow-up the effect of low doses of iodized vegetable oil (IVO) on urinary excretion of iodine (UIC) in studied population of 2 kind of IVO: 1st enriched with Lipiodol (IVOL); 2nd enriched with Brassiodol (IVOB). 3 population groups were randomly selected in Bucharest, an area where the median UIC is 128 µg/l: 1st (n=34) received 60 µg iodine/day in 1 ml IVOL in dairy; 2nd (n=40) received 60 µg iodine in 1 ml of IVOB in dairy; 3rd group (n=34) received 50 µg iodine per day IVOB in cooking meals. Anthropometric measurements were made. Serial UIC level for each subject were done 1 month before IVO adm -12 samples and 1 month during IVO adm -12 samples. The administration of daily low doses of IVO is followed by an increase of UIC ( $p < 0.001$ ). There were no significant statistical differences between UIC of the 2 groups that received IVO enriched with 2 different vehicles of iodination ( $p = 0.24$ ) and also between the 2 forms of administration (dairy products and cooking meals;  $p = 0.55$ ). IVO constitutes an additional measure to iodized salt for iodine deficit prophylaxis in the future.*

**Key words:** iodine deficiency, iodized vegetable oil, iodine deficit prophylaxis

## INTRODUCTION

An iodine deficiency in the diet of a population may entail not only a change of the thyroid function in all age groups but also a delay in the development of the brain of fetus and children, whose clinical demonstration will be an irreversible mental retardation. This complication can only be avoided if a correction of the iodine deficiency is made very early during the intellectual development, which means during pregnancy and childhood (Delange, 1994).

In 2003, 2 billion inhabitants of 136 countries of the world were affected by iodine deficiency that is 33 % of the world population. It seemed consequently that iodine deficiency was the main cause of potentially avoidable mental retardation (Dr. de Benoist, personal communication).

---

„Metal Elements in Environment, Medicine and Biology”, Tome VIII, pp.87-94, Publishing House “Eurobit” Timișoara, 2008

Proc. of 8<sup>th</sup> Int. Symp. of Roumanian Academy - Branch Cluj-Napoca and Branch Timișoara, 2008, December 5-6, Timișoara, Roumania

At the general assembly of the World Health Organization (WHO) and the World Summit for Children in 1990, it was agreed that the most effective strategy aiming at the long-term elimination of iodine deficiency in the world was the establishment of programs of universal salt iodization, that is the enrichment in iodine of salt intended for food, feed and for the food industry (WHO, UNICEF and ICCIDD, 1994; Mannar and Dunn, 1995).

The salt iodization campaigns had a remarkable effect on the progressive elimination of iodine deficiency in the world and constitute an unprecedented success of public health in the field of the micronutrient deficiencies. Indeed, the household consumption of iodized salt in deficient countries had increased from less than 10 % in 1990 to 67 % in 1999, with a remarkable figure of 90 % in Latin America (WHO, UNICEF and ICCIDD, 1999; Delange et al, 2001).

However, a reassessment of the iodine nutrition in the world in 2005 showed that 54 countries were always affected by iodine deficiency, in particular in Central and Western Europe (Andersson et al, 2005). It was thus suggested strengthening the establishment of the salt iodization programs in these countries while organizing in parallel a systematic iodine supplementation for pregnant and lactating women and young children. An additional possibility is the introduction in these countries of iodine fortified food others than salt (de Benoist and Delange).

Romania is one of the countries of Central Europe which has suffered a lot due to iodine deficiency (Simescu et al, 2002). Until 2002, a national voluntary salt iodization program brought a partial correction to this deficiency but collided with a series of difficulties of logistic order (Dunn, 1987). In 2002, the Romanian government adopted a Decision on compulsory addition of salt in iodine (Decision 568 of 5 June 2002). This resulted in 2005-2006 in a correction of iodine deficiency in almost all the districts where the iodine status was estimated (Dr. Simescu, personal communication), but monitoring the iodine uptake in pregnant women obtained by determination of urinary iodine concentration in a representative group showed a persistence of low values (median UIC 55 µg/l in 2002, 55 µg/l in 2005 and 80 µg/l in 2006-2007). Iodine deficiency in pregnant women represents an emergent health problem.

The aim of the study: to estimate additional contribution to the control of iodine deficiency resulting from the iodine fortification of another basic food of the population namely sunflower oil; to follow-up the effect of low doses of iodized vegetable oil (IVO) on urinary excretion of iodine (UIC) of 2 kind of IVO: first enriched with Lipiodol (poppy seed oil as vehicle of iodination); second enriched with Brassiodol (utilizes rapeseed oil as vehicle of iodination).

## MATERIALS AND METHODS

The iodized sunflower oil is produced from sunflower (*Heliantus annuus* L.) seed. It is free from any contaminants or toxic compounds and fulfill the requirements and regulations of National Food Law (HG 150/2004, HG 412/2004, HG 924/2005, HG 1198/2002) ) and National Oil Standards ( Ordin 454/917/2001-22/2002, Ordin 495/997/2002-10/2003). The product contains iodized poppy seed oil in form of fatty acid ethylesters in order to provide the requested iodine content.

In this study, low daily intakes of iodized oil (organic iodine) were administered to studied population with a recent correction of their iodine deficiency.

Two groups of schoolchildren aged 6-14 y.o. and 1 group of young adults 24-40 y.o. were randomly selected in Bucharest, an area where the median UIC is 128 µg/l: first group of schoolchildren (n=34) received a complement of 60 µg iodine/day in 1 ml IVO (enriched with Lipiodol) mixed in dairy dessert; second group of young adult (n=40) received a complement of 60 µg iodine in 1 ml of IVO enriched with Brassiodol, mixed in dairy dessert; third group of schoolchildren (n=32) received 50 µg iodine per day from IVO enriched with Brassiodol used in cooking meals. At the beginning of the study anthropometric measurement were registered. Serial determinations of the UIC level for each child, as an index of the effect of iodine supplementation and an estimation of the iodine stored in the body, were done 3 times/week, for 4 weeks (12 samples), before IVO administration and during 4 weeks (12 samples) under administration of vegetable oil. Urine samples were also taken twice a month during 6 months after administration of iodized oil in 2 groups of schoolchildren (the study is ongoing).

The samples of urine were taken in the morning at getting up time with the aim to reduce the variability of the urinary iodine due to the volume and were kept at -20°C without preservative. For determination of UIC we used Sandell and Kolthoff method.

Statistical analysis of the results was made using SPSS program.

## RESULTS AND DISCUSSIONS

The data of the first group of schoolchildren which received IVO enriched with Lipiodol showed an increase of UIC on the second day after administration and the persistence of significant increased values during the month of administration ( $p < 0,001$ ). Elimination of UIC after a month of IVO administration showed still a significant high median value (median UIC = 177.5 µg/l,  $p=0,14$ ). During the 3 to 6 months after IVO administration median UIC came back to values registered before IVO administration (table 4 and fig. 1).

The median UIC before IVO administration was 140 µg/l (table 2) and during IVO administration 192.5 µg/l (table 3). Median UIC of serial determination in first group of children before IVO administration are showed in table 1.

**Table 1:** Serial determination of urinary iodine concentration in first group of children in first month before iodized vegetable oil administration

| Date       | Mean UIC (µg/l) | Median UIC (µg/l) |
|------------|-----------------|-------------------|
| 23.04.2007 | 166             | 150               |
| 26.04.2007 | 130.3           | 122.5             |
| 28.04.2007 | 139             | 127.5             |
| 30.04.2007 | 132.8           | 120               |
| 02.05.2007 | 128.4           | 115               |
| 04.05.2007 | 121.3           | 113.8             |
| 07.05.2007 | 126.9           | 116.3             |
| 09.05.2007 | 171.8           | 152.5             |
| 11.05.2007 | 159.7           | 150               |
| 14.05.2007 | 131.3           | 130               |
| 16.05.2007 | 159.6           | 150               |
| 18.05.2007 | 164.3           | 167.5             |

**Table 2:** UIC value before IVO administration in first group children


| No of samples | Mean UIC (µg/l) | Median UIC (µg/l) | SD (µg/l) |
|---------------|-----------------|-------------------|-----------|
| 347           | 144             | 140               | 39.8      |

**Table 3:** UIC value during IVO administration in first group children

| No of samples | Mean UIC (µg/l) | Median UIC (µg/l) | SD (µg/l) |
|---------------|-----------------|-------------------|-----------|
| 345           | 196.1           | 192.5             | 38.2      |

**Table 4:** UIC value 6 month after IVO administration in first group children

| Month after IVO administration | No of samples | Mean UIC (µg/l) | Median UIC (µg/l) | SD (µg/l) |
|--------------------------------|---------------|-----------------|-------------------|-----------|
| 1 <sup>st</sup> month          | 42            | 169             | 177.5             | 44.6      |
| 2 <sup>nd</sup> month          | 56            | 147.3           | 140               | 29.3      |
| 3 <sup>rd</sup> month          | 57            | 136.8           | 127.5             | 24.8      |
| 4 <sup>th</sup> month          | 54            | 14.2            | 135               | 26        |
| 5 <sup>th</sup> month          | 54            | 134.7           | 125               | 24.5      |
| 6 <sup>th</sup> month          | 26            | 139.3           | 135               | 25.7      |

**Fig.1:** Evolution of UIC in first studied group

In second group of young adults who received IVO enriched with Brassiodol in amount of 60 µg iodine mixed in dairy products, median UIC before IVO

administration was 135 µg/l (table 5) and during IVO administration 192.5 µg/l (table 6).

**Table 5:** UIC value before IVO administration in second group of young adults

| No of samples | Mean UIC (µg/l) | Median UIC (µg/l) | SD (µg/l) |
|---------------|-----------------|-------------------|-----------|
| 107           | 134.50          | 135.00            | 12.7      |

**Table 6:** UIC value during IVO administration in second group of young adults

| No of samples | Mean UIC (µg/l) | Median UIC (µg/l) | SD (µg/l) |
|---------------|-----------------|-------------------|-----------|
| 220           | 190.30          | 192.50            | 7.30      |

In third group of schoolchildren who received IVO enriched with Brassiodol in amount of 50 µg iodine per day mixed in cooking meals, median UIC before IVO administration was 131.25 µg/l (table 7) and during IVO administration 195 µg/l (table 8). The median UIC during IVO administration is significantly higher than basal values ( $p < 0,001$ ). The follow-up 5 months after IVO administration revealed the persistence of high median UIC at 4 months ( $p=0,03$  at 4 months). The values of UIC 5 months after IVO administration are showed in table 9 and fig. 2.

**Table 7:** UIC value before IVO administration in third group of schoolchildren

| No of samples | Mean UIC (µg/l) | Median UIC (µg/l) | SD (µg/l) |
|---------------|-----------------|-------------------|-----------|
| 68            | 136.65          | 131.25            | 25.17     |

**Table 8:** UIC value during IVO administration in third group of schoolchildren

| No of samples | Mean UIC (µg/l) | Median UIC (µg/l) | SD (µg/l) |
|---------------|-----------------|-------------------|-----------|
| 377           | 193.04          | 195.00            | 18.24     |

**Table 9:** UIC value 5 months after IVO administration in third group of schoolchildren

| No of samples | Mean UIC (µg/l) | Median UIC (µg/l) | SD (µg/l) |
|---------------|-----------------|-------------------|-----------|
| 41            | 193.11          | 195.00            | 8.33      |
| 47            | 186.76          | 187.50            | 6.73      |
| 54            | 175.42          | 175.00            | 6.57      |
| 50            | 167.20          | 167.50            | 6.86      |
| 52            | 143.1           | 142.5             | 10.3      |




**Fig.2.** Evolution of UIC in 3<sup>rd</sup> studied group

**Table 10:** Median urinary iodine concentration and SD in 3 studied groups

| No. of subjects | Iodine source       | Iodine dosage | Baseline 4 weeks |       | During daily oral FIP 4 weeks |       |
|-----------------|---------------------|---------------|------------------|-------|-------------------------------|-------|
|                 |                     |               | Median UIC(µg/l) | SD    | Median UIC(µg/l)              | SD    |
| 34              | I. FIP Lipiodol     | 60 µg/day     | 140              | 39.8  | 192.5                         | 38.2  |
| 40              | II. FIP Brassiodol  | 60 µg/day     | 135              | 12.7  | 192.5                         | 7.3   |
| 34              | III. FIP Brassiodol | 50 µg/day     | 131.25           | 25.17 | 195                           | 18.24 |

The samples of UIC during the month preceding the administration of IVO reflect environmental availability of iodine. The results showed in table1, 2, 5 and 7 are concordant with previous reported data regarding the iodine uptake availability in Bucharest population.

UIC measured during the month of IVOL and IVOB administration revealed early increased values – first day after onset of administration – and the persistence of I127 high output during the month of administration.

After IVOL administration we observed the persistence of significant values of iodine concentration in urine only in the first month. We didn't observe significant high values in the next 5 months.

After IVOB administration we observed a prolonged elimination of iodine in urine. Median UIC in 1<sup>st</sup>, 2<sup>nd</sup> and 3<sup>rd</sup> and 4<sup>th</sup> month after IVOB administration showed significant high values.

The administration of daily low doses of IVO is followed by an increase of UIC ( $p < 0.001$ ). There were no significant statistical differences between UIC of the 2 groups that received IVO enriched with 2 different vehicles of iodination ( $p = 0.24$ ) and also between the 2 forms of administration (dairy products and cooking meals;  $p = 0.55$ ).



IVO constitutes an important additional measure for iodine deficit prophylaxis in the future with beneficial effects especially in population groups which need iodine supplementation (e.g. pregnant women) and in population groups with salt intake restriction.

## CONCLUSIONS

1. The administration of daily low doses of IVO is followed by an increase of UIC ( $p < 0.001$ ).
2. There were no significant statistical differences between the 2 groups that received IVO enriched with 2 different vehicles of iodination ( $p = 0.24$ ) and also between the 2 forms of administration (dairy products and cooking meals;  $p = 0.55$ ).

## REFERENCES

1. Andersson M., Takkouche B., Egli I., Allen H.E., de Benoist B. - Current global iodine status and progress over the last decade towards the elimination of iodine deficiency, Bull WHO, 83, 2005
2. Anonymous - Iodized water to eliminate iodine deficiency, IDD Newsletter, 13, 1997, 33-39.
3. de Benoist B., Delange F. - Iodine supplementation of pregnant and lactating women and children under 2 years, Public Health Nutrition, in press.
4. Delange F. - The disorders induced by iodine deficiency, Thyroid, 1994, 4, 107-128.
5. Delange F., de Benoist B., Pretell E., Dunn J. - Iodine deficiency in the world: where do we stand at the turn of the century, Thyroid, 2001, 11, 437-447.
6. Dunn J.T. - Iodized oil in the treatment and prophylaxis of IDD, pp127-134, In: The prevention and control of iodine deficiency disorders (eds Hetzel B.S., Dunn J.T., Stanbury J.B.), Amsterdam, Elsevier, 1987
7. Elnagar B., Eltom M., Karlsson F.A., Ermans A.M., Gebre-Medhin M., Bourdoux P.P. - The effects of different doses of oral iodized oil on goiter size, urinary iodine, and thyroid-related hormones, J. Clinical Endocrinology Metab, 1995, 80, 891-897.
8. Furnee C.A., Pfann G.A., West C.E., van der Haar F., van der Heide D. - New model for describing urinary iodine excretion its use for comparing different oral preparations of iodized oil, Am J. Clin Nutr, 1995, 61; 1257-1262.
9. Ingenbleek Y., Jung L., Ferard G., - Brassiodol: A New Iodised Oil for Eradicating Endemic Goiter, The Journal of Trace Elements in Experimental Medicine 13:85-96, 2000.
10. Mannar V.M.G., Dunn J.T. - Salt iodization for the elimination of iodine deficiency., MI, ICCIDD, UNICEF, WHO Geneva, 1995
11. Raclot T., Mioskowski E., Bach A.C., Groscolas R. - Selectivity of fatty acid mobilization; a general metabolic feature of adipose tissue, Am J. Physiol, 1995, 269, R1060-R1067.
12. Simescu M., Popescu R., Ionitiu D., Zbranca E., Grecu E., Marinescu E., Tintea L., Nicolaescu E., Purice M., Popa M. and Gutekunst R. - The status of iodine nutrition in Romania, pp 383-388, In : Iodine deficiency in Europe. A Continuing Concern. (eds Delange F., Dunn J.T., Glinioer D.), Plenum Press publ, 1993.

13. Simescu et al. - Iodized oil as a complement to iodized salt in schoolchildren in endemic goiter in Romania, *Horm. Res.*, 2002, 58, 78-82.
14. Simescu et al. - ICCIDD/ETA Meeting, Naples.
15. Snedecor G.W., Cochran W.G. - *Statistical methods*, 6th ed. Ames, Iowa; Iowa State University Press, 1967.
16. Wei J., Li J. - Metabolism of iodized oil after oral administration in guinea pigs, *Nutr Rep Int*, 1985, 31, 1085-1092.
20. Zimmermann M.B., Adou P., Torresani T., Zeder C., Hurrell R.F. - Effect of oral iodized oil on thyroid size and thyroid hormone metabolism in children with concurrent selenium and iodine deficiency, *EJCN*, 2000, 54, 209-213.
21. Zimmermann M.B., Wegmueller R., Zeder C., Torresani T., Chaouki N. - Rapid relapse of thyroid dysfunction and goiter in school-age children after discontinuation of salt iodization, *Am J Clin Nutr*, 2004, 79, 642-645.
22. \*\*\*WHO, UNICEF and ICCIDD: Indicators for assessing Iodine Deficiency Disorders and their control through salt iodization, WHO publ. WHO/NUT/94.6, 1-55, 1994.
23. \*\*\*WHO, UNICEF and ICCIDD - Progress towards the elimination of Iodine Deficiency Disorders (IDD). WHO publ. WHO/NHD/99.4, 1-33, 1999

# MINERAL IMBALANCE – REFLECTION TO BLOOD VESSELS

Tatjana LALIC <sup>1\*</sup>, Ivana DJUJIC <sup>2</sup>

*1\* Institute of Chemistry, Technology and Metalurgy - Science and Technology Park, RS-11080 Belgrade, Serbia; 2. Department of Chemistry, Institute of Chemistry, Technology and Metalurgy, University of Belgrade, Njegoseva 12, RS-11080 Belgrade, Serbia*

\* E-mail : tatjanalali@yahoo.com

## ABSTRACT

The purpose of this study was to research reflection of disturbances in chemical elements content to health status of blood vessels. In this study, we used two non invasive diagnostic methods particularly efficient in early diagnostic of different pre-pathological disturbances and pre clinical stage of disease development, when disease symptoms are not expressed or absent: Bio-resonance method and elemental hair analyze. Examination were conducted on 103 subjects (67 females and 36 males) aged 12-86 years.

In the largest number of subjects, early forms of abnormalities were discovered in functioning of myocardium (58%); abdominal and thoracic aorta (63%); arteries coronaries (66%); brain arterial circulation (69%); venous circulation (75%); arterial blood pressure (50% hypertension, 15% hypotension, 45% variable), pancreas (69%). These subjects were without symptoms, detectable disorders in total plasma Mg concentration and dysfunction of tested organs by classical diagnostic methods. Only Mg concentration in their hair was higher than normal. With these very comfortable methods for patients, we can very early detect pre pathological disturbances and pre clinical stage in health status. Also we can design nutrition program for specific person and follow results.

**Key words:** hair, elements, imbalance, blood vessel, bio-resonance,

## INTRODUCTION

It is now clear that disturbances in trace elements are associated with vascular diseases risk. Conducted studies on vascular diseases shows that they are the first cause of morbidity and mortality in our and many Western countries.

The main risk factors for vascular diseases, hypertension, dyslipidemia, diabetes and obesity related to diet. Conducted studies shows that the great variation in vascular disease death rates among various nations exists. Other studies shows that between death rates of venous diseases, cholesterol and other risk factors on one side and the dietary intake of 34 elements exist relationships (Yacowitz et al., 1965; Klevay, 1977b). Among the conducted study most related to imbalances in Ca, Mg, K, Na, Zn, Se, Cu, Li, Mo, V, Cr, Si, Cd, Pb and Sn.

The mechanisms by which particular elements or their compounds may affect vascular disease risk are not clear, but it is likely that they involve effects on enzymes, hormones, and messenger molecules. Observations of known enzymes

showed that about 30% enzymes contain mineral elements and/or require minerals for activity.

The vascular diseases are the first cause of morbidity and mortality in Serbia and Western populations. Researches conducted in Serbia showed that: 95% of the population takes in about 40% of the RDA for selenium (Se); 60% females take in less iron (Fe) than is RDA; 85% of the population take in from 75-85% of the RDA for zinc (Zn); 15% have a lower intake of copper (Cu) and chromium (Cr) than is the RDA (Djujić et al., 2000). Hair analyses of toxic elements, on the other side indicates that contamination with some of toxic elements as is As richer in males the maximal clinical reference level for biomonitoring of 0,1 mg/kg. Comparison of smokers with non-smokers showed that they have higher concentrations of Pb, Cd and Al in hair (Jeremic et al., 2004).

The relationship between hair element concentrations and human health is a complex process related to exposure, absorption, and tissue distribution of essential and toxic elements. Studies correlate elements in hair with exposure, disease and physiologic or pathologic effects of nutritional excesses or deficiencies. A long-term record that provides hair analysis reflects normal and abnormal metabolism, assimilation and exposure (Suzuki, 1982; Airey, 1983).

Hair analysis is useful for biological monitoring of most of the toxic metals and as well as a prognostic tool to ascertain whether an individual has a specific biochemical uniqueness, which can then be addressed in a therapeutic or prophylactic program. Levels of elements in hair are often even up to 300 times higher than those in serum or urine. As such hair is sensitive barometer of early imbalances and often reflects them before symptoms appear.

The BIO-RESONANCA is one of the most promising non-invasive diagnostic and therapeutic method in the field of energy-information, medicine. It is based on the conception of integrity of human organism and as such offers possibilities for sophisticated assessments.

For diagnostic and activation natural mechanisms, directed onto restoring and maintenance of healthy conditions it uses processes of energy – information interchange that are a base of the functioning organism. Examinations conducted by the practitioners that use Bio-resonant diagnostic showed that such one screening test provides very useful information's that help in identifying the underlying root of patient's complaint (Gotovski et al., 2000; Djujić and Lalic., 2006; Lalic and Djujić, 2008).

The aim of the present study was to evaluate interrelations between blood vessels health status markers obtained by Bio-resonance diagnostic and concentrations of essential and toxic elements in hair of subjects from Serbia.

## **MATERIALS AND METHODS**

The studied population were 103 subjects (66 females and 37 males) aged 12-86 years from various urban residential regions of Serbia, aged 8-78 years. Volunteers were, according to their ages, divided in 5 groups (<20; 21-35; 36-50; 51-65; >65 years).

The material for analysis was hair samples 3-5 cm in length taken from at least six spots of the occipital scalp. About 0,5 g of sample was collected from each subject. The samples were thoroughly washed according to the method advised by the IAEA Advisory Group (1985) (acetone, 3X water, acetone), dried at 105<sup>0</sup> C to

constant weight, and mineralised by using wet aching procedure. In the hair were analysed following elements: Al, As, Ca, [Cd, Co, Cr, Cu, Fe, Hg, I, K, Li, Mg, Mn, Na, Ni, P, Pb, Se, Si, Sn, Ti, V and Zn. For analytical determination of all elements except iodine was used atomic emission spectrometry with inductively coupled argon plasma (ICP-AES) (spectrometer Optima 2000 DV, Perkin Elmer, USA), while I was determined by ion-selective electrodes. The accuracy of the method was confirmed by simultaneous analysis of the certified reference material (Chinese human hair GBW09101 powder). Hair element concentrations were expressed as µg/g dry weight.

For noninvasive diagnostic of functional abnormalities in cardiovascular system functioning and biochemical parameters in tissues Device Sensitive Imago was used. electromagnetic oscillations that living body radiate in wide frequency spectrum, that govern practically all biochemical and morphological processes in organism were measured by interface block "AUR-uM", model 3x'30s (Producer – Center of New Technology. Omsk, Russia), equipped with programs that energoinformational data transform to diagnostic and corrective therapy procedures (RUS Patents No.: 2001610661 and 2001110-691, 2001).

As such it assures quick diagnostic of organs functional status and internal interconnections between diseases of organs and systems of organism. In this study we examined functional disturbances in vascular tissues and fallowed coefficients of distribution of other, for vascular status validation, relevant parameters (myocard; aorta; coronary artery; average artery; brain blood vessels; average vene; liver; thymus; thiroidea; chromosomes 1,2,3 and 19; cholesterol in myocard, aorta, brain blood vessels and liver; LDH in average artery; c-reactive protein in myocard, average artery and liver; alkaly phosphatase in average artery and liver; creatinin phospho-kinase in myocard and average artery; potassium in myocard and average artery; Ca and Mg in average artery). Beside mentioned parameters in examined population group were checked to parasites, bacteria and viruses.

## RESULTS AND DISCUSSIONS

Investigations of data obtained for functional disturbances in vascular sistem by bio-resonance diagnostic for different age groups (1/ <20; 2/ 21-35; 3/36-50; 4/ 51-65; 5/ >65) are presented In Tables 1 – 5, while data obtained for biochemical indices of significance for assesment risk for vascular sistem functioning are presented in Tables 6 - 10.

Correlation analysis of data obtained by bioreyonce diagnostic showed that between maz parameters exist correlations (Table 11).

Disturbances from "normal range" obtained in hair elements in different age groups were, in dependence of grade of deficiency or sufficiency, divided in 4 groups (1 – appears only changes in the metabolism of the element and may be transient; 2 – the impairment of the most sensitive biochemical functions and the most labile sources of elements are affected, but this may be compensated by other systems; 3 - appears consequential defects important for health - metabolic, immunological, cognitive, or development processes that do not express overt symptoms; 4 - appears identifiable symptoms that increase with the severity of deficiency - clinical phase) are presented in Table 12.

**Table 1.** Functional disturbances in vascular system functioning for group 1\*, expressed in %

| Parameter        | Coefitients** |       |       |   |   |
|------------------|---------------|-------|-------|---|---|
|                  | 0             | 1     | 2     | 3 | 4 |
| Aorta            | 28,57         | 28,57 | 42,86 | 0 | 0 |
| Coronry arteries | 42,86         | 28,57 | 28,57 | 0 | 0 |
| Brain arteries   | 28,57         | 42,86 | 28,57 | 0 | 0 |
| Averige arteria  | 28,57         | 42,86 | 28,57 | 0 | 0 |
| Averige veins    | 28,57         | 57,14 | 14,29 | 0 | 0 |
| Chromosome 1     | 14,28         | 85,71 | 0     | 0 | 0 |
| Chromosome2      | 14,28         | 85,71 | 0     | 0 | 0 |
| Chromosome 3     | 42,86         | 57,14 | 0     | 0 | 0 |
| Chromosome19     | 42,86         | 57,14 | 0     | 0 | 0 |

\*12-20 ages; n=7

\*\*0 - physiological functional activity; 1 - initial compensatory disturbances of function without symptoms; 2- functional compensatory disturbances with symptoms; 3 - pathological disturbances of homeostasis – compensatory mechanisms disturbed; 4 - compensatory adaptable processes ended – irreversible changes incompatible with live

**Table 2.** Functional disturbances in vascular system functioning for group 2\*, expressed in %

| Parameter        | Coefitient** |       |       |      |   |
|------------------|--------------|-------|-------|------|---|
|                  | 0            | 1     | 2     | 3    | 4 |
| Aorta            | 25           | 50    | 18,75 | 6,25 | 0 |
| Coronry arteries | 31,25        | 46,88 | 18,75 | 3,13 | 0 |
| Brain arteries   | 18,75        | 50    | 28,13 | 3,13 | 0 |
| Averige arteries | 18,75        | 43,75 | 37,5  | 0    | 0 |
| Averige veins    | 12,5         | 62,5  | 18,75 | 6,25 | 0 |
| Chromosome 1     | 18,75        | 81,25 | 0     | 0    | 0 |
| Chromosome2      | 18,75        | 81,25 | 0     | 0    | 0 |
| Chromosome 3     | 31,25        | 65,63 | 3,13  | 0    | 0 |
| Chromosome19     | 36,32        | 63,68 | 0     | 0    | 0 |

\*21-35 ages; n=32

\*\*0 - physiological functional activity; 1 - initial compensatory disturbances of function without symptoms; 2- functional compensatory disturbances with symptoms; 3 - pathological disturbances of homeostasis – compensatory mechanisms disturbed; 4 - compensatory adaptable processes ended – irreversible changes incompatible with live

**Table 3.** Functional disturbances in vascular system functioning for group 3\*, expressed in %

| <b>Parameter.</b> | <b>Coefitient**</b> |          |          |          |          |
|-------------------|---------------------|----------|----------|----------|----------|
|                   | <b>0</b>            | <b>1</b> | <b>2</b> | <b>3</b> | <b>4</b> |
| Aorta             | 22,22               | 44,44    | 22,22    | 11,11    | 0        |
| Coronry arteries  | 22,22               | 27,78    | 38,89    | 11,11    | 0        |
| Brain arteries    | 22,22               | 38,89    | 33,33    | 5,56     | 0        |
| Averige arteries  | 0                   | 50       | 33,33    | 16,67    | 0        |
| Averige veins     | 11,11               | 33,33    | 55,55    | 0        | 0        |
| Chromosome 1      | 22,22               | 77,78    | 0        | 0        | 0        |
| Chromosome2       | 22,22               | 7,78     | 0        | 0        | 0        |
| Chromosome 3      | 5,56                | 94,44    | 0        | 0        | 0        |
| Chromosome19      | 26,46               | 73,54    | 0        | 0        | 0        |

\*36-50 ages; n=18

\*\*0 - physiological functional activity; 1 - initial compensatory disturbances of function without symptoms; 2- functional compensatory disturbances with symptoms; 3 - pathological disturbances of homeostasis – compensatory mechanisms disturbed; 4 - compensatory adaptable processes ended – irreversible changes incompatible with live

**Table 4.** Functional disturbances in vascular system functioning for group 4\*, expressed in %

| <b>Parameter</b> | <b>Coefitient**</b> |          |          |          |          |
|------------------|---------------------|----------|----------|----------|----------|
|                  | <b>0</b>            | <b>1</b> | <b>2</b> | <b>3</b> | <b>4</b> |
| Aorta            | 2,86                | 22,86    | 31,43    | 31,43    | 2,86     |
| Coronry arteries | 8,57                | 40       | 28,57    | 17,14    | 5,71     |
| Brain arteries   | 5,71                | 28,57    | 48,57    | 11,43    | 5,71     |
| Averige arteries | 0                   | 42,86    | 37,4     | 20       | 0        |
| Averige veins    | 5,71                | 25,71    | 45,71    | 17,14    | 5,71     |
| Chromosome 1     | 0                   | 100      | 0        | 0        | 0        |
| Chromosome2      | 5,71                | 94,29    | 0        | 0        | 0        |
| Chromosome 3     | 2,86                | 97,14    | 0        | 0        | 0        |
| Chromosome19     | 14,29               | 85,71    | 0        | 0        | 0        |

\*51-65 ages; n=35

\*\*0 - physiological functional activity; 1 - initial compensatory disturbances of function without symptoms; 2- functional compensatory disturbances with symptoms; 3 - pathological disturbances of homeostasis – compensatory mechanisms disturbed; 4 - compensatory adaptable processes ended – irreversible changes incompatible with live

**Table 5.** Functional disturbances in vascular system functioning for group 5\*, expressed in %

|                  | <b>Coefitient**</b> |          |          |          |          |
|------------------|---------------------|----------|----------|----------|----------|
| <b>Parameter</b> | <b>0</b>            | <b>1</b> | <b>2</b> | <b>3</b> | <b>4</b> |
| Aorta            | 0                   | 9,09     | 36,36    | 17,14    | 0        |
| Coronry arteries | 0                   | 9,09     | 81,82    | 9,09     | 0        |
| Brain arteries   | 0                   | 0        | 45,46    | 45,46    | 9,09     |
| Average arteries | 0                   | 18,18    | 72,73    | 9,09     | 0        |
| Average veins    | 0                   | 18,18    | 45,46    | 27,27    | 9,09     |
| Chromosome 1     | 0                   | 100      | 0        | 0        | 0        |
| Chromosome2      | 0                   | 100      | 0        | 0        | 0        |
| Chromosome 3     | 0                   | 100      | 0        | 0        | 0        |
| Chromosome19     | 9,09                | 90,9     | 0        | 0        | 0        |

\*66-86 ages; n=11

\*\*0 - physiological functional activity; 1 - initial compensatory disturbances of function without symptoms; 2- functional compensatory disturbances with symptoms; 3 - pathological disturbances of homeostasis – compensatory mechanisms disturbed; 4 - compensatory adaptable processes ended – irreversible changes incompatible with live

Data presented for the youngist group of examiners (<20 years) (tables 1 and 6) comparison with data for older group of examiners (20-35 years) (tables 2 and 7) showed that vascular sistem functions in youngist group are more disturbed than in older group. In order to find explanation for this we compared incidence of inflammations with viruses and bacteria in both groups and discovered that 42% of young participants have infection with adenovirus, often together with streptococcus, while in older group 25% has chronic inflammation with adenovirus and 15,6% with candida.

Disturbances in elements present in hair analysis (Table 12) showed that youngist group of examiners suffer from stronger deficiency of many essential elements (I, Co, Se, Mn, Zn, Si, Ca) except Cu that is find in excess. At the some time they have higher concentrations of some of potentially toxic elements than older group (Li, As, Al). Obviously life styel of compared groups differ, as well as their habits. Younger generation was less physicaly active, eat more fast food and various artificiel drinks, has more direct contacts with peoples with various inflammatory processes what obviously reflects negatively to vascular sistem functioning and corresponding biochemical parameters.

**Table 6.** Disturbances in biochemical parameters in group 1\*, expressed in %

|                                 | <b>Coefitient*</b> |                |              |
|---------------------------------|--------------------|----------------|--------------|
| <b>Parameter</b>                | <b>&lt;0,8</b>     | <b>0,8-1,0</b> | <b>&gt;1</b> |
| LDH average arteria             | 0                  | 28,75          | 71,43        |
| Alk. Phos. average arteria      | 57,14              | 28,57          | 14,29        |
| Creat.Phos.Kin. average arteria | 28,57              | 57,14          | 14,29        |
| Cholesterol average arteria     | 14,29              | 85,71          | 0            |
| C-react. prot. average arteria  | 42,86              | 57,14          | 0            |
| Ca average arteria              | 28,57              | 57,14          | 14,29        |
| K average arteria               | 0                  | 85,71          | 14,29        |
| Mg average arteria              | 14,29              | 71,43          | 14,29        |
| Cholesterol aorta               | 28,57              | 57,14          | 14,29        |
| Cholesterol brain arteries      | 0                  | 14,29          | 85,72        |

\*0,4-0,8 normal; 0,8-1 physiological tendency to higher value; &lt;1 pathological changes



**Table 7.** Disturbances in biochemical parameters in group 2\*, expressed in %

| Parameter                       | Coefficient* |         |       |
|---------------------------------|--------------|---------|-------|
|                                 | <0,8         | 0,8-1,0 | >1    |
| LDH average arteria             | 15,63        | 50,00   | 34,38 |
| Alk. Phos. average arteria      | 68,75        | 31,25   | 0     |
| Creat.Phos.Kin. average arteria | 43,75        | 40,63   | 15,63 |
| Cholesterol average arteria     | 59,38        | 34,38   | 6,25  |
| C-react. prot. average arteria  | 65,63        | 31,25   | 3,13  |
| Ca average arteria              | 46,88        | 37,50   | 15,63 |
| K average arteria               | 68,75        | 21,88   | 9,38  |
| Mg average arteria              | 62,50        | 28,13   | 9,38  |
| Cholesterol aorta               | 62,50        | 25,00   | 12,50 |
| Cholesterol brain arteries      | 6,25         | 34,38   | 59,38 |

\*0,4-0,8 normal; 0,8-1 physiological tendency to higher value; <1 pathological changes

**Table 8.** Disturbances in biochemical parameters in group 3\*, expressed in %

| Parameter                       | Coefficient* |         |       |
|---------------------------------|--------------|---------|-------|
|                                 | <0,8         | 0,8-1,0 | >1    |
| LDH average arteria             | 5,55         | 44,44   | 50,00 |
| Alk. Phos. average arteria      | 50,00        | 33,33   | 16,67 |
| Creat.Phos.Kin. average arteria | 33,33        | 38,89   | 27,78 |
| Cholesterol average arteria     | 44,44        | 27,78   | 27,78 |
| C-react. prot. average arteria  | 44,44        | 33,33   | 22,22 |
| Ca average arteria              | 61,11        | 22,22   | 16,67 |
| K average arteria               | 66,67        | 22,22   | 11,11 |
| Mg average arteria              | 50,00        | 27,78   | 22,22 |
| Cholesterol aorta               | 55,56        | 33,33   | 11,11 |
| Cholesterol brain arteries      | 0            | 22,22   | 77,78 |

\*0,4-0,8 normal; 0,8-1 physiological tendency to higher value; <1 pathological changes

Intercomparison of data for groups 3 (36-50 years), 4 (51-65) and 5 (>65 years) showed that with ages I deficiency became more serious, as well as that many other elements have decreasing or increasing trends. Changes in macroelements clearly indicate that acidity of the organism increases with ages, while disturbances in microelement indicate that with aging antioxidative and immunological defence become weaker.

**Table 9.** Disturbances in biochemical parameters in group 4\*, expressed in %

| Parameter                       | Coefficient* |         |       |
|---------------------------------|--------------|---------|-------|
|                                 | <0,8         | 0,8-1,0 | >1    |
| LDH average arteria             | 14,29        | 31,43   | 54,29 |
| Alk. Phos. average arteria      | 71,43        | 22,86   | 5,71  |
| Creat.Phos.Kin. average arteria | 28,57        | 51,43   | 20,00 |
| Cholesterol average arteria     | 48,57        | 37,14   | 14,29 |

**Table 9.** Disturbances in biochemical parameters in group 4\*, expressed in % (cont.)

| Parameter                      | Coefficient** |       |       |
|--------------------------------|---------------|-------|-------|
| C-react. prot. average arteria | 51,43         | 40,00 | 8,57  |
| Ca average arteria             | 48,57         | 42,86 | 8,57  |
| K average arteria              | 45,71         | 37,14 | 17,14 |
| Mg average arteria             | 54,29         | 25,71 | 20,00 |
| Cholesterol aorta              | 51,43         | 34,29 | 14,29 |
| Cholesterol brain arteries     | 8,57          | 14,29 | 77,14 |

\*0,4-0,8 normal; 0,8-1 physiological tendency to higher value; <1 pathological changes

**Table 10.** Disturbances in biochemical parameters in group 5\*, expressed in %

|                                 | Coefficient* |         |       |
|---------------------------------|--------------|---------|-------|
| Parameter                       | <0,8         | 0,8-1,0 | >1    |
| LDH average arteria             | 0            | 54,55   | 45,46 |
| Alk. Phos. average arteria      | 72,73        | 18,18   | 9,09  |
| Creat.Phos.Kin. average arteria | 36,36        | 36,36   | 27,27 |
| Cholesterol average arteria     | 54,55        | 36,36   | 9,09  |
| C-react. prot. average arteria  | 63,64        | 27,27   | 9,09  |
| Ca average arteria              | 72,73        | 27,27   | 0     |
| K average arteria               | 45,45        | 27,27   | 27,27 |
| Mg average arteria              | 63,64        | 9,09    | 27,27 |
| Cholesterol aorta               | 27,27        | 45,46   | 27,27 |
| Cholesterol brain arteries      | 0            | 0       | 100   |

\*0,4-0,8 normal; 0,8-1 physiological tendency to higher value; <1 pathological changes

**Table 11.** Correlations obtained between parameters determined by Bio-resonance

|                                                       |      |
|-------------------------------------------------------|------|
| Average arteria cholesterol/ Average arteria Ca       | 0,79 |
| Average arteria cholesterol/ Average arteria K        | 0,75 |
| Average arteria cholesterol/ Average arteria Mg       | 0,46 |
| Average arteria c-react.prot/ Average arteria K       | 0,63 |
| Average arteria c-react.prot/ Average arteria Ca      | 0,56 |
| Average arteria c-react.prot/c-react prot mocard      | 0,86 |
| Average arteria cholesterol/Average arteria alk. phos | 0,59 |
| Liver c-react.prot/Liver cholesterol                  | 0,50 |
| Ages/aorta function                                   | 0,46 |
| Ages/Average arteria                                  | 0,37 |
| Ages/Miocard function                                 | 0,49 |

Correlation analysis data presented in table 11 showing that disturbances in vascular system functioning highly correlates with Ca, K and Mg and cholesterol in arteries, moderate with c-reactive protein and alkaline phosphatase

**Table 12.** Average disturbances from normal range in hair elements concentrations in investigated groups of residents.

| <b>Ages</b> | <b>Average disturbance grades in hair elements</b>                                                                                             |
|-------------|------------------------------------------------------------------------------------------------------------------------------------------------|
| <20         | I (-1,8); Co (-1,2); Cu(+1,0); Se (-2,6); Zn(-1,4);Mn(-1,0); Cr (-0,6); Si(-1,1); Fe (+0,8); Mg (-0,9); Ca (-1,1); P(+0,6); Na(+0,8); K(-0,9)  |
| 21-35       | I (-2,5); Co (-1,1); Cu(+0,6); Se (-2,1); Zn(-1,0); Mn(-0,6); Cr (-1,1); Si(-1,6); Fe (+0,5); Mg (+1,4); Ca (+2,2); P(+1,1); Na(+0,3); K(-0,6) |
| 36-50       | I (-3,0); Co (-1,0); Cu(+0,3); Se (-1,9); Zn(-1,6); Mn(-0,7); Cr (-0,7);Si(+0,3); Fe (+0,8); Mg (+2,2); Ca (+1,8); P(+0,9); Na(+0,9); K(+0,4)  |
| 51-65       | I (-3,2); Co (-1,1); Cu(-1,2); Se (-2,3); Zn(-1,8);Mn(-1,2); Cr (-1,3); Si(+0,9); Fe (+1,6); Mg (+2,8); Ca (+3,0); P(+2,1); Na(-0,5); K(+0,9)  |
| >66         | I (-2,2); Co (-1,0); Cu(-1,5); Se (-2,6); Zn(-1,2); Mn(-1,6); Cr (-1,0); Si(+0,8); Fe (+2,4); Mg (+3,0); Ca (+3,6); P(+2,6); Na(+0,3); K(+1,3) |

Monitoring of data obtained for inflammations influenced by viruses and/or bacteria in older age groups showed that with aging decrease no of infections with adenovirus, but increases infection iduced by bacteria (in age group 36-50 27,8% with Candida; 51-65 14,3% Candida often with Echerichia; >65 27% Streptococcus and/or Staphylococcus).

## CONCLUSIONS

In the largest number of subjects, early forms of abnormalities were discovered in functioning of abdominal and thoracic aorta (63%); arteries coronaries (66%); brain arterial circulation (69%); venous circulation (75%); arterial blood pressure (50% hypertension, 15% hypotension, 45% variable). These subjects were without symptoms, detectable disorders and dysfunction of tested organs by classical diagnostic methods. Only hair concentrations in elements important for acid[base balance, antioxidative defense and immunological status regulation in their hair were disrupted.

## REFERENCES

1. Airey D. - Mercury in human hair due to environment and diet: a review, Env Health Perspectives, 1983, 52,303-316,.
2. Djujić I, Djermanović V, Jozanov-Stankov O. - Dietary intake of macro and trace elements in Serbia, pp 787-796, In: M. Anke et al. (HRSG), Mengen-und Spurenelemente, 20. Arbeitstagung, Verlag Harald Schubert, Leipzig, 2000.
3. Djujić S.I., Lalić D.T., - Significance of elemental hair analysis for early screening and localization of medical problems, pp. 33-42, In: Metal Elements in Environment, Medicine and Biology, Tome VII, Eds Garban Z. and Dragan P., Romanian Academy Branch Timisoara, Publ."Eurobit", 2006,.

4. Gotovski Y.V., Kosareva L.B., Frolova L.A. - Resonance-frequency diagnostics and therapy of fungi, viruses, bacteria, protozoa and helminthes Methodic recommendations, M.: "IMEDIS", 2000, 70 c;
5. Jeremić S., Brković V., Djujić I. - Toxic elements in scalp hair of Serbian residents (data for 2004), pp.279-292, in: Metal Elements in Environment Medicine and Biology, Eds.: Z.Garban, P.Dragan et al., 6th Int.Symp. of Roumanian Academy-Branch Timisoara, Vol. VI, 2004.
6. Klevay L.M. - The role of copper and zinc in cholesterol metabolism, pp. 227-252, In: Advances in Nutrition Research (Draper HH, ed.), vol. 1, 1st ed., Plenum Publishing Corp., New York, 1977b:.
7. Lalic T., Djujic I. - Elements in hair of population with different grade of hypothyroidism, Cell Biol Toxicol, 2008, 24,S42.
8. Suzuki T., Yamamoto R. - Organic mercury levels in human hair with and without storage for eleven years, Bull Environ Contam Toxicol, 1982, 28, 186-188,.
9. Yacowitz H, Fleischman AI, Bierenbaum ML. - Effects of oral calcium upon serum lipids in man, Br Med J, 1965, 1,1352-1354.
- 10.\*\*\* IAEA Advisory Group - Report on the Second Research Co-ordination Meeting, Neuherberg, FRG, 1985.
- 11.\*\*\* RUS patent No 2001610661, "I-M-A-G-O technology", Center of New Technology OMSK, Russia, 2001.

## FEW WORDS ABOUT METAL COMPOUNDS IN CANCER TREATMENT

Radostina ALEXANDROVA<sup>1</sup>, Konstanta TIMCHEVA<sup>2</sup>, Reni KALFIN<sup>3</sup>, Elena-Maria MOSOARCA<sup>4</sup>, Ramona TUDOSE<sup>4</sup>, Otilia COSTISOR<sup>4</sup>, Daniela C. CULITA<sup>5</sup>, Gabriela MARINESCU<sup>5</sup>, Luminita PATRON<sup>5</sup>, Maria-Lallia KANTOURI<sup>6</sup>

1. Institute of Experimental Pathology and Parasitology, Bulgarian Academy of Sciences, Acad. G. Bonchev, Block 25, BG – 1113 Sofia, Bulgaria; 2. National Specialized Hospital for Active Treatment in Oncology, Sofia, Bulgaria; 3. Institute of Neurobiology, Bulgarian Academy of Sciences, Acad. G. Bonchev, Block 23, BG-1113 Sofia, Bulgaria; 4. Institute of Chemistry Timisoara of the Romanian Academy, 24 Mihai Viteazu Blvd, RO-300223 Timisoara, Roumania; 5. Institute of Physical Chemistry "Ilie Murgulescu", Splaiul Independentei 202, sect.6, RO-060021 Bucuresti, Romania; 6. Aristotle university of Thessaloniki, Chemistry Department, Lab. of Inorganic Chemistry P.O. Box 135, GR/54 124 Thessaloniki, Greece

\* E-mail : rialexandrova@hotmail.com

### ABSTRACT

*Presently, there is large interest in the research of metal complexes as possible anticancer agents. In our investigations we have evaluated the effect of > 50 complexes of Zn(II), Cu(I, II), Co(II), Ni(II) and Fe(II, III) with different ligands (Mannich bases, aminoacids, cholic acids, mixed ligands) on viability and proliferation of cultured human and animal tumour cells. The most promising cytotoxic and antiproliferative agents were found among mixed ligand metal complexes.*

**Key words:** malignant neoplasms, anticancer chemotherapy, metal compounds, cytotoxic and antiproliferative properties, tumour cell lines

### INTRODUCTION

Cancer, a group of more than 100 different diseases, manifests itself as uncontrolled cellular reproduction, local tissue invasion and distant metastases. Cancer therapy employs surgery, radiation, chemotherapy and immunotherapy. Treatment becomes complicated because malignant cells are not pathogens that have specific cure, but are the body's own cells that must be killed or physically removed (Levi et al., 2001). While therapeutical results obtained by surgery and radiotherapy - which are locoregional interventions - are close to their maximum accomplishment, success of drug therapy - the only systemic approach - is far from being satisfactory (Kostova, 2006).

#### Cancer chemotherapy meets problems

Despite decisive progress in understanding the molecular biology of cancer development, cytotoxic anticancer drugs continue to be the cornerstone of modern antitumour therapies (Gonzalez-Nicolini, Fussenegger, 2005). Several obstacles can prevent successful chemotherapy of neoplastic diseases including: 1) Tumour heterogeneity phenomenon (Alexandrova, 2001); 2) Multidrug resistance (Alexandrova, 1998; Sarkadi et al., 2004; Leslie et al., 2005); 3) Undesirable side

effects - the occurrence of myelosuppression, mucositis, cardiodepression, peripheral neuropathy, and central neurotoxicity can complicate treatment especially in older patients (65 years of age and older) (Repetto, 2003; Ocean, Vohdat, 2004). Consequently, the need to find a safe and highly effective cure for neoplastic diseases remains a major challenge for modern science (Levi et al., 2001; Gonzales-Nicolini, Fussenegger, 2005; Kostova, 2006).

### Metal compounds as antitumour agents

The earliest reports on the therapeutic use of metals or metal-containing compounds in cancer and leukemia date from the sixteenth and nineteenth centuries (Kopf-Maier, 1994; Desoize, 2004). They were forgotten until the 1960s, when the anti-tumour activity of the inorganic complex cis-diammine-dichloroplatinum (II) (cisplatin) was discovered.

**Table 1.** Metal complexes with Mannich bases

| Code  | Formula                                             | Molecular weight (g/mol) |
|-------|-----------------------------------------------------|--------------------------|
| TS-1  | $\text{Cu}_2(\text{BAMP})(\text{NCS})_4$            | 846.00                   |
| TS-2  | $\text{Cu}_2(\text{BAMP})\text{I}_3$                | 1007.00                  |
| TS-3  | $\text{Co}_2(\text{BAMP})\text{Cl}_4$               | 746.00                   |
| TS-4  | $\text{Co}(\text{BAMP})(\text{NCS})_2$              | 661.00                   |
| TS-5  | $\text{Fe}_2(\text{BAMP})\text{Cl}_4$               | 740.00                   |
| TS-6  | $\text{Fe}_2(\text{BAMP})\text{Cl}_6$               | 811.00                   |
| TS-7  | $\text{Cu}(\text{TAMEN})(\text{NO}_3)_2$            | 1046.00                  |
| TS-8  | $\text{Co}_2(\text{TAMEN})\text{Cl}_4$              | 1121.00                  |
| TS-9  | $\text{Co}_2(\text{TAMEN})(\text{NCS})_4$           | 1211.00                  |
| TS-10 | $\text{Fe}_2(\text{TAMEN})\text{Cl}_6$              | 1186.00                  |
| TS-11 | $\text{Ni}(\text{TAMEN})(\text{ClO}_4)_2$           | 1119.69                  |
| TS-12 | $\text{Ni}(\text{TAMEN})(\text{NCS})_2$             | 1036.69                  |
| TS-13 | $\text{Ni}_2(\text{BAMP})(\text{CH}_3\text{COO})_4$ | 839.38                   |
| TS-14 | $\text{Ni}_2(\text{BAMP})(\text{Cl})_4$             | 745.38                   |
| TS-15 | $\text{Fe}(\text{TAMEN})(\text{NO}_3)_3$            | 1103.85                  |
| TS-16 | $\text{Fe}(\text{BAMP})(\text{NO}_3)_3$             | 727.85                   |
| TS-17 | $\text{Cu}(\text{TAMEN})(\text{ClO}_4)_2$           | 1124.55                  |
| TS-18 | $\text{Cu}(\text{TAMEN})(\text{CH}_3\text{COO})_2$  | 1043.55                  |
| TS-19 | $\text{Cu}_2(\text{BAMP})(\text{ClO}_4)_4$          | 1011.10                  |
| TS-20 | $\text{Co}(\text{TAMEN})(\text{ClO}_4)_2$           | 119.93                   |
| TS-21 | $\text{Fe}_2\text{BAMPpy}_2\text{Cl}_4$             | 898.33                   |
| TS-22 | $\text{Fe}_2\text{BAMPpy}_2\text{Cl}_6$             | 969.23                   |
| TS-23 | $\text{Co}_2\text{BAMPpy}_2\text{Cl}_4$             | 904.50                   |
| TS-26 | $\text{Cu}_2\text{BAMPdipyCl}_4$                    | 911.71                   |
| TS-27 | $\text{Fe}_2\text{TAMENDipyCl}_4$                   | 1270.75                  |
| TS-28 | $\text{Fe}_2\text{TAMENDipyCl}_6$                   | 1341.66                  |

### The antitumour power of platinum compounds

The square planar platinum (II) complex, cis-[PtCl<sub>2</sub>(NH<sub>3</sub>)<sub>2</sub>] (cis-DDP, cisplatin), was first synthesized in 1844 by Michael Peyrone in Turin and is historically known as Peyron's chloride. Fifty years later in his theory of coordination chemistry Alfred Werner correctly assigned to this compound a cis-geometry, which was confirmed in an X-ray crystal-structure determination reported in 1966. The biological activity of cis-platin was accidentally discovered in 1965 by Barnet Rosenberg and his

coworkers who reported first the inhibitory activity of this compound on *E. coli* division. In the seventies efficacy in human cancer patients was established. A little bit later these scientists performed experiments with sarcoma 180 and leukemia L1210 bearing mice and the results obtained led to cisplatin entering phase I clinical trials in 1971. Approval to cisplatin for treatment of testicular and ovarian cancer was given in 1978 (Rosenberg et al., 1965; Lippert, 1999; Desoize, 2004; Abu-Surrah and Kettunen, 2006).

Evidence for antitumour activity in humans was first reported against testicular and ovarian cancer, in terms of both objective response and of prolonged survival. Because of its marked renal toxicity, cisplatin was almost withdrawn from clinical trials. Interest was rekindled when hyperhydration with isotonic saline circumvented this problem. One of the attractions of cisplatin, besides its efficacy, is that its toxicity is different from that of other anticancer drugs, thus making it attractive for combination regimens. Today cis-platin is the world's leading anti-tumour drug for the chemotherapy of human cancer. This platinum compound continues to be used in 50-70% of all cancer patients especially against testicular and ovarian cancer, and it is also used in case of head and neck cancer, cervical and bladder carcinomas, bronchogenic carcinoma, small-cell and non-small cell lung cancers, lymphoma, osteosarcoma, melanoma and neuroblastoma (Wong and Giandomenico, 1999; Galanski et al., 2003; Abu-Surrah and Kettunen, 2006).




**Fig. 1.** N,N'-bis(4-antipyrilmethyl)-piperazine (BAMP)

Although cis-platin is highly effective anticancer agent, it is not ideal because of:

- 1) Low or no activity against some of the most common and socially important malignant neoplasms such as breast, gastrointestinal and renal cancers, leukemia;
- 2) The development of drug resistance during treatment is not rare;
- 3) It has major toxicity limitations of which nephrotoxicity is the most notable, although nausea and vomiting, peripheral neuropathy, and myelotoxicity can also raise major concerns (Galanski et al., 2003; Abu-Surrah and Kettunen, 2006).

The landmark discovery of the antitumour activity of cisplatin was an extremely fortunate one for a number of reasons: 1) Cis-platin gave and still gives patients suffering from various types of cancer hope for a prolonged and a better quality of life, and the chance of a cure; 2) It had an impact on inorganic and coordination chemistry that cannot be overestimated; 3) It is a wonderful example of that basic

research is capable of achieving and a pledge for the support of basic science (Lippert, 1999).




**Fig. 2.** N,N'-tetra-(antipyryl-1-methyl)-1,2-diaminoethane (TAMEN)

Since the introduction of cisplatin around three thousand platinum derivatives have been synthesized and tested against cancer cells; but, at most, only thirty compounds have reached clinical trials and more than half of those have already been rejected. The main aim of these intensive investigations was to obtain drugs with at least an equal activity but reduced toxicity compared to cisplatin. Today, six are used clinically: three of them – cisplatin, carboplatin and oxaliplatin, are used world-wide, whereas the other three are applied in a few countries - nedaplatin in Japan, lobaplatin in China, and heptaplatin – in south Korea. Some platinum complexes are still under clinical investigation, including those developed for oral administration (Desoize and Madoulet, 2002; Abu-Surrah and Kettunen, 2006; Choy, 2008).

### **The antitumour properties of non-platinum compounds**

The unresolved problems (narrow spectrum of cancers that can be treated; side effects; resistance phenomena) in platinum-based anticancer therapy have stimulated increased research efforts in the development of novel non platinum-containing metal species as cytostatic agents. As a result, a wide variety of ruthenium (Alessio et al., 2004; Kostova, 2006b), gallium (Collery, 2002; Jacupec, Keppler, 2004), titanium (Desoize, 2004), gold (Messori, Marcon, 2004; Kostova, 2006a), vanadium (Chatterjee, Bishayee, 1998; Evangelou, 2002), lanthanides (Kostova, 2005), zinc (Alexandrova et al., 2002), copper (Alexandrova et al., 2003), cobalt (Alexandrova et al., 2004), nickel (Alexandrova et al., 2006), iron (Alexandrova et al., 2005) compounds have been reported to express promising antitumour properties in preclinical studies. Some of them (titanium, ruthenium, gallium



compounds) have already been evaluated in Phase I and Phase II trials (Desoize, 2004; Abu-Surrah, Kettunen, 2006; Ott, Gust, 2007).

In the search for new agents with anticancer efficacy we have investigated the antitumour activity of more than 50 metal compounds such as:

1. Zn(II), Cu(II) and Co(II) complexes with L picolinic or DL aspartic acid) [28];
2. Zn(II), Cu(II), Co(II), La(III) complexes with cholic, dehydrocholic and lithocholic acids;
3. Cu(I, II), Co(II), Ni(II) and Fe(II, III) complexes (TS1-TS20, Table 1) with Mannich-bases N,N'-bis(4-antipyrilmethyl)-piperazine (BAMP, Fig. 1) and N,N'-tetra-(antipyril-1-methyl)-1,2-diaminoethane (TAMEN, Fig.2)
4. Cu(I, II), Co(II) and Fe(II, III) mixed ligand complexes containing BAMP or TAMEN as well as pyrimine (py) or 2,2-dipyridyl (dipy) as a coligand (TS, Table 1).
5. Co(II) complexes with mixed ligands 2-hydroxy-benzophenones and nitrogenous bases enR (still under investigation).

The following model systems were used in our experiments: 1) Permanent cell lines from human (breast cancer MCF-7, glioblastoma multiforme 8 MG BA, carcinoma of the larynx Hep-2, hepatoma HepG2, rhabdomyosarcoma RD64, squamous cell carcinoma A431 and erythroleukemia K562), rat (transplantable sarcoma induced by *Rous sarcoma virus* strain *Schmidt-Ruppin*), murine (myeloma P3U1) and chicken (transplantable hepatoma induced by the myelocytomatosis virus Mc29) origin; 2) Primary cultures from 11 day old 151 White Leghorn chickens.

The influence of the compounds on cell viability and proliferation as well as cytopathological changes were studied by MTT test, neutral red uptake cytotoxicity assay, trypan blue dye exclusion technique, colony-forming method, autoradiography, neutral and alkaline variants of single cell gel electrophoresis and acridine orange staining.

The results that we obtained revealed that:

1. According to their ability to reduce cell viability and proliferation, the examined metal complexes with aminoacids were graded as follows:  
Zn(II) complexes > Cu(II) complexes > Co(II) complexes
2. Co(II) and Ni(II) complexes with BAMP expressed more pronounced cytotoxic and antiproliferative properties as compared to TAMEN-containing complexes of the same metals. At the same time Fe(II, III) complexes with TAMEN are found to be active than iron complexes with BAMP.
3. Applied independently, the ligands (BAMP, TAMEN, lithocholic and dehydrocholic acids) were found to be less effective cytotoxic and antiproliferative agents as compared to the corresponding metal complexes;
4. The compounds with the most promising antineoplastic properties in cultured tumour cells were found among mixed ligand metal complexes. Thus, the  $CC_{50}$  of  $Cu_2(BAMP)dipyCl_4$  determined on 24 h were found to vary between 3  $\mu g/ml$  (LSCC-SF-Mc29) and 34  $\mu g/ml$  (8 MG BA) and on 48 h between 1  $\mu g/ml$  (K562) and 23  $\mu g/ml$  (8-MG-BA). Applied at concentration of 1 to 10  $\mu g/ml$  this compound completely inhibited the growth of tumor cells in semisolid medium and reduced significantly the incorporation of  $^3H$ -thymidine. Shrinkage of the cell and its nucleus, plasma membrane blebbing, chromatin condensation and segmentation of the nuclei as well as formation of apoptotic bodies were observed after acridine orange staining in the cells treated with  $\geq 5 \mu g/ml$   $Cu_2(BAMP)dipyCl_4$ . DNA damages were found by

single cell gel electrophoresis in 60% of the tumor cells cultured for 48 h in the presence of this compound (10 µg/ml). The mixed ligand copper (II) complex expressed more pronounced cytotoxic and cytostatic activities against LSCC-SF-Mc29 hepatoma and P3U1 mouse myeloma cells than on primary chicken embryo cells and BALB/c 3T3 mouse fibroblasts. Some of the Co(II) complexes with mixed ligands 2-hydroxy-benzophenones and nitrogenous bases enR were found to reduce cell viability by 50% (IC<sub>50</sub>) when applied at concentrations < 5 µg/ml for 24 h. These compounds were also proved to induce DNA damages (double stranded breaks) in the treated (5, 20 and 50 µg/ml for 24 h) cells.

The need to find a safe and highly effective cure for neoplastic diseases remains a major challenge for modern science. The ideal anticancer treatment by any compound should fulfill certain criteria and exert the following selective effects on malignant cells: 1) reduction of cellular growth rate (antiproliferative effect); 2) cytotoxic action expressed either by necrosis or apoptosis; 3) reduction and/or inhibition of the invasive or metastatic potential of the cells; 4) reduced or absent potency to induce cellular resistance; 5) low side effects (Messori and Marcon, 2004). Developing metal compounds as drugs is not an easy task, because accumulation of metal ions in the body can lead to deleterious effects. Future investigations will clarify the potential antitumour properties and mechanism of action of cobalt and cobalt compounds and will facilitate the design of drugs with improved antineoplastic activity and reduced general toxicity.

**Acknowledgements.** This study was supported by Grant C1402/2004 from the National Science Fund in Bulgaria; Bilateral project between Bulgarian Academy of Sciences and Aristotle University of Thessaloniki, Greece; Bilateral project between Bulgarian Academy of Sciences and Romanian Academy.

## REFERENCES

1. Abu-Surrah, A.S., Kettunen M. : Platinum group antitumor chemistry: Design and development of new anticancer drugs complementary to cisplatin. – *Curr. Med. Chem.*, 13, 2006, 1337-1357.
2. Alessio, E., Mestroni G., Bergamo A., Sava G. : Ruthenium antimetastatic agents. - *Curr. Topics Med. Chem.*, 4, 2004, 1525-1535.
3. Alexandrova R., Rashkova G., Alexandrov I., Tsenova W., Tudose R., Costisor O. : Briefly about copper. *Exp. Pathol. Parasitol.* 6-12 (2003) 3-12.
4. Alexandrova R., Tudose R., Arnaudova E., Costisor O., Patron L.: Cobalt. *Exp. Pathol. Parasitol.* 7-2 (2004) 3-14.
5. Alexandrova, R. : Multidrug resistance and P-glycoprotein. *Minireview. Experimental Pathology and Parasitology*, 1, 1998, 62-66.
6. Alexandrova, R. : Tumour heterogeneity. *Exp. Pathol. Parasitol.*, 2001, 4/6, 57-67.
7. Alexandrova, R., Rashkova G., Salkova D., Sainova I. : Something more about zinc. *Exp. Pathol. Parasitol.*, 2002, 5/8, 17-24.
8. Alexandrova, R., Rashkova G., Patron L., Costisor O. : Iron. *Exp. Pathol. Parasitol.*, 8/3, 2005, 49-60.
9. Alexandrova, R., Costisor O., Patron L. : Nickel. *Exp. Pathol. Parasitol.*, 2006, 9/1, 64-74.
10. Chatterjee, M., Bishayee A. : Vanadium – a new tool for cancer prevention. In . *Vanadium in the environment*. J.O. Nriagu, ed. John Wiley&Sons. NY, 1998.

11. Choy H, Park C., Yao M.: Current status and future prospects for satraplatin, an oral platinum analogue. *Clin Cancer Res.*, 14, 2008, 1633-8.
12. Collery P., Keppler B., Madoulet C., Desoize B. : Gallium in cancer treatment. *Crit. Rev. Oncol. Hematol.*, 42, 2002, 283-296.
13. Desoize B : Metals and metal compounds in cancer treatment. - *Anticancer Res.*, 24, 2004, 3-18.
14. Desoize B., Madoulet C. : Particular aspects of platinum compounds used at present in cancer treatment. - *Crit. Rev. Oncol. Hematol.*, 42, 2002, 317-325.
15. Desoize B., Madoulet C. : Particular aspects of platinum compounds used at present in cancer treatment. - *Crit. Rev. Oncol. Hematol.*, 42, 2002, 317-325.
16. Evangelou A.M. : Vanadium in cancer treatment. *Crit. Rev. in Oncol. Hematol.*, 42, 2002, 249-265.
17. Galanski M., Arion V.B., Jakupec M.A., Keppler B.K. : Recent developments in the field of tumor-inhibiting metal complexes. - *Curr. Pharm. Des.*, 9, 2003, 2078-2089.
18. Gonzalez-Nicolini V., Fussenegger M. : In vitro assays for anticancer drug discovery - a novel approach based on engineered mammalian cell lines. - *Anticancer Drugs*, 16, 2005, 223-228.
19. Jakupec, M.A., B.K. Keppler. Gallium in cancer treatment. *Curr. Top. Med. Chem.*, 4, 2004, 1575-1583.
20. Kopf-Maier, P. Complexes of metals other than platinum as antitumor agents. - *Eur. J. Clin. Pharmacol.*, 47, 1994, 1-16.
21. Kostova, I. Gold coordination complexes as anticancer agents. - *Anti-Cancer Agents - Med. Chem.*, 6, 2006, 1-14.
22. Kostova, I. Gold coordination complexes as anticancer agents. - *Anti-Cancer Agents - Med. Chem.*, 6, 2006a, 1-14.
23. Kostova, I. Lanthanides as anticancer agents. - *Curr. Med. Chem. - Anti-Cancer Agents*, 5, 2005, 1-12.
24. Kostova, I. Lanthanides as anticancer agents. - *Curr. Med. Chem. - Anti-Cancer Agents*, 5, 2005, 1-12.
25. Kostova, I. Ruthenium complexes as anticancer agents. - *Curr. Med. Chem.*, 13, 2006b, 1085-1107.
26. Leslie, E.M., R.G. Deeley, S.P.C. Cole. Multidrug resistance proteins: role of P-glycoprotein, MRP1, MRP2, and BCRP (ABCG2) in tissue defense. *Toxicol. Appl. Pharmacol.*, 204, 2005, 216-237.
27. Levi, M.S., R.F. Borne, J.S. Williamson. A review of cancer chemopreventive agents. - *Curr. Med. Chem.*, 8, 2001, 1349-1362.
28. Lippert, B. Cisplatin. Chemistry and biochemistry of a leading anticancer drug. Lippert, B. (ed). Verlag Helvetica Chimica Acta, Zurich, 1999.
29. Messori, L., G. Marcon. Gold complexes as antitumor agents. - *Met Ions Biol. Syst.*, 42, 2004, 385-424.
30. Ocean, A.J., L.T. Vahdat. Chemotherapy-induced peripheral neuropathy: pathogenesis and emerging therapies. - *Support Care Cancer*, 12, 2004, 619-625.
31. Ott, I., R. Gust. Non platinum metal complexes as anti-cancer drugs. - *Arch. Pharm. Chem. Life Sci.*, 340, 2007, 117-126.
32. Peyrone, M. Über die einwirkung des ammoniaks auf platinchlorid. - *Annalen der Chemie und Pharmacie*, L1, 1844, 1-29.
33. Repetto, L. Greater risks of chemotherapy toxicity in elderly patients with cancer. - *J. Support. Oncol.*, 4 Suppl. 2, 2003, 18-24.

34. Rosenberg, B., L. Van Camp, T. Krigas. Inhibition of cell division in *Escherichia coli* by electrolysis products from platinum electrode. - *Nature*, 205, 1965, 698-699.
35. Sarkadi, B., C. Ozvegy-Laczka, K. Nemet, A. Varadi. ABCG2 - a transporter for all seasons. *FEBS Lett.*, 567, 2004, 116-120
36. Wong, E., C.M. Giandomenico. Current status of platinum-based antitumor drugs. *Chem. Rev.*, 99, 1999, 2451-2466.

# A STUDY ON THE POSSIBLE PREVENTION OF ENVIRONMENTAL POLLUTION WITH MINING WASTES FROM COPPER MINES THROUGH AGRICULTURAL VALORISATION

Ioan BORCEAN\*, Gheorghe DAVID, Simona NIȚĂ, Adrian BORCEAN, Narcis BĂGHINĂ, Lucian NIȚĂ

*Department of Phytotechnics, Faculty of Agriculture, University of Agricultural Sciences and Veterinary Medicine of Banat, Calea Aradului nr. 119, RO-300 475 Timișoara, Roumania*

\* E-mail: borcean\_ioan@yahoo.com

## ABSTRACT

*We tested the impact of mining wastes from copper mines in southern Banat on wheat, maize, oat, and bean crops. Mining waste analysis results point out that this material can be used for agricultural purposes since heavy metal content is within accepted limits. We applied waste amounts of 1t/ha, 2t/ha, and 3t/ha combined with variable nitrogen rates on a constant phosphorus and potassium agro-fund ( $P_{60}K_{60}$ ).*

**Key words:** agricultural use, mining wastes, copper mine

## INTRODUCTION

Our starting point in this study was to see if the thousands of tonnes of mining wastes from the copper mines in the Moldova Nouă and Sasca Montană area can be valorised as a source of oligo-elements and macro-elements in agriculture.

This hypothesis could be tested since the material is not radio-active and heavy metal content is below accepted limits at European level.

The material was found to have an alkaline reaction as a result of high contents of calcium, and because flotation was done with basic elements; this indicates that it is possible to use the material to correct soil acid reaction in the hill area of western Romania.

## MATERIALS AND METHODS

Table 1 presents analysis results concerning waste sample chemical element content.

The experiments were bi-factorial, with Factor A represented by mining waste (we tested amounts of 1t/ha, 2t/ha, and 3t/ha) applied on a constant potassium and phosphorus agro-fund ( $P_{60}K_{60}$ ) and with Factor B represented by variable rates of nitrogen, i.e.  $N_0$ - $N_{150}$  in wheat, maize, and oat and  $N_0$ - $N_{90}$  in bean.

The type of soil on which our research was carried out is a typical brown eumeso-basic soil with a pH 5.6, a humus content of 2.61%, and a clayish-sandy texture.

**Table 1.** The element content of mining sterile from Moldova Nouă

| Element | The content in the mining steril store |        |
|---------|----------------------------------------|--------|
|         | Test 1                                 | Test 2 |
| Si      | 41                                     | 53     |
| Al      | 2                                      | 11     |
| Fe      | 1,5                                    | 5      |
| Ti      | 0,2                                    | 0,25   |
| P       | 0,08                                   | 0.06   |
| Ca      | 25                                     | 32     |
| Mg      | 2,4                                    | 1,7    |
| Mn      | 0,11                                   | 0.09   |
| S       | 0,30                                   | 0,47   |
| As      | 0,001                                  | 0,003  |
| Bi      | 0,01                                   | 0,005  |
| Mo      | 0,004                                  | 0,004  |
| Cu      | 0,1                                    | 0,08   |
| Zn      | 0.07                                   | 0,02   |
| Cr      | 0,002                                  | 0,004  |
| Ni      | 0,001                                  | 0,002  |
| Water   | 8,0                                    | 9,0    |

There were traces of B, Pb, Cd, Se, and Ag.

## RESULTS AND DISCUSSIONS

Results in winter wheat are presented in Figure 1, which show that the smallest crops and, therefore, the lowest valorising of fertilisers were on an agro-fund with no mining wastes. On the agro-funds treated with 1.2 and 3 t/ha of mining wastes, fertiliser valorisation was higher, as a result of the positive impact of the mining wastes neutralising soil acid reaction, and of the fact that micro- and macro-elements improved plant nutrition.

For a rate of  $N_{150}$ , yields obtained on the agro-funds treated with 2 t/ha and 3 t/ha were pretty close, which means that increasing waste is not motivated. At the same time, this allows avoiding the danger of increasing heavy metal content in the soil.

Figure 2 shows that the lowest yields in maize were on the agro-fund not treated with mining wastes while the highest yields were obtained on the agro-fund treated with 3 t/ha of mining wastes and fertilised with  $N_{150}$ - $N_{200}$ .

Results obtained in oat Figure 4 point out that the highest yields were on the agro-fund treated with 2 t/ha and 3 t/ha of mining wastes and fertilised with  $N_{150}$ .

Results obtained in bean Figure 4 show a positive response within the range  $N_0$ -  $N_{100}$ , on all mining waste funds, with the best results on the fund treated with 3 t/ha.




**Fig. 1.** Response curves wheat, crops to nitrogen fertilisation with variable levels of mining wastes on a constant agro-fund fertilised with P<sub>60</sub>K<sub>60</sub>




**Fig. 2.** Response curves maize, crops to nitrogen fertilisation with variable levels of mining wastes on a constant agro-fund fertilised with P<sub>60</sub>K<sub>60</sub>




**Fig. 3.** Response curves oat, crops to nitrogen fertilisation with variable levels of mining wastes on a constant agro-fund fertilised with P<sub>60</sub>K<sub>60</sub>




**Fig. 4.** Response curves of bean, crops to nitrogen fertilisation with variable levels of mining wastes on a constant agro-fund fertilised with P<sub>60</sub>K<sub>60</sub>

## CONCLUSIONS

Mining wastes from the copper mines in the Moldova Nouă and Sasca Montană area are not radio-active and heavy metal content is below European accepted limits.

Alkaline reaction of the mining wastes and their content in micro- and macro-elements make it possible to use them on acid reaction soils.

Applying mining wastes in rates of 2-3 t/ha lead to a better valorisation of the nitrogen fertilizers applied on an agro-fund of P<sub>60</sub>K<sub>60</sub>.

## REFERENCES

1. Baker A.J., Walker P.L. :Ecophysiology of mental uptake by tolerant plants, In Heavy Mental Tolerance in Plants.Evolutionary Aspects Shaw A.J.,Ed. R.C.Press,Boca Raton, Fl., p.155 - 177
2. Bogdanovici Darinka , Ubavic M.,HadzicV. :Heavy Metals in soil, Heavy Metals in the Enviroment.Novi Sad,1997, p.97-143
3. Borcean I.,Borcean A., Goian M., Şuveţi Simona : A possible valorification of mine steril in copper mines, pp. 99-102, in Metal Elements in Environment Medicine and Biology, Tome III (Eds. Garban Z., Dragan P.),Publishing House Eurobit, Timisoara, 1998
4. Borcan I.,Borcean A., Goian M.: Mine steril of banatit from the souther part of Banat country, a valuable amendament and source of lements, Proceedings of the Symposium Restoration Ecology, Timisoara, Ed.Orizonturi Universitare, 2001, p .98.
5. Chisaliță I. : Cercetări privind stabilizarea haldelor de steril de la Moldova Noua cu ajutorul vegetației forestiere si influența acesteia asupra mediului, Braşov, Teza de doctorat, Universitatea Transilvania, 2001.



# STUDY OF THE COMPLEXES FORMED BETWEEN SOME METAL IONS WITH THE CONDENSED PHOSPHATES BY MEANS OF GEL- CHROMATOGRAPHY

Aurel CONTREA

*Department of Biochemistry, Faculty of Animal Sciences and Biotechnologies, Univesity of Agricultural Sciences and Veterinary Medicine of Banat, Calea Aradului, Nr. 119, RO-300 645 Timisoara, Roumania*

## ABSTRACT

*The Hummel and Dreyer effect were used to the study of complex formation between  $\text{Cu}^{2+}$ ,  $\text{Zn}^{2+}$  and  $\text{Fe}^{3+}$  ions and diphosphate, triphosphate and polyphosphate by steric-exclusion gel-chromatography on Sephadex columns.*

*The stability constants of the metal complexe were determined and the effect of pH on the complexation equilibria were investigated.*

*The method is useful for the physico-chemical analysis of trace minerals in the presence of condensed phosphates.*

**Key words:** complex metal-polyphosphates, gel chromatography, stability constants.

## INTRODUCTION

The liniary polyphosphates forms soluble complexes with the divalent and trivalent ions (1,5). The usually study methods for the forming of complexes as the spectrophotometry, nephelometry, potentiometry, conductometry a.o. , being static methods, permitted not the simultaneous research of the of the resulted complexes formed by the metal ions with the many mixed ligands.

Hummel and Dreyer (1962) were applied for the first time the sterical exclusion chromatography to the reversible interaction between the molecules and the substances with little molecular weight molecules, method which enable the direct observation of the complex forming which are in equilibrium with the free metal ion, maintained to a predetermined constant value.

Yoza and his coworkers were developped this experiment based on a dynamic method, which were applied for the stability constants determination of the magnesium polyphosphates (Kouchiyama et all., 1978).

In the present study, the complex formation of the copper, zinc and iron, transition metals having trace minerals role in the biological systhems with the diphosphate, triphosphate and the high polycondensed polyphosphate (Kurrol salt) were investigated.

## MATERIALS AND METHODS

We used diphosphate, triphosphate (obtained by means of repeated recrystalisations of the technical tripolyphosphate high polymerised), potassium polyphosphate (Kurrol salt) and Blue Dextran 2000.

The chromatography were made on Sephadex G – 15, Sephadex G-25 and Seohadex G-50 fine gells columns with 900x15 mm, placed in equilibrium with buffer solutions used for the samples elution, in which the metal ion concentration were  $5 \cdot 10^{-5}$  M. The eluent flow speed were insured with a peristaltic pump. The sample solution of 1 ml. formed of the polyphosphatic ligands in presence of an metal excess were introduced in the top of the column.

The eluent fractions of 2 ml were collected with an fraction collector. The eluated  $\text{Fe}^{3+}$  fractions were determined photocolometrically by means of the ammonium sulphocyanure in presence of aluminium nitrate, for the iron delivering from the condensed phosphate (H.E.Wirth, 1942).

The phosphorus were determined with the  $\text{Mo}^{\text{V}} - \text{Mo}^{\text{VI}}$  reagent (Yoza et all., 1977), using a colorimetrical automatic line.

The copper and zinc were determined by means of an atomic absorption spectrograph. The used eluents were:

0.08 M NH OH – 0,02 M  $\text{NH}_4\text{Cl}$  (pH = 10)

0.01 M HNO – 0,1  $\text{NaNO}_3$ ) (pH = 2)

A and B contained  $5 \cdot 10^{-5}$  M metall ions.

## RESULTS AND DISCUSSIONS

The study of the ferric ions complexes with the condensed phosphates in the fine Sephadex G – 25, pH = 2 make evident the forming of the soluble complexes with the all ligands used. The ferric ions endure hydrolyse, forming insoluble hydrated polymers at pH values upper to 2.5 (Spiro et all., 1966).

When we graphical represent the metal concentration in the eluent as function of the total eluent, we obtain the elution profile, known as the Hummel Dreyer effect,

The elution prophile of a so experiment presents an positive peak for the metal complex ML, which walk speedy together with the ligand, as compared with the free metal ion M, which go slower.

The elution volumes of the free ferric ion and of the iron – polyphosphates complexes decreased in speed in reverse order with their molecular weights:  $\text{Fe} > \text{FeP} > \text{FeP} > \dots > \text{FeP}$ .

Yoza, 1975, showed that through the introduction in the sample of a ligands mixture with various molecular weights together with a metal excess, which insure to forming of the all complexes and to remain also an excess, so appear an positive peak at the metall ion elutions position, fact confirmed also diphosphate and triphosphate elution.

In this manner we may study through a single experiment many complexes of an metal ion with various polyphosphatic ligands in mixture., as well as the competitive phenomena which may interweave between many metal ions for the binding site of the same ligand.

In the case when the metal ions are not in excess in the sample, after the peak of the Fe-triphosphate complex, the iron concentration of the elution continues to be constant, and afterwards it decreased at the elution position of the free iron ion, forming an negative peak, which corresponds to the iron amount consumed from the elution solution.

The study of the concentration at the elution of the  $\text{Fe}^{3+}$  complex of the high polymerised Kurrol salt is useful as a marker for the determination of the zero volume

of the fine Sephadex G-25 column, this being excluded from the gel phase at the position of the corresponding position of the Blue Dextran.

The possibility to metal-complexes forming of the polyphosphates with copper ions were studied with the fine Sephadex gel G-25 at pH = 10. The elution profile make evident also in this case the soluble complexes forming with the studied polyphosphates.

The elution position of the  $\text{Cu}^{2+}$  -complex with the Kurrol salt were controlled also by means of the phosphorus determination in the eluted fractions, obtaining a perfect concordance. The fine Sephadex G-25 enables the copper complexes separation, formed in the presence of a ligands mixture.

For to render evident the resolution power of the metal complexes with the condensed phosphates also on another Sephadex type, we worked at the zinc complexes with a fine gel Sephadex G-15 and at pH = 10. The quantitative valuation of the gel chromatographic behaviour of the complexes needs to know some parameters of the columns as well as the void volume  $V$  and the elution position of the free (not complexed) Zn ion. The elution profile of the zinc complex with the Kurrol salt make obvious the fact that the his retain volume are independent of the complexed iron and of the eluent pH-value.

We have studied the Zn- complexes with the the diphoaphate and the triphosphate, in presence of a zinc ions excess in the eluted sample.

When the chromatography were made on a gel with the large inside pores, as well as the fine Sephadex G-50, these enables to separate the Zn-complexes with the studied polyposphatic ligands.

One can observe that in this case an increase the separation space amongst the retention volume of the  $\text{Zn-P}_3$  complexes and the exclusion of the  $\text{ZnP}_n$  complex, fact which enables to separate from a mixture also the metal complexes with ligands having a greather condensation degree as three (tetraphosphate, pentaphosphate, hexaphosphate, etc.)

The so kind of oligomeric polyphosphates are encountered in some technical matters, as well as the polyphosphate glasses with trace minerals, as well as the liquid complex fertilizers with trace minerals, based on the ammonium polyphosphoric acids.

The analytical methods presented in this paper may be apply also to the technical analysis of the this industrial products.

The study by means of the sterical gel-exclusion chromatography enable together to characterize the formed complexes, respective the metal-ligands ratio, as well as to determine the stability constants.

The metal-ligand complex formation may be represented by the relation:  $\text{M} + \text{L} = \text{ML}$  (1)

The stability constant of the metal-ligand complex are definable by the known relation:  $K = [\text{ML}] / [\text{M}][\text{L}]$  (2)

After Yoza,(1977), in the study of the metal-ligand complexes by means of the sterical - exclusion in the gel chromatography, when the complex migration accomplish in the presence of a predicted concentrations an constante value of the stability contant in the eluant, nominated as  $[\text{M}]_0$  the stability constant value result by means of the relation:  $K = [\text{ML}] / [\text{M}_0](\text{[L]}_t - [\text{ML}])$  (3)

In which  $L_t$  = the total concentration of the introduced ligand in the sample.

Because during the complex migration, it is permanently in contact with the free metal ions  $[M_0]$ , the complex stability are insured, this being into a permanent dynamic equilibrium.

When the complex ML elution would be effectuate with an eluant which not contain free M and L, this successive dissociate into his constituents during the migration, to get to a complicated elution profile, depending so from the stability constant, as well as from the kinetical factors. (11)

When in the (3) relation we introduce the notations:

$[ML] = Q_{ML}$  = the ligand amount binded in the total quantity applied in the sample, the relation become:  $K = Q_{ML} / [M_0](Q_{Lt} - Q_{ML})$  (4)

The  $Q_{ML}$  value were graphical determined from the positive peak area, measured by means of planimetry, from the basal line of the metal concentration in the eluent and were report to the standardisation area.

The standardisation area we chose in the horizontal portion of the elution profile, in the predecessor zone of the exclusion peak of the columns, at which correspond a known amount of the metal ion ( $\mu M$ ) contained in the respectively eluant volume.

The medium number of the binded metal ions by the ligand designate as n are given by the relation:  $n = Q_{ML}/Q_{Lt} = K[M]_0 / 1 + K[M]_0$  (5)

The n values of the stability constants and of the log K for the copper, zinc and iron with the diphosphate, tripolyphosphate and the high polymerised polyphosphate, determined by means of the sterical exclusion chromatography are shown in the table 1.

**Table 1.** Stability constants of the Fe, Cu and Zn complexes

| 1  | 2                 | 3   | 4     | 5    | 6     | 7                | 8    | 9     |
|----|-------------------|-----|-------|------|-------|------------------|------|-------|
| 1  | Fe-P <sub>2</sub> | 4,9 | 0,993 | 1    | 0,007 | $3.1 \cdot 10^6$ | 6.49 | 1,738 |
| 2  | Fe-P <sub>3</sub> | 4.9 | 0.972 | 1    | 0.021 | $1.0 \cdot 10^6$ | 6.02 | 1.959 |
| 3  | Fe-P <sub>n</sub> | 6.1 | 0.940 | 4.24 | 0.287 | $2.3 \cdot 10^5$ | 5,35 | 0.222 |
| 4  | Cu-P <sub>2</sub> | 4.8 | 0.860 | 1    | 0.140 | $1.3 \cdot 10^5$ | 5.11 | 1.720 |
| 5  | Cu-P <sub>3</sub> | 5.0 | 0.560 | 1    | 0.440 | $2.6 \cdot 10^4$ | 4.41 | 1.120 |
| 6  | Cu-P <sub>n</sub> | 5.0 | 0.360 | 8.47 | 5.702 | $1.3 \cdot 10^3$ | 3.10 | 0.043 |
| 7  | Zn-P <sub>2</sub> | 4.6 | 0.438 | 1.47 | 0.562 | $1.7 \cdot 10^4$ | 4.23 | 0.617 |
| 8  | Zn-P <sub>3</sub> | 5.9 | 0.660 | 1    | 0.340 | $3.3 \cdot 10^4$ | 4.52 | 0.660 |
| 9  | Zn-P <sub>3</sub> | 4.8 | 0.159 | 1    | 0.431 | $2.4 \cdot 10^4$ | 4.37 | 0.854 |
| 10 | Zn-P <sub>n</sub> | 4.6 | 2.294 | 8.47 | 0.933 | $5.4 \cdot 10^4$ | 4.73 | 0.271 |

The stability constants calculated for the mentioned complex ions with the diphosphate and triphosphate are comparable with the data from the bibliography sources, which are determined for the stability constants by means of the other physico-chemical methods (Bobtelski and Kertes, 1955, Martell and Calvin, 1956).

In the case of the metal complexes with the Kurrol salt, the stability constants are smaller as compared with these for the diphosphate and triphosphate. The calculated values of the n for the copper and iron complexes with the diphosphate and the triphosphate shows a metal-ligand ratio equal to 2:1, and for the zinc complexes with the same ligands, the ratio of 1:1.

In the case of the Kurrol salt complexes with the metal ions, the  $n$  values are very small.

Supposing that the molecular weight of the Kroll salt are  $10^6$  (Kouchiyama and Dreyer, 1978), the medium number of the binded metal ions to a  $P_n$  molecule (not to a  $PO_3$  unit), results in the case of the copper a value of 390; results in the case of the zinc of 161 and in the case of iron of 845. These values being smaller as compared with the 1440 value obtained of Yoza for the magnesium complex with the Kurrol salt.

For the iron complex with the Kurrol salt realized at the pH value of 2 for the eluant, results from the graph of the per cent concentration for the various diphosphate specimen as function of the pH value, calculated by means of the diphosphoric acid dissociation constants, results that the complexed iron species at this pH value are  $H_2P_2O_7$ .

The stability constants determination by means of the steric exclusion chromatography in gel, give the possibility to study also of the polyphosphate complexes with a condensation degree upper to the studied, and which may be obtained in very small quantities, only by the chromatography in column of the ion exchangers, quantities which are insufficients for the other usually determination methods.

## CONCLUSIONS

1. The sterical exclusion gel chromatography enable to study the forming of the metal ions complexes with the oligomeric polyphosphates also in the case of the transition metals with the trace element role in the biological systems.
2. The method insure a good separation of the complex compounds, by means of a judicious choose the type of the Sephadex gel, as function of the middle molecular weight of the ligands used for the complexation.
3. This method give the possibility to determine the stability constants values and the ratio between the metal and the ligand ratio.
4. To know the metal-ligand ratio in the complexes enable a indirect dosage of the polyphosphoric mixtures in solution, replacing the phosphorus dosage, fact which increased the speed, the sensibility and the precision of the dosage by mean of the metal ions from the automatic dosage by means of the eluated fractions by mean of the others methods, as well as the atomic absorption spectrography.

## REFERENCES

1. Bobtelski M, Kertes S. J.: Appl. Chem. 4, 419, 1954, 5, 125, 1955.
2. Forth W, Rummel W.: Physiological Rew. 53,724,1973.
3. Hummel Y. P., Dreyer W.Y.: Biochim. Biophys Acta. 63,530,1962.
4. Kouchiyama K, Yoza W., Ohashi S. J. : Chromatogr. 147, 271, 1978.
5. Martell A., Calvin M.: Chemistry of the Meta-Chelate Compounds. Prentice Hall. Inc. Englewood Cliffs. Ed..III. 1956.
6. Pfanstiel R., Iler R.K.: J. Am. Chem. Soc. 74, 6059, 1952.
7. Quimby Q. T., Mabis A.Y.:Lampe. Anal. Chem. 26, 661, 1954.
8. Atma Ram, Kumar S., Sinha B.C. : Indian J. of Chemistry. Vol. I. (6).

9. Spiro T. G., Allerton S.E., Renner Y., Terzis A., Bils. R. Saltman P. J. Am. Chem. Soc. 88, 8721.
10. Wirth H.E.: Ind. Eng. Chem. Anal. Ed. 14, 722, 1942.
11. Yoza N.: J. Chromatogr. 86, 325, 1973.
12. Yoza N.: J. Che. Educ. 54, 284, 1977.
13. Yoza N.: Kouchiyama, Miyajima T., Ohashi S. Anal. Lett. 8, 641, 1975.
14. Yoza N.: Ishibashi K., Ohashi S. J. Chromatogr. 134, 497, 1977.

# REFINED SUNFLOWER OIL: AN EXCELLENT VEHICLE FOR IODINE FORTIFICATION

Adina CREANGĂ<sup>1\*</sup>, Nathalie HENIN<sup>2</sup>, Mihaela SIMESCU<sup>3</sup>, Bruno DE BENOIST<sup>4</sup>

1. Bunge Romania, Pipera North Center, 1st floor, 2 //II Pipera Tunari, RO-014 251 Voluntari, Ilfov, Roumania; 2 "Bunge Europe" 13, Rue de Florissant, Geneve, Switzerland; 3. "National Institute of Endocrinology, C.I. Parhon, Bucharest, Blvd. Aviatorilor 34-36, Sector 1, RO-011 863, Roumania; 4. Bunge Switzerland, 13, Rue de Florissant, Geneve, Switzerland

\* E-mail: adina.creanga@bunge.com;

## ABSTRACT

Iodine is an essential element for normal physical and mental development. More than 2 billions individuals worldwide have insufficient iodine intake. Iodine deficiency disorders (IDD) is a public health priority not only as a main cause of preventable brain damage in children, but also because of its consequences on social and economic development through limitation of human resources potential. Salt iodization is the most common strategy for eliminating iodine deficiency and was mandatory or voluntary implemented in some countries. Because of increased concerns related to salt consumption as a risk factor to cardiovascular diseases, countries are encouraged by WHO to reduce salt consumption and identify other vehicles but salt for iodine fortification (WHO, 2008). A promising option is refined sunflower oil enriched in iodine. Given in complement to iodized salt, it can substantially contribute to increase iodine intake and therefore improve iodine status of the population. It also has the advantage over salt to provide other nutrient beneficial to health such as unsaturated fatty acids and vitamin E.

**Key words:** Iodine deficiency, food vehicle for iodine fortification, iodine status, iodized salt, refined vegetable iodized oil, unsaturated fatty acid

## INTRODUCTION

Regarding iodine deficiency the followings are mentioned :

- a) 40 million people are at risk of iodine deficiency in Europe (both Eastern and Western)
- b) In Europe, iodine deficiency disorders (IDD) is usually moderate or mild with limited manifestations such as poor pregnancy outcome, reduced physical growth or mental impairment: 10-15 IQ points may be lost in iodine-deficient communities!
- c) Subtle degrees of mental impairment as observed in mild forms of IDD may lead to poor school performance, reduced intellectual ability and impaired work capacity
- d) The fact that some countries have been successful in eliminating IDD over the last few years such as Romania does not preclude that some areas within the country might be still affected by moderate or even severe iodine deficiency.

- e) Even in countries that have achieved iodine sufficiency, the iodine nutrition status of pregnant and lactating women, and infants may still be inadequate. National commitments towards adequate iodine nutrition and its prevention is much weaker in most European countries than elsewhere in the world. According to WHO, "iodine deficiency and its many consequences are still a public health problem in much of Europe".

## **1. FOOD FORTIFICATION - BASIC PRINCIPLES**

### **1.1. Food fortification is a public health approach to the correction of micronutrients deficiencies**

Food fortification is justified where there is "a demonstrated need for increasing the intake of an essential nutrient in one or more population groups. This may be in the form of actual clinical or sub clinical evidence of deficiency, estimates indicating low levels of intake of nutrients, or possible deficiencies likely to develop because of changes taking place in food habits" (Codex Alimentarius, 1994).

The goal of fortification is to increase the micronutrients intake of an acceptable proportion of the population to an adequate level, without causing an unacceptable risk of excessive intake

### **1.2. Oil is an appropriate vehicle for food fortification because it:**

- Can be distributed widely to reach key target groups throughout the country.
- It is consumed daily in fairly constant amounts.
- Has no objectionable change in taste, appearance or color after fortification.
- Retains an appropriate level of the nutrient after further processing or cooking.
- Is not consumed in amounts that would expose to the risk of over consumption and therefore of iodine intakes beyond the safe level.
- Is an homogenous product

## **2. REFINED SUNFLOWER OIL ENRICHED IN IODINE:**

### **2.1. Product description :**

#### **Floriol Iod Protect launched in Romania in 2006**

Its Ingredients are: Refined sunflower oil; Ethyl esters of iodized fatty acids; The iodine concentration is 1 mg/kg and two tablespoons (20g) of Floriol Iod Protect bring 15% of daily recommended intake in iodine

### **2.2. Nutritional information**

A tablespoon of this product brings a few calories into the organism. It contains more unsaturated fatty acids than saturated ones and is rich in vitamin E. Details are given in Table 1.



**Table 1.** Composition of the iodine enriched sunflower oil

| Mean nutritional value      | 10 g (one table spoon) | 100 g   |
|-----------------------------|------------------------|---------|
| Energetic value             | 90 kcal or 370 kJ      | 3700 kJ |
| Proteins                    | 0                      | 0       |
| Carbohydrates               | 0                      | 0       |
| Fats                        | 10 g                   | 100 g   |
| Saturated fatty acids       | 1.2 g                  | 12 g    |
| Monounsaturated fatty acids | 2.6 g                  | 26 g    |
| Polyunsaturated fatty acids | 6.2 g                  | 62 g    |
| Cholesterol                 | 0 g                    | 0 g     |
| Fibers                      | 0 g                    | 0 g     |
| Sodium                      | 0 g                    | 0 g     |
| Vitamin E                   | 7.0 mg                 | 70 mg   |
| Iodine                      | 10 mcg                 | 100 mcg |

### **2.3. Health claims on Floriol Iod Protect**

“Recent studies shows that IDD is a worldwide public health problem, including in Romania.

Iodine is an important element for maintaining health, contributing to healthy physical and mental development.

Iodine is an essential nutrient during the overall life cycle especially in adolescent girls as future mothers, in pregnant and lactating women and in young children. Iodine deficiency may result in irreversible brain damage during fetal development and infancy.

## **CONCLUSIONS**

### **The refined sunflower oil enriched in iodine:**

1. Has a similar behavior as regular sunflower oil during storage and cooking;
2. Due to covalent bonds iodine-fatty acids, is very stable during cooking and frying; the losses are much smaller comparing with iodized salt ( aprox.10%).
3. Is a complementary source of iodine for optimum iodine nutrition especially in pregnant women and young children;
4. Is a naturally source of essential fatty acids and vitamin E;

## REFERENCES

1. de Benoist B., Andersson M., Egli I., Takkouche B., Allen H. (ed) - Department of Nutrition for Health and Development, World Health Organization, Geneva
2. Delange F. - The disorders induced by iodine deficiency, *Thyroid*, 1994, 4, 107-128
3. Delange F., Hetzel B. - The iodine deficiency disorders, in: *The thyroid and its diseases*, Chapter 20, <http://www.thyroidmanager.org/thyroidbook.htm>, 2004.
4. Simescu M., Varcui M., Nicolaescu E., Gnat D., Podoba J., Mihaescu M., Delange F. - Iodized oil as a complement to iodized salt in schoolchildren in endemic goiter in Romania, *Horm. Res.*, 2002, 58 (2), 78-82.
5. Zimmermann - Iron status influences the efficacy of iodine prophylaxis in goitrous children in Cote d'Ivoire, *Int. Vitam. Nutr. Res.*, 2002, 72, 19-25.
6. \*\*\*WHO/UNICEF/ICCIDD - Assessment of iodine deficiency disorders and monitoring their elimination, Geneva, World health Organization, 2001.
7. \*\*\*WHO - Iodine status worldwide, WHO Global database on iodine deficiency, Publ. by WHO Geneva, 2008.

# PHOSPHONATE LIGANDS GRAFTED ON STYRENE-DIVINYLBENZENE COPOLYMERS USED IN METAL IONS SEPARATION FROM WATER

Corneliu-Mircea DAVIDESCU<sup>1</sup>, Adriana POPA<sup>2</sup>, Petru NEGREA<sup>1</sup>,  
Geza BANDUR<sup>1</sup>, Gheorghe ILIA<sup>2</sup>, Radu ARDELEAN<sup>1</sup>

\* E-mail : corneliu.davidescu@rektorat.upt.ro

1 Faculty of Industrial and Environmental Chemistry University „Politehnica” of Timișoara, 6, V. Pârvan Blv., RO-300223, Timișoara, Roumania; 2. Institute of Chemistry, 24, Mihai Viteazul Blv., RO-300223, Timișoara, Roumania

## ABSTRACT

*Phosphonates grafted on styrene-divinylbenzene supports were prepared and their use in Cu(II) separation from aqueous solutions was investigated. The polymeric materials were characterized by determination of the functionalization degree based on the phosphorus content and thermal stability by thermogravimetric analysis. The sorption capacity of the resins functionalized with phosphonate pendant groups towards Cu(II) was studied.*

**Keywords:** phosphonates, styrene-divinylbenzene microbeads, removal of metal ions

## INTRODUCTION

The synthesis of new polymeric matrix based sorbents, incorporating ligands which are immobilized by chemical bonds and are highly selective towards metal ions, received lately great interest due to their potential use in the industrial separation and concentration of metals and in trace metal analysis.

Nonspecific sorbents, such as activated carbon, metal oxides, silica and ion-exchange resins have been used (Reed B.E., 1993). Specific sorbents consist in ligands (e.g. ion-exchange materials or chelating agents) interacting specifically with the metal ions, and a carrier matrix which may be an inorganic material (e.g. aluminum oxide, silica or glass) or polymer microbeads (e.g. polystyrene, styrene copolymers, cellulose, poly(maleic anhydride) or polymethylmethacrylate (Kantipuly et al., 1990; Goto et al., 1993; Popa et al., 2003; Popa et al., 2004; Popa et al., 2007; Popa et al., 2008).

Classes of ligands that have been developed for this purpose include amines, amidoximes, phenols and their derivatives, and phosphonates. Phosphoric acid derivatives are especially versatile, displaying a wide range of ionic selectivities in direct relationship with the ligand structure.

The determination of Cu(II) in natural waters is important whereas this trace element constitutes an environmental problem and there is a lack of information on the behaviour and toxic potential of this metal on the metabolism of living organisms (Korn M.G.A. et al., 2004).

In this work we describe the multiphase system synthesis of phosphonates grafted on styrene-divinylbenzene copolymer-supports and their characterization (degree of functionalization and thermal stability).

The aim of this work was to assess the possibility of application of chelating resins obtained by functionalization of styrene-divinylbenzene copolymers with phosphonate groups in copper separation and preconcentration. Atomic absorption spectrometry is nowadays the most common method for trace metal determination. The quantitative analysis of metal ions was carried out by atomic adsorption spectroscopy on SpectrA 110 coupled with GTA-110.

## MATERIALS AND METHODS

The functionalization with phosphonate pendant groups was realized by the polymer-analogous Michaelis-Becker reaction presented in Scheme 1.




**Fig. 1.** Michaelis-Becker polymer-analogous functionalization reaction.  
Phosphite used: (CH<sub>3</sub>O)<sub>2</sub>POH, (C<sub>2</sub>H<sub>5</sub>O)<sub>2</sub>POH, (i-C<sub>8</sub>H<sub>17</sub>O)<sub>2</sub>POH, (C<sub>2</sub>H<sub>5</sub>-C<sub>6</sub>H<sub>12</sub>O)<sub>2</sub>POH

### *Synthesis of the phosphonates grafted on styrene-7% divinylbenzene copolymers*

The synthesis of the phosphonates grafted on a macroporous styrene-7%divinylbenzene copolymer-support was performed by the method previously described (Popa A. et al., 2003; Popa A. et al., 2004). A 5-g sample of chloromethylated styrene-7%divinylbenzene copolymer and 100 ml of solvent (1,4 - Dioxane) were added into a round bottom flask fitted with reflux condenser, mechanical stirrer and thermometer. Then, the necessary amounts of phosphite and sodium carbonate were added in order to achieve the molar ratio of chloromethyl:diphosphite:sodium carbonate = 1:1.5:1.5 Tetramethylammonium iodide was used as catalyst. The mixture was maintained under stirring for 24 h at 30°C. After cooling, the copolymer beads were separated by filtration, washed with ethanol (2x50 mL), 1,2-dichloromethane (2x50 mL) and ethyl ether (2x50 mL), and dried at 50°C for 24 h.

### *Procedure for Cu(II) removal from aqueous solutions using phosphonates grafted on polymeric supports (Popa A. et al., 2007; Popa A. et al., 2008)*

Cu(II) ion separation from aqueous solutions was performed by a batch technique at room temperature. 0.5 grams of styrene-7%divinylbenzene copolymer functionalized with phosphonate pendant groups were added to a volume solution containing 100 mg Cu(II) in 0.1 N HNO<sub>3</sub> and maintained under mechanical stirring for 7 hours. The residual Cu(II) content in solution was determined on samples (1 mL) taken from the mixture as follows: at 10 min. interval in the first hour, at 15 min.

interval in the second hour and at 30 min. interval in the last 5 hours. Each sample was diluted (ratio 1:25) and the concentration of Cu(II) was determined by atomic adsorption spectroscopy on SpectrA 110 coupled with GTA-110.

## RESULTS AND DISCUSSIONS

The main characteristics of the copolymer functionalized with phosphonates groups are presented in Table 1.

**Table 1.** Characteristics of the styrene-7%divinylbenzene copolymers grafted with phosphonate groups obtained by polymer-analogous Michaelis – Becker reaction.

| Product code | Phosphite                                                                          | P (%) | G <sub>F</sub> (mmoles phosphonate/g of copolymer) |
|--------------|------------------------------------------------------------------------------------|-------|----------------------------------------------------|
| Me-PS        | (CH <sub>3</sub> O) <sub>2</sub> POH                                               | 5.49  | 1.78                                               |
| Et-PS        | (C <sub>2</sub> H <sub>5</sub> O) <sub>2</sub> POH                                 | 6.09  | 1.98                                               |
| i-Oct-PS     | (C <sub>8</sub> H <sub>17</sub> O) <sub>2</sub> POH                                | 4.37  | 1.40                                               |
| EtHex-PS     | (C <sub>2</sub> H <sub>5</sub> -C <sub>6</sub> H <sub>12</sub> O) <sub>2</sub> POH | 4.13  | 1.33                                               |

One of the key properties of a polymeric chelating agent used in metal ions separation is its thermal stability. The thermal stability of the original copolymer support (PS) and that of the copolymers functionalized with phosphonate groups was evaluated by thermogravimetric analysis on a Netzsch TG 209, in nitrogen atmosphere, at a heat rate of 5K/min from 20°C to 600 °C. The results are presented in Table 2.

**Table 2.** Weight loss of the polymeric materials

| Product  | Weight loss (%) |          |          | T <sub>inflexion</sub> (°C) |
|----------|-----------------|----------|----------|-----------------------------|
|          | 20-200°C        | 20-300°C | 20-400°C |                             |
| PS       | 3.25            | 14.25    | 23.72    | 427.7                       |
| Et-PS    | 4.57            | 11.20    | 94.72    | 366.7                       |
| i-Oct-PS | 5.27            | 11.91    | 44.60    | 399.1                       |
| EtHex-PS | 5.11            | 17.99    | 40.85    | 404.3                       |

As can be seen the weight loss up to 200°C is minor, indicating a thermal stability qualifying the products for use in small and large scale separation processes. This type of behavior is characteristic for styrene-divinylbenzene copolymers. At higher temperatures the weight loss is larger indicating significant degradation of the polymeric network. The thermal stability is affected by the nature of the phosphonate ligands. In all cases, a decrease of the thermal stability was observed as a result of the functionalization with phosphonate pendant groups in comparison with the starting chloromethylated copolymer.

The ability of the functionalized copolymers to act as specific sorbents/chelating agents for Cu(II) ions from aqueous acidic solutions (pH=1) was investigated.

The concentrations of Cu(II) (mg/L) in solution were determined by atomic absorption spectrometry on a Varian SpectrAA 110 coupled with GTA-110, at the wavelength  $\lambda = 324,8$  nm.

The amount of Cu(II) separated, q, was calculated as:

$$q = \frac{C_0 - C_t}{G} \cdot V \cdot 10^{-3} \quad \text{mg Cu(II)/g of copolymer}$$

where:  $C_0$  is the initial concentration of Cu(II) in solution (mg/L);  
 $C_t$  is the concentration of Cu(II) in solution after  $t$  minutes (mg/L);  
 $V$  is the volume of the solution (mL);  
 $G$  is the mass of resin sample (g).

The adsorption rates of Cu(II) ions by the polymer-grafted phosphonates with different functional groups are shown in Figure 3.




**Fig.3.** Specific adsorption capacity as a function of time, for the resins bearing phosphonate groups, at pH=1 and 25 °C.

As can be seen, the kinetic curves are Langmuir type isotherms, suggesting adsorption in monomolecular layer followed by saturation of the active centers. Such a behaviour was expected due to the fact that water is not a good swelling solvent for the polymer support. In such conditions only the active centers located at the surface of the polymer beads or, eventually, in the larger pores are available.

The amount of Cu(II) ions adsorbed per mass unit of copolymer (*ie.* adsorption capacity) increase with the exposure time and finally reach a constant value corresponding to the saturation of the active centers located in the porous structure of the polymer beads.

The total adsorption capacity of the resins functionalized with phosphonate groups (in mg. Cu(II)/g of copolymer) was: 2.90 mg/g for Me-PS, 2.78 mg/g for Et-PS, 3.58 mg/g for i-Oct-PS and 2.77 mg/g for EtHex-PS and was attained in all cases after around 360 minutes of exposure.

The efficiency of metal retention ability (mmoles Cu(II)/mmole phosphonate) of the functionalized polymers was found to depend on the nature of the phosphonate group, as shown in Figure 4.




**Fig. 4.** Efficiency of the retention ability of the polymers bearing phosphonate pendant groups (mmoles Cu(II)/mmole phosphonate).

The efficiency was better in case of the phosphonates with large, bulky or branched alkyl groups. The best results were obtained with the copolymer functionalized with iso-octylphosphonate groups.

## CONCLUSIONS

All the copolymers functionalized with phosphonate groups reported in the present study were proved to be efficient sorbents for Cu(II) ions from aqueous solutions.

The polymer-grafted chelating agents investigated are thermally and mechanically stable.

The specific retention capacity of Cu(II) was suggesting classical monomolecular layer adsorption.

The efficiency of metal retention by the functionalized polymers was found to depend on the nature of the phosphonate group.

Further studies will focus on the retention capacity of polymer-grafted phosphonates towards other types of metal ions and evaluation of the polymeric sorbents in column separations.

## REFERENCES

1. Goto A., Morooka A.S., Fukamachi M., Kusakabe K., Kago T.: Sep. Sci. Technol., 28, (1993), 2229.
2. Kantipuly C., Katragadda S., Chow A., Goser H.D.: Talanta, 37, (1990), 491.
3. Korn M.G.A., Santos Jr. A.F.: J.Braz. Chem.Soc., 2004, 15(2), 212-218
4. Popa A., Ilia Gh., Iliescu S., Davidescu C.M., Pascariu A., Bora A.: Rev. Chim., 54(10), (2003), 834.
5. Popa A., Ilia Gh., Iliescu S., Davidescu C.M., Macarie L., Plesu N.: Rev. Chim., 55(6), (2004), 456.
6. Popa A., Davidescu C.M., Negrea P., Bandur G.: Proceedings: New trends and strategies in the chemistry of advanced materials with relevance in

biological systems, technique and environmental protection, (2007), 127  
(ISBN 978-973-52-0217-0)

7. Popa A., Davidescu C.M., Negrea P., Ilia Gh., Katsaros A. , Demadis K.D.:  
Ind. Eng. Chem. Res. (2008), 47, 2010-2017
8. Reed B.E.,Matsumoto M.R.: Sep. Sci. Technol., 28, (1993), 2179.



# DEVELOPMENT OF FUNCTIONAL FOODS FROM SOY EXTRACTS ENRICHED WITH MAGNESIUM IN BREAST CANCER

Simona DRĂGAN<sup>1\*</sup>, Carmen SOCACIU<sup>2</sup>, Iosif GERGEN<sup>3</sup>, Lavanya YOGI<sup>1</sup>,  
Cristina A. DEHELEAN<sup>1</sup>

\* E-mail: sdragan@cardiologie.ro

1. *Department of Medical Policlinics, University of Medicine and Pharmacy "Victor Babes", Eftimie Murgu Square no. 2, RO-300041 Timisoara, Roumania;* 2 *Faculty of Agriculture, University of Agricultural Sciences and Veterinary Medicine, Calea Manastur no.3-5, RO-400 372 Cluj-Napoca, Roumania;* 3. *Department of Analytic Chemistry, Faculty of Food Products Technology, University of Agricultural Sciences and Veterinary Medicine of Banat, Calea Aradului nr. 119 RO-300 645 Timisoara, Roumania*

## ABSTRACT

*Reactive oxygen species (ROS) play a key role in pathogenesis of degenerative diseases. Evidence offered by antioxidant dietary supplements is intriguing. Vitamin E, ascorbic acid and carotenoids were inversely associated with cardiovascular and cancer mortality in some studies, while negative results have been reported from intervention trials on beta-carotene and vitamin E. Metabolically, nutrients act in combination at physiologic doses, whereas pharmacological doses of a single nutrient, like those used in the above studies, may provoke a nutritional imbalance. Pharmacologic doses of a single carotenoid, such as beta-carotene, can result in competition at the receptor site with the other 50 naturally occurring dietary carotenoids and thus inhibition of their protective effects. The available data justify the conclusion that a balanced mixture of natural antioxidants provides the best protection against cancer and cardiovascular disease. Integral antioxidative capacity and antiradical properties of various extracts from plants can be measured by DPPH (11-diphenyl-2-picril-hydrazyl) scavenging test in order to obtain the most potent mixtures. These can be incorporated as bioactive components in functional foods when prescribing healthy diets. We have studied four soy extracts in order to find the most potent one to be incorporated in foods prepared for patients with breast cancer, preferablz breakfast cereals with high magnesium content by adding nuts and raisins and and dried buckthorn berries for high vitamin C and E content.*

**Key words:** functional foods, soy extracts, breast cancer

## INTRODUCTION

According to data officially reported by the WHO Regional Office in 2002, cancer is responsible for close to one fifth of all deaths, ranging to about 30% in Western European countries (13). The most common cause of death from cancer in males is lung cancer, while in females it is breast cancer. Over the last 150 years, fat

consumption has increased substantially in Western and Eastern European countries, which also coincides with a dramatic increase in cancer incidence. A review of epidemiological studies strongly suggests an inverse relationship between the consumption of fresh fruits and vegetables and the incidence of cancer. Fruits and vegetables are rich in fiber, antioxidants, and other plant-derived substances, or phytonutrients, that are thought to have cancer-protective properties (1).

Psychological stress effects, which are often underestimated, should be taken into account among patients bearing free radical-mediated diseases. Psychological stress gives rise to a prooxidant state. Radical oxygen species (ROS) are implicated in mediation of all degenerative diseases, including the atherosclerotic process of coronary heart disease, cancer and neurodegenerative diseases (7). Preservation of key target functions, in the reversible phase of promotion of carcinogenesis, is the major strategic goal of defense against ROS. The human diet contains an array of different compounds that possess either antioxidant activities or the apparent ability to scavenge ROS (11). Examples of opportunities for modulation of target function by candidate food components, as well as possible markers related to defence against ROS are summarized in Table 1.

**Table 1.** Opportunities for modulation of target function by food components in the defense against ROS

| <b>Target functions</b>                                                           | <b>Possible markers</b>                                         | <b>Candidate food components</b>                                                  |
|-----------------------------------------------------------------------------------|-----------------------------------------------------------------|-----------------------------------------------------------------------------------|
| Preservation of structural and functional activity of DNA                         | Measurement of damaged components of DNA                        | Vitamin E, vitamin C<br>Carotenoids, polyphenols including flavonoids             |
| Preservation of structural and functional activity of polyunsaturated fatty acids | Measurement of hydroperoxides or lipid derivatives              | Vitamin E, vitamin C<br>Carotenoids, polyphenols including flavonoids             |
| Preservation of structural and functional activity of circulating lipoproteins    | Measurement of hydroperoxides and oxidized apoproteins of lipid | Vitamin E, vitamin C<br>Carotenoids, polyphenols including flavonoids             |
| Preservation of structural and functional activity of proteins                    | Measurement of damaged proteins or components                   | Vitamin E, vitamin C<br>Selenium<br>Carotenoids, polyphenols including flavonoids |

Reproduced from: Consensus Document: Scientific Concepts of Functional Foods in Europe, BJ Nutrition, 81, 1999 (2).

Diet is estimated to contribute to up to 50% of all newly diagnosed breast cancer cases (3,10). Consumption of phytoestrogens, particularly soy products, as well as vegetables, is higher in Asia than in the Western world. Soy-based diets are high in genistein (4, 5, 7-trihydroxyisoflavone), which has been widely studied for its potential anticancer properties (4,5,9,12). Daidzein and genistein are isoflavone phytonutrients derived from soybeans. Laboratory tests have found that selenium and magnesium reduce the incidence of new breast cancers when taken with adequate dietary supplies of vitamins A, C, and E.

The aims of our research were:

1. Development of functional foods from soy matrixes with nuts and raisins for high magnesium content
2. Analysis of antioxidant activity of functional foods (DPPH)

## **EXPERIMENTAL PART**

### **Obtaining the extracts from soybeans.**

Four types of soybeans extract were used:

- degreased soy extract containing glucosylated isoflavonoids,
- hydrolyzed soy extract containing isoflavonoid aglycons,
- extract of hydrolyzed soy pellets containing isoflavonoid aglycons,
- non-degreased soy extract obtained through the following process: 50g of ground soybeans were mixed with 150ml methanol solution (120 ml MeOH, 30 ml H<sub>2</sub>O, 0,2 ml HCl concentrate) and then centrifuged. Supernatant was used as an extract.

### **Obtaining the soy texturate**

The soy texturate sample was obtained as follows: 10g of sample with 50 ml 0.1% HCl in methanol 80% were sonicated for 60 minutes, after which the extract was filtered and its volume measured. 50ml more extraction solvent were added to the sample, 60 minutes sonication, then filtering. The two volumes of solvent cumulated, and the final extract volume was concentrated on rotavapor half way through. For all samples, the UV-Vis spectrum within the range of 200-600nm was recorded.

### **Obtaining the functional food as breakfast cereal**

The soy texturate was enriched with raisins for higher magnesium content and dried buckthorn berries for vitamin C and E content.

### **Determining the antioxidant activity by the DPPH method**

One of the highlighting methods for the *in vitro* antioxidant activity, namely the one of flavonoids, is the DPPH method. It consists of the ability of antioxidants (flavonoids) to eliminate the free radical DPPH<sup>•</sup> (11-diphenyl-2-picril-hydrazyl). This is a paramagnetic free radical, relatively stable, which accepts an electron from a hydrogen radical in order to form a stable diamagnetic molecule (Diagram 1). The antioxidant flavonoids donate the hydrogen, thus reducing the radical, a phenomenon perceived through the changing in colour from violet to yellow. The decrease in optical density within the characteristic wavelength range during the radical-antioxidant reaction is due to the fact that the DPPH<sup>•</sup> radical absorbs at 515nm, while after it being reduced, this absorption disappears.

The activity of *in vitro* free radical scavenging by use of the DPPH method was at first determined with flavonoid standards ( quercetine, genistein and daidzein) then compared to the soy extracts.

Chemical reagents and standards used:

- 2,2-dyphenil-1-picril-hydrazyl (DPPH)-0,031g of DPPH reagent were dissolved into 99,969ml of distilled water ( a concentration of 80µM DPPH was obtained). Then 1 ml of this solution was mixed with 9ml methanol and sonicated in order to produce stable free radicals.
- Genistein - the following concentrations were used: 500 ppm, 100 ppm and 50 ppm.

- Daidzein - the following concentrations were used: 4mg/100 ml; 2 mg/100 ml.
- Quercetine - the following concentrations were used: 2mM, 1mM, 0,5mM and 0,25mM.

Method: 2,8ml of the DPPH radical solution were mixed in turn with 400 µl of isoflavone standards ( genistein and daidzein) of different concentrations, and with the soy extracts. The mixture was energetically agitated then kept in the dark for 30 minutes, after which the absorbance at 515nm was measured with a spectrophotometer (JASCO V-530). For the blank, 2,8ml of DPPH radical solution were used in mixture with 400 µl of methanol.

The degree of discoloration of DPPH (%) was calculated according to the following formula:

$$\% \text{discoloration} = [1 - A_{\text{sample}} / A_{\text{blank}}] \times 100$$

Where:

$A_{\text{blank}}$  – absorbance of the blank solution;

$A_{\text{sample}}$  – absorbance of standard solution and soy extracts with DPPH after 30 minutes.

## RESULTS AND DISCUSSIONS

The degree of discoloration of the DPPH by the isoflavonoid standards (genistein and daidzein), as well as the soy extracts, are all presented in Table 2.

**Table 2 .** Values of absorbance ( $A_{\text{sample}}$  and  $A_{\text{blank}}$ ) and degree of DPPH discoloration for standards and soy extracts.

| Specification                     | Dilutions    | * $A_{\text{sample}}$ | $A_{\text{blank}}$ | %discoloration DPPH |
|-----------------------------------|--------------|-----------------------|--------------------|---------------------|
| Genistein                         | 500 ppm      | 0,4450                | 0,6520             | 31,74               |
|                                   | 100 ppm      | 0,5246                |                    | 19,53               |
|                                   | 50 ppm       | 0,5311                |                    | 18,54               |
| Daidzein                          | 4 mg/ 100 ml | 0,4787                | 0,6520             | 26,5                |
|                                   | 2 mg/100 ml  | 0,5554                |                    | 14,81               |
| Quercetine                        | 2mM          | 0,0426                | 0,6596             | 93,54               |
|                                   | 1mM          | 0,0315                |                    | 95,22               |
|                                   | 0,5mM        | 0,0281                |                    | 95,73               |
|                                   | 0,25mM       | 0,0268                |                    | 95,93               |
| Degreased soy extract             |              | 0,1535                | 0,6271             | 75,52               |
| Non-degreased soy extract         |              | 0,1393                | 0,7112             | 80,41               |
| Hydrolyzed soy extract            |              | 0,0967                | 0,6271             | 84,57               |
| Extract of hydrolyzed soy pellets |              | 0,2121                |                    | 66,17               |

\* absorbance after 30 minute

The DPPH method is used for investigating the activity of radical scavenging of several natural compounds such as the fenolic compounds, or by some plant extracts. The DPPH radical is being scavenged by the antioxidants through their capacity of giving away one hydrogen, with DPPH-H formation. The change in colour from dark violet to yellow after reduction can be quantified by the decrease of absorbance to 515nm.




**Fig. 1.** Various degrees of DPPH discoloration by the four soy extracts studied

In figure 2, the various degrees of DPPH discoloration by the soy extracts are presented comparatively. The highest degree of DPPH discoloration was shown by the hydrolyzed soy extract. This extract was characterized by TLC, by highlighting the presence of genistein and daidzein. The capacity of DPPH radical inhibition by the isoflavonoid aglycons is very small in comparison with the one of the extracts containing these isoflavonoids.

While characterizing TLC extracts, other unidentified compounds were highlighted, which could be responsible for the high antioxidant activity of the extracts. Moreover, the degreased soy extract is shown to have less antioxidant activity than the non-degreased one. Through degreasing, phytosterols – bioactive substances with antioxidant activity – are being eliminated.

The antioxidant activity also depends upon the concentration and structure of isoflavonoids. For example, the glucose binding in isoflavonoid aglycons will result in a decrease of antioxidant activity (6).

The soy beans contain approximately 2% isoflavonoids in the form of aglycons, thus the non-hydrolyzed extracts will present a less intense antioxidant activity than the hydrolyzed ones.

#### **The relationship between the chemical structure of quercetine and isoflavonoids and their antioxidant activity.**

In Table 3, on the first column, the structural characteristics that need to be fulfilled by a flavonoid in order to function as an antioxidant are being shown, while in the other columns the structural characteristics fulfilled by the studied flavonoids (quercetine, genistein and daidzein) will be ticked.

**Table 3.** Comparison between the structural characteristics of quercetine and isoflavonoids regarding their antioxidant capacity

| <b>Structural characteristics of flavonoids in free radical scavenging</b> | <b>Quercetine</b> | <b>Genistein</b> | <b>Daidzein</b> |
|----------------------------------------------------------------------------|-------------------|------------------|-----------------|
| Cathecol group in ring B                                                   | √                 | -                | -               |
| Pyrogallol group in ring B                                                 | -                 | -                | -               |
| The double bond between carbon atoms 2 and 3                               | √                 | √                | √               |
| Ketonic group in position 4                                                | √                 | √                | √               |
| -OH group in position 3                                                    | √                 | -                | -               |
| -OH group in position 5                                                    | √                 | √                | -               |
| -OH group in position 7                                                    | √                 | √                | √               |

Out of the three flavonoids studied here, quercetine fulfills almost all conditions (with the exception of the presence of pyrogallol group in ring B) in order to be considered a powerful antioxidant, followed by genistein and daidzein. In Table 4 it shows that quercetine has the highest capacity of DPPH discoloration (ranging from 95,93% to 93,54%, according to concentration levels). Genistein fulfills only 4 out of the 7 conditions necessary in order to be a good antioxidant, also reflected in the degree of DPPH discoloration (ranging from 31,74% to 18,54%, according to concentration levels). The weakest antioxidant is daidzein, as it fulfills only 3 out of the 7 necessary conditions, with a DPPH discoloration degree between 14,81% and 26,5%.

### HPLC chromatographic analysis of soy samples

Genistein being the most hydrophile, it will easily absorb into the thin intestine as aglycon after the hydrolysis of glycosilate conjugate, with exceptional therapeutical effects: estrogen-like activity, breast, colon or prostate cancer prevention, antioxidant activity, prevention of premenopause and osteoporosis symptoms as well as cardiac diseases (8).

The analysis of isoflavones is usually done with the help of chromatographic methods: thin layer chromatography and HPLC -High Performance Liquid Chromatography (Figure 3).

The genistein quantity oscilates between 15-45 mg/100 g, while the daidzein will range between 9-30 mg/100 g. The Daidzein:Glycitin:Genistein ratio can oscilate according to the soy variety from 1.2:1:2.1 to 35.4:1:60.4 .

### UV-VIS Spectrophotometric analysis of soy samples

Figure 4 presents the comparative image of the spectras at different dilutions of the soy texturate. Specific peaks ensuring the recognition of characteristic phytochemical compounds (spectral markers) for 250-265/abs.2,4 bandwidth (amino-acid-peptides) were identified.




**Fig. 2.** HPLC chromatogram of a soy extract: 1- genistein; 2-glycitin; 3-daidzein.




**Fig. 3.** Comparative image of spectra at different dilutions of soy texturate

## CONCLUSIONS

1. Based on data in the literature, it appears that genistein can act as both an estrogen and an antioxidant and an anti-proliferative agent. These effects may be both dose and tissue dependent. Incorporation of natural active components with therapeutic effects in functional foods, based on results of multidisciplinary research will be a challenge for both food and drug industries.

2. It is clear from epidemiologic data that Asian women living in Asia (where a diet high in soy is consumed) have a decreased risk for breast cancer. It is also

possible that soy contains other factors besides genistein that oppose the estrogenic effects of genistein and actually reduce breast cancer risk. Additionally, other environmental or lifestyle factors may be more related to the low breast cancer risk seen in this population than soy/genistein consumption. Fortunately, genistein continues to be an active area of research interest, and therefore an explanation for the dual nature of genistein may not be too far away.

3. The adding of soy flours to the wheat flour for bread as a functional food, as well as the use of soy texturate incorporated in cereal flakes enriched with nuts, raisins and buckthorn berries will give them a pleasant, nutty taste, a brownish colour and will maintain humidity for longer; also, as soy is gluten-free, it is highly recommended to women who are allergic to gluten. The volume of the bread prepared with soy will be less, but the freshness will maintain for longer.

**Acknowledgments** : This investigation was supported by grant CNCSIS 275/2005

## REFERENCES

1. Cassileth, B.R., Lucarelli C.D.: Herb-Drug Interactions in Oncology. Hamilton, Canada: B. C. Decker, 2003, pp.1.
2. Huang M.T., et al. : Canc Res., 51, 1991, pp.813.
3. Izzo A.A., Ernst E. : Drugs, 61, 2001, pp.2163.
4. Kienle G.S., Berrino, F., Bussing, A. et al. : Eur J Med Res, 8, 2001, pp.109.
5. Kiuchi, F., et al. : Chem Pharm Bull, 40, 1992, pp.387.
6. Labriola D, Livingston R. : Oncology (Huntingt) 13, 1999, pp.1003.
7. Lamartiniere C.A., Murrill W.B., Manzolillo P.A., Zhang J.X., Wei H., Brown N.M. Proc. Soc. Exp. Biol. Med., 217, 1998, pp.358.
8. Pierce J.P., Faerber S., Wright F.A. et al.: Control. Clin. Trials, 23, 2002, pp.728.
9. Ruby A.J., et al. : Cancer Lett., 94, 1995, pp.79.
10. Susan M., Rao, M.N.A.: Arzneim-Forsch/Drug Res. 42, 1992, pp.962.
11. Wattenburg L.W. :Can. Res. (suppl) 52, 1992, pp.2085s.
12. Young I.S., Woodside J.V.: J. Clin. Pathol. 54, 2001, pp.76.
13. \*\*\*Consensus Document: Scientific Concepts of Functional Foods in Europe, BJ Nutrition, 81, 1999.
14. \*\*\*WHO Europe. European Health Report 2002, WHO Regional Publications, European Series No 97, Copenhagen



# BIOLOGIC EVALUATION OF PLATINUM AND PALLADIUM COMPLEXES USING THE STATIC CYTOMETRY

Eva FISCHER-FODOR <sup>1\*</sup>, Natalia MIKLASOVA <sup>2,3</sup>, Piroska VIRAG <sup>1</sup>, Corina TATOMIR <sup>1</sup>,  
Ioana BRIE <sup>1</sup>, Maria PERDE <sup>1</sup>, Valentin CERNEA <sup>1</sup>, Luminita SILAGHI-DUMITRESCU <sup>3</sup>,

1. Research Department, I. Chiricuță Cancer Institute, Str. Republicii Nr. 34-36, RO-400 015 Cluj-Napoca, Roumania;

2. Faculty of Pharmacy, Comenius University, SK-81806 Bratislava, Slovakia; 3. Department of Chemistry, Faculty of Chemistry and Chemical Engineering, Babes Bolyai University, RO-400 028 Cluj Napoca, Roumania

\* E-mail: fischer.eva@iocn.ro

## ABSTRACT

*The present paper examines the CellScan technique utilization in cancer research in order to evaluate the antiproliferative activity of novel synthesized metal-based compounds and with the purpose to improve treatment of malignant diseases with standard platin-based drugs. The CellScan is a laser scanning static cytometer which allows repetitive spectroscopic measurements in fluorescent marked cells. It was demonstrated the potential of CellScan technology to detect the in vitro effects of the chemotherapy on tumor cells, making this method a valuable tool for chemosensitivity tests. The detection of cell viability and apoptosis is possible based on the transformations which occur in the cytoplasm matrix of tumoral cells influenced by cytotoxic compounds. The measurement of fluorescence changes due to this phenomenon is possible with the CellScan system. The potential of this technology to detect the in vitro effects of the inhibitory molecules on tumor cells was demonstrated, making this method a valuable tool for chemosensitivity tests.*

**Key words:** chemosensitivity, CellScan, fluorescence polarization, platinum complexes, cytotoxicity

## INTRODUCTION

Metal-based chemotherapeutic agents, above all platinum complexes, are an important class of antitumour compounds that have had significant impact on the treatment of a variety of tumours for nearly 30 years. The unique efficiency observed with standard platinum-based drugs such as cisplatin, *cis*-diamminecyclobutanedicarboxylato platinum(II) (carboplatin) and 1,2-diaminocyclohexaneoxalato platinum(II) (oxaliplatin) in cancer treatment fuelled the development of new platinum analogues (DeVita, 2004). In the last decade, it was demonstrated that several newly synthesised platinum compounds exhibit biological activity against tumoural cells, and their antimitotic properties were proved by several procedures (Fukuda, 1995; Barbara, 2006; Carland, 2005; Akamatsu, 1993; Gay, 2005; Robillard, 2003; Cemazar, 2006).

One of the major objectives of the cancer research is to find the best therapeutically scheme for the cancer patient and in this trend we performed chemosensitivity tests "ex vivo" on the tumoral cells harvested from the patients in order to establish the most efficient drug among the compounds recommended by the international treatment protocols and to improve the remission of the disease and the survival of the patient.

Another important goal is to develop new compounds with antiproliferative effect and to predict their prospective application as anticancer drugs. We synthesised and we tested "in vitro" three platin complexes and three palladium complexes with organometal ligands with the intention to characterize their antimitotic potential. Tertiary arsine complexes of platinum and other metals were recently reported (Moldovan, 2008; Phadnis 2005; Kwong, 2005) but there is no information regarding their biological activity. Among phenylarsines, information on biological activity was reported for phenylarsine oxide and dichlorophenylarsine (Frost, 1990; Geran, 1972; Takahashi, 2003; Begun, 1994; Verspohl, 2006), and the phenylarsine compounds attached to biomolecules like peptides and enzymes were studied (Schmidt, 2007; Wang, 2007). Phenylarsine oxide derivatives are well known as anti-trypanosome agents and they are used as drugs (Mottram, 1985; Fairlamb, 1989; Berger, 1994). We can achieve both aims using the static cytometry. The detection of fluorescence changes due to this phenomenon is possible with the CellScan system. The

specificity and the sensitivity of the test made by CellScan are greater than those with other methods. Simplicity, rapidity and non-invasive character of these methods make the system to be a new perspective in clinical studies and also in the basic research areas.

## MATERIALS AND METHODS

We analyzed the effect of three standard antineoplastic drugs: cisplatin, carboplatin and oxaliplatin on primary cell lines obtained from the biopsies of cancer patients. These drugs are used in the therapeutic scheme of several localizations, among which we select ovary and digestive tumors.

Also we synthesised a new group of platinum and palladium complexes with general formula  $[MX_2L_2]$  where  $M = Pt, Pd$ ;  $X = I, Cl$ ;  $L =$  diphenylarsine ligand, in order to determine their potential biological activity. These complexes have organometallic ligands with structures which lead us to presume that they exhibit biological effects: 2-PrOC<sub>6</sub>H<sub>4</sub>AsPh<sub>2</sub> (**L1**), 2-MeOC<sub>6</sub>H<sub>4</sub>AsPh<sub>2</sub> (**L7**) and 2-HOC<sub>6</sub>H<sub>4</sub>AsPh<sub>2</sub> (**L11**) [Fig.1 and 2]. Each structure contain two identical tertiary arsine groups bonded to platin, respectively to palladium, the only difference between these groups being one moiety, which is isopropyl for compound L1, methyl for L7 and hydroxyl for compound L11, molecular structures of the six complexes are analogous.




Fig.1. Structure of ligands L1 and L7



Fig.2. Structure of ligand L11

We performed cell cultures using MLS tumoral ovary cell line and Hfl-1 normal human lung fibroblast cell line. Primary cell cultures were performed using ovary tumor biopsies obtained with the patient informed consent. To obtain the primary cell suspensions, we performed mechanic-enzymatic disaggregating procedures using the Medimachine equipment. Cell cultures were performed in class II laminar hoods, cells were kept in sterile conditions in incubators with stable temperature (37°C), humidity and CO<sub>2</sub> concentration (5%). For the experiments, tumoral and normal cells were placed on 6-well and 96-well plates and they were treated with the novel synthesized compounds, and with the metal-based drugs, respectively. Measurements were made after 24 hours of incubation with the platinum and palladium complexes.

Cell line Hfl-1 was acquired by Sigma Aldrich Corporation from European Collection of Cell Cultures; MLS ovary cell line was a donation of Dr. Yael S. Schiffenbauer from MedisEI Company, Lod, Israel.

Cell cultures were performed using Dulbecco's Modified Eagles Medium, RPMI 1640 respectively F-12 cell growth media from Sigma Aldrich Corporation. Phosphate Buffered Salt Solution, Trypsine EDTA, MTT salt, Ethidium bromide, Low Melting Point Agarose and Type I Agarose, FDA and Rh 123 fluorescent dyes were provided also by Sigma Aldrich. NaCl p.a., NaOH p.a., EDTA-2 HCl p.a. was obtained from Merck&Co. Inc., and Trizma HCl from Fluka Analytics. Supplier of all disposables used for the biological experiments was Nalgene Nunc International. Chemotherapeutic drugs carboplatin (Carboplatin) and oxaliplatin (Eloxantine) were procured from Sindan Pharma.

### Instrumentation:

For this purpose we used the CellScan system, which permit detection of fluorescence intensity (FI) and polarization (FP) of different cell types. There is a strong correlation between inhibition of cell growth induced by different chemotherapeutic agents and the resultant changes in FI and FP. Thus, changes can be revealed by staining treated cells with fluorescent dyes such Fluorescein diacetate (FDA) and Rhodamine 123 (Rh123). Hyperpolarization of the fluorescence intensity detected by the CellScan system can identify apoptotic cells within a complex cell population; it is a marker of early apoptotic processes.

The CellScan apparatus is a laser scanning, static, multiparameter cytometer (Harel, 2005), measuring optical parameters: fluorescence intensity (FI) and polarization fluorescence (FP) of individual, fluorescent-labelled cells. By using this system in cancer research, many diagnosis and chemosensitivity tests became available. The simplicity of this method makes the system to be a new perspective in clinical studies and also in the basic research area.




**Fig. 3.** The cell carrier: A. Tumor cell trapped in the hole of the cell carrier. B. Arrangement of apertures on the cell carrier surface. C. The cell carrier is placed on a plastic grid. D. The carrier can be loaded using pressure gradient. Image kindly provided by Medis El Technologies.

The CellScan system enables repetitive spectroscopic measurements in intact living cells within a population under physiological conditions. It incorporates a unique "cell carrier" [Fig.3] which allows repeated high-precision measurements by fixing the location of the cell in the cell carrier. The "core" of the carrier is a 2 mm squared grid of precisely dimensioned conic apertures : 5-8 µm in diameter, spaced at 20 µm, which can hold 10000 cells, each loaded and maintained in a separate trap. Using the cell carrier, the system

can define and retain the locations of individual cells and the resulting measurements (Deutsch, 1994)

The most widespread equipments which quantify optical properties of cells are spectrophotometers, flow cytometers, and confocal laser scanning microscopes. It is possible to make an analogy with the well-known flow-cytometer; both are able to study the functions of the fluorochrome-labelled cells. The differences are that the CellScan have a static configuration, provide the results for each individual cell, not only for the whole population, fewer cells are enough to make a determination, and the measurement is repeatable in the same cells (Deutsch, 1996; Rahmani, 1996; Zurgil, 2002).

This device includes an advanced optical system. The individual trapped cells are illuminated by a He-Cd laser beam source which scans each hole and displays the spread of intensity (FI) on the x-axis and polarization (FP) on the y-axis on the screen. The measurements are made using a preset photon counting technique, in which the same number of photons is counted for each cell, regardless emission's intensity; so that the measurement has the same precision both for the weakly or strongly emitting cells.

## RESULTS AND DISCUSSIONS

We performed static cytometry measurements on primary tumor cultures treated with platin-based drugs, and both on secondary tumor and normal cell cultures, treated with the novel synthesized compounds, but also with the chemotherapeutic drugs, to compare their antiproliferative effect with the standard drugs.

The fluorescence polarization (FP) indicates the entropy, the relative rotational freedom of the fluorescent molecules. The FP of a fluorochrome increases with viscosity or with the restrictions of molecular rotation. Cell activation and apoptosis involve conformational changes of cytosolic enzymes and proteins, the intracellular matrix reorganization. The more the molecule rotational movement is restricted, the higher is the FP value, and vice versa. The fluorescence intensity (FI) of the fluorochrome reflects intracellular content.

The dyes used are Fluorescein diacetate (FDA) and Rhodamine 123 (Rh123). Fluorescein diacetate (FDA) is a non-fluorescent dye, which is converted through enzymatic hydrolysis in its fluorescent form only in the living cells. It is an indicator of cell viability. Rh 123 is a lipophilic cationic fluorochrome that selectively accumulates in the mitochondria of the living cells and whose specific uptake or release depends upon mitochondrial membrane potential, thus reflecting mitochondrial activity (Zurgil, 1996). A drop in the FI of Rh123 was attributed to early apoptosis (Kaplan, 1997; Fixler, 1997).

The CellScan system provides us histograms [Fig.4] representing the polarization and the intensity of every single tested cell from the cell population of the culture.




**Fig. 4.** Histogram provided by the CellScan: the FP and FI for each cell, measured at two wavelengths it is represented by red and green dots. Here are showed untreated (control) cells.




**Fig. 5.** Histogram provided by the CellScan: the polarization and the intensity for each cell, measured at two wavelengths are represented by red and green dots. In figure .a untreated (control) MLS cells are shown. The figure b and c represents FI and FP of MLS cells treated with complex PTL1, respectively PTL11.

In clinical oncology drugs are used in concordance with the international chemotherapy protocols for a certain type of tumor (Schiffenbauer, 2002). A recent study showed that FDA fluorescence hyperpolarization is a measure of apoptosis. Alterations of the mitochondrial structure and function are the early events in apoptosis in different cell types (Green, 1998; Lizard, 1995; Zurgil, 2000). These alterations were characterized by distinct changes in the mitochondrial membrane, associated with the dissipation of the membrane potential.

The Cell Scan can detect changes induced by anti-neoplastic drugs in the fluorescence intensity and polarization of fluorescent-labelled tumour cells in order to investigate the effect of anti-neoplastic drugs on cancer cells. T47D and T80 human breast cancer cell lines were exposed to Navelbine and to 5-Fluorouracil and the fluorescence properties of the treated cells, stained with FDA and Rh123, were measured by the CellScan.

A strong correlation was found between the inhibition of cell growth induced by the two drugs and the hyperpolarization and the changes in fluorescence intensity. The T80 human breast cancer cells showed after *in vitro* treatment a significant reduction in FI, directly proportional with the exposure and concentration of anti-neoplastic drug, while T47D cells, known as 5Fu-resistant cells, does not show a significant drop in the FI. Changes in FP are also relevant and well correlated with the drug efficacy. An increase in average FP of treated T80 cells (stained either with Rh123 and FDA) versus untreated control was observed simultaneously with the FI decrease when the cells are responding to the drugs.

Another study pointed out the novelty and the advantages of the method which may be used to assist in the prediction of the response to anti-cancer agents (Chaitchik, 2005). This procedure requires only a 48 hours tumoral primary cell culture, and it is independent of proliferation and colony-formation assays, long and difficult. This test has a significantly lower duration as the previous chemosensitivity tests and enables more samples to be tested in a

shorter period of time. The test requires low number of cells, which happens frequent in the case of tumour biopsies.

Using this method, it is possible differentiation between necrotic and apoptotic cells. The CellScan test can detect in this manner the pre-lytic changes induced by the drugs and can measure the sensitivity of cancer cells to them.

In our laboratory we performed studies on tumors prevealed from cancer patients (Soritau, 2005) demonstrating that the CellScan technique is adequate for chemosensitivity tests. This gives a possibility to eliminate ineffective drugs of chemotherapeutic protocols and thus to improve the treatment efficacy. Eighty-eight tumors were tested for *in vitro* response to drugs, using the fluorescence intensity and polarization measurements. Tumors were mechanically dissociated, exposed to different drugs for 24 hours, and stained with two different markers of apoptosis: FDA and Rh123. The FI decrease observed simultaneously with FP increase of the stained cells indicates an inhibition in cell proliferation. The patients were treated with standard protocol regimens. For the patients with advanced cancers (stage III and IV), a correlation between measurements using FDA fluorochrome and the clinical result of the treatment was found.




**Fig. 6.** IC50 values for the MLS tumor cells treated with platinum and palladium complexes

For the novel synthesized compounds we establish the IC50 value, the concentration which inhibits cell proliferation with 50% [Fig. 6], and the correspondent polarization values using the static cytometry. In compare with the untreated, control cells, hyperpolarization of PtL1 treated cells was 5,1%, for the PtL7 treated cells 2,5% and for the PtL11 3,4% (Fischer, 2008). Compounds PtL7 and PdL7 showed a very good potential to

induce early apoptosis, their polarization values obtained for the Rh123 dye was 7,8 [Table1], and 8,3%, respectively. These two compounds, containing the L7 ligand present the best potential to induce programmed cell death.

**Table 1.** Polarization values for tumoral MLS and normal Hfl-1 cells treated with novel synthesized compounds.

| Stimulant   | FP values [a.u.] |             |
|-------------|------------------|-------------|
|             | MLS cells        | Hfl-1 cells |
| Control     | 0.362            | 0,293       |
| Oxaliplatin | 0.358            | 0,310       |
| Carboplatin | 0.398            | 0,345       |
| PtL 1       | 0.371            | 0,317       |
| PtL7        | 0.391            | 0,305       |
| PtL11       | 0.352            | 0,322       |

2000; Collins,2004). Data are correlated with the values obtained using the alternative methods.

To verify the accuracy of the new method, we correlated the results obtained in this way with those determined with customary procedures such as the MTT (MTT = 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide) viability test (Mosmann, 1983) and one of the standard methods for assessing DNA damage, namely, single-cell gel ("comet") assay (Tice,

## CONCLUSIONS

The Cell Scan technology is a valuable tool in cytometry assays, superior in many aspects to the precedent scanners and comparable in efficiency to the well-known flow cytometers. It enables repetitive measurements on the same cells and makes possible the follow-up of a cell population in time. The cell carrier gives the CellScan the capability to investigate the same individual cells after one or several desired manipulations, facilitating the study of cell dynamics and simplifies repetitive examination, even when a small number of cells are available.

The significance of the test lies in its technical simplicity, rapidity, accuracy and the possibility to repeat the measurements in the same cells. The SCM test's aim is to distinguish between patients with neoplastic diseases and healthy controls. Moreover, it can suggest the localization of the cancer completing in this way the other investigations and can precede many clinical and imagistic procedures. The SCM test is a potential tool for screening and early detection of cancer, which in turn may lead to a significant reduction in mortality rates.



The specificity and the sensitivity of the tests made by CellScan were greater than those performed using the spectrofluorimeters. The kinetic information we can obtain is useful in understanding the process of lymphocyte activation and has clinical relevance as well. The results showed the possibility of a different, high specificity method for detection of cancer by a blood test.

The six novel synthesized platinum and palladium compounds are biologically active against tumoral cells; their cytotoxicity is comparable with standard drugs. Measurements using the CellScan technology are well correlated with the results provided by other bioassay methods. The studied compounds, especially PtL7 and PdL7 are suitable to be tested as new drugs.

It was demonstrated the potential of the CellScan system to detect the in vitro effects of the chemotherapy performed on cancer cells. The system can be used to assess drug sensitivity of cancer cells derived from solid human tumours, biopsies in order to predict the in vivo effects of the treatment. The test has the ability to differentiate between drug sensitivity and resistance in cancer cells.

**Acknowledgements:** The present study was founded by CNCSIS Complex Exploratory Research Project "Metalloomics". We appreciate the support of Cell Kinetics Department from MedisEl Company, Lod, Israel.

## REFERENCES

1. Akamatsu K., H. Saito, T. Tsunenari, T. Matsumoto, K. Morikawa, M. Koizumi, H. Mitsui and K. Koizumi - *Anticancer Res.*, 1993, 13, 2261-2265.
2. Barbara C., P. Orlandi, G. Bocci, A. Fioravanti, A. Di Paolo, G. Natale, M. Del Tacca and R. Danesi - *Eur. J. Pharmacol.*, 2006, 549, 27-34.
3. Begun N. : *Am. J. Physiol.*, 1994, 267, E14-23.
4. Berger B. J. , A. H. Fairlamb - *Antimicrob. Agents Chemother.*, 1994, 38, 1298-302.
5. Carland M., K. J. Tan, J. M. White, J. Stephenson, V. Murray, W. A. Denny, W. D. Fayden - *J. Inorg. Biochem.*, 2005, 99, 1738-1743.
6. Cemazar M., Z. Pipan, S. Grabner, N. Bukovec, G. Sersa - *Anticancer Res.*, 2006, 26, 1997-2002.
7. Chaitchik S., O. Soritau, E. Fischer, D. Pelau, L. Lazar, Y.S. Schiffenbauer - *J. Clin. Oncol.*, 2005, 23(165): 2125.
8. Collins A. R. - *Mol. Biotechnol.*, 2004, 26, 249-261.
9. Deutsch M., A. Weinreb, *Cytometry*, 1994, 16: 214-226.
10. Deutsch M., I. Ron, A. Weinreb, R. Tirosh, S. Chaitchik, 1996, 23: 159-165.
11. Fairlamb A. H., G. B. Henderson and A. Cerami, *Proc. Natl. Acad. Sci. U S A.*, 1989, 86: 2607.
12. Fischer -Fodor E. , N. Moldovan, P. Virag, O. Soritau, I. Brie, P. Lönnecke, E. Hey-Hawkins, L. Silaghi Dumitrescu. : *Dalton Trans.*, 2008 : 6393-6400.
13. Fixler D, Tirosh R, Eisenthal A, Marder O, Irlin Y, Lalchuk S, Deutsch M. *Biol Cell*, 1997, 89: 443-452.
14. Frost S. C, M. S. Schwalbe, *Biochem. J.*, 1990, 269, 598-595.
15. Fukuda M., Y. Ohe, F. Kzawa, M. Oka, K. Hara and N. Saijo, *Anticancer Res.*, 1995, 15, 393-398.
16. Gay M. , A. M. Montaña, V. Moreno, M. J. Prieto, J. M. Pérez, C. Alonso, *Bioorg. Med. Chem.*, 2006, 14, 1565-72; M. Gay, A. M. Montaña, V. Moreno, M. J. Prieto, R. Llorens, L. Ferrer, *J. Inorg. Biochem.*, 2005, 99, 2387-2394.
17. Geran R. I., N. H. Greenberg, M. M. Macdonald, A. M. Schumacher, B. J. Abbott, *Cancer Chemother. Rep.*, 1972, 3, 1-85.
18. Green D.R, Reed J.C, Mitochondria and apoptosis. *Science* 1998, 281(5381):1309-1312.
19. Harel M., B. Gilburd, Y. S. Schiffenbauer, Y. Schoenfeld. *Clin. Dev. Immunol.*, 2005, 12(3): 187-195.
20. Kaplan MR, Trubniykov E, Berke G. *J. Immunol. Meth.*, 1997, 201: 15-24.
21. Kwong F. Y., C. W. Lai, M. Yu, D.-M. Tan, F. L. Lam, A. S. C. Chan, K. S. Chan, *Organometallics*, 2005, 24, 4170-4178.
22. Lizard G., S. Fournel., L. Genestier., N. Dgedin, C. Chaput, M. Flacher. *Cytometry*, 1995, 21: 275-283.
23. Moldovan N., P. Lönnecke, I. Silaghi-Dumitrescu, L. Silaghi-Dumitrescu, E. Hey-Hawkins, *Inorg. Chem.*, 2008, 47, 1525-31.
24. Mosmann T., *J. Immunol. Meth.*, 1983, 65, 55-63.
25. Mottram J. C. , G. H. Coombs, *Exp. Parasitol.*, 1985, 59(2):151-60.
26. Phadnis P. P., V. K. Jain, A. Klein, M. Weber, W. Kaim, *Inorganica Chim. Acta* 2003, 346, 119-128; P. P. Phadnis, V. K. Jain, T. Schurr, A. Klein, F. Lissner, T. Schleid and W. Kaim, *Inorganica Chim. Acta.*, 2005, 358, 2609-2617.
27. Rahmani H., M. Deutsch, I. Ron, S. Gerbat, R. Tirosh, A. Weinreb, S. Chaitchik, S. Lalchuk, *Eur. J. Cancer*, 1996, 32A (10): 1758-1765.
28. Robillard M. S., M. Bacac, H. van den Elst, A. Flamigni, G. A. van der Marel, J. H. van Boom, and J. Reedijk, *J. Comb. Chem.*, 2003, 5, 821-825.
29. Schmidt A C., M. Neustadt, M. Otto, *J. Mass Spectrom.*, 2007, 42:771-80.
30. Schiffenbauer Y.S., E. Trubniykov, Ben-Tzion Zacharia, S. Gerbat, Zvi Rehavi, G. Berke, S. Chaitchik. *Anticancer Res.*, 2002, 22: 2663-2670.
31. Soritau O., Fischer E, Tanasescu R., Muresan M.A., Pelau D., Lazar L, Chaitchik S, Schiffenbauer Y.S, 16th International Congress on Anti-Cancer Treatment, 2005, Abstract Book: 346.
32. Takahashi H., K. Suzuki and H. Namiki, *Cell. Struct. Funct.*, 2003, 28, 123-130.
33. Tice R. R., E. Agurell, D. Anderson, B. Burlinson, A. Hartmann, H. Kobayashi, Y. Miyamae, E. Rojas, J. C. Ryu and Y.F. Sasaki, *Envir. Molec. Mutagen.*, 2000, 35: 206-221.
34. DeVita V. T., S. Hellman, S. Rosenberg in: *Cancer: Principles and Practice of Oncology*, ed Lippincott, Williams and Wilkins, Philadelphia, 7th ed, 2004.ch 15, sect 5.
35. Verspohl E. J. - *Endocrin. J.*, 2006, 53, 21-26.
36. Wang Z., H. Zhang, X. F. Li and X. C. Le - *Rapid. Commun. Mass Spectrom.*, 2007, 21:3658-3666.
37. Zurgil N, Shafran Y, Fixler D, Deutsch M. - *Biochem Biophys Res Commun*, 2002, 290: 1573-1582.
38. Zurgil N, Z. Schiffer, Y. Shafran, M. Kaufman, M. Deutsch. - *Biochem Biophys Res Comm* 268: 155-163, 2000.
39. Zurgil N, Deutsch M, Tirosh R, Brodie C. - *Cell Struct Funct*, 1996, 21: 271-276.

# FOOD SUPPLEMENTS WITH CHROMIUM AND SELENIUM CONTENT

Gabriela GARBAN\*

*Environmental and Food Hygiene, Institute of Public Health Timișoara, Blvd. Dr. V. Babes Nr. 16-18, RO – 300 226 Timisoara, Roumania*

\* E-mail : garban\_g@yahoo.com

## ABSTRACT

*Food supplements, according to Directive 2002/46/EC means foodstuffs the purpose of which is to supplement the normal diet and which are concentrated sources of nutrients or other substances with a nutritional or physiological effect, alone or in combination, marketed in dose form, namely forms such as capsules, pastilles, tablets, pills and other similar forms, sachets of powder, ampoules of liquids, drop dispensing bottles, and other similar forms of liquids and powders designed to be taken in measured small unit quantities; "nutrients" means the following substances: (i) vitamins, (ii) minerals. This Directive was transposed also by the Romanian sanitary authorities and decided to start monitoring and inspecting food supplements with vitamins and minerals at national level. In this context, the specialists from the Institute of Public Health Timișoara elaborated a methodology which was transmitted to each County Directorate of Public Health and of Bucharestts. The aim of this work was to identify at national level the number of manufacturers, importers, wholesalers and retailer units of food supplements. In the present paper there will be discussed only the food supplements with chromium and selenium content.*

**Key words:** chromium, selenium, food supplements

## INTRODUCTION

Mineral insufficiency and trace element insufficiency states are actually more likely to occur than are vitamin insufficiency. Those at increased risk of such insufficiencies include people who eat low-calories diet, the elderly, pregnant women, people taking certain drugs (e.g. diuretics), vegetarians and those living where the soil is deficient in certain minerals. Because of differing geologic conditions minerals and trace elements may scarce in the soil of certain region and rich in those of other regions (DiSilvestro, 2001; O'Dell and Sunde, 1997; Garban, 2004).

Any sort of stress deplete the vitamin and mineral stores of organism more rapidly. With age the organs tend to function less efficiently: the metabolisms slows, physical activity diminishes, food intake goes down.

Various diseases reduce also the mineral nutrients intake and especially those that interfere with the ingestion, digestion, absorption and requirement of nutrients: celiac disease, Crohn disease, irritable bowel syndrome, lactose intolerance, bacterial, viral and parasitic infections.

Evidence from recent studies reveal that mineral trace element supplementation may help to prevent various forms of cancer, heart disease and some other degenerative processes.

In this context the importance of food supplements intake with mineral content is discussed.

## MATERIALS AND METHODS

At national level the Ministry of Health of Roumania adopted a law concerning food supplements which was based on the EU directive. After this directive the health authorities decided to start in the framework of the National Health Programme 1.4 the monitoring and inspecting food supplements at national level. In this context, a methodology was elaborated by the specialists from the Institute of Public Health Timișoara and widespread to 41 County Directorate of Public Health and to the Directorate of Public Health of Bucharest. The task of specialists was to found out the number of manufacturing units, importing units, wholesalers and retailers of food supplements. Thenafter, to identify the composition of food supplements based on vitamins and minerals (Garban, 2006; Garban, 2008).

According to the methodology the specialists sent the obtained data to the Institute of Public Health where they were processed.

## RESULTS AND DISCUSSIONS

Data were received from 36 County Directorate of Public Health and the Directorate of Public Health of Bucharest. The total number of identified food supplements was 916 but only 121 were in accordance with the national law (Ordin MSP 1069/2007) referring to food supplements which transpose EU Directive (Garban, 2008).

According to the Directive 46/2002/EC only a limited number of vitamins (13) and minerals (15) are accepted as ingredients in food supplements –Table 2.

**Table 2.** Vitamins and minerals allowed by law in food supplements

| <b>Vitamins</b>       | <b>Minerals</b> |
|-----------------------|-----------------|
| Vitamin A (µg RE)     | Calcium (mg)    |
| Vitamin D (µg)        | Magnesium (mg)  |
| Vitamin E (mg α-TE)   | Iron (mg)       |
| Vitamin K (µg)        | Copper (µg)     |
| Vitamin B1 (mg)       | Iodine (µg)     |
| Vitamin B2 (mg)       | Zinc (mg)       |
| Niacin (mg NE)        | Manganese (mg)  |
| Pantothenic acid (mg) | Sodium (mg)     |
| Vitamin B6 (mg)       | Potassium (mg)  |
| Folic acid (µg)       | Selenium (µg)   |
| Vitamin B12 (µg)      | Chromium (µg)   |
| Biotin (µg)           | Molybdenum (µg) |
| Vitamin C (mg)        | Fluoride (mg)   |
|                       | Chloride (mg)   |
|                       | Phosphorus (mg) |



Minerals are essential nutrients to life and are needed in small amounts and beside vitamins they constitute the class of micronutrients.

These nutrients participate in a multitude of biochemical and physiological processes necessary for the maintenance of health. For the human organism there are essential about twenty eight elements, e.g. Ca, Mg, Na, K, Zn, Fe, Cu etc.

As to minerals in food supplements one have to mention that there are a lot of data concerning the products with calcium, magnesium, iron, copper, iodine, zinc, manganese etc. content. But, food supplements with chromium, selenium, molybdenum are less known.

Among the identified food supplement there were found 75 containing chromium, without specification of their chemical form – details in Table 2.

**Table 2.** Food supplements with Cr and Se content on the Romanian market

| <b>Specification</b>           | <b>Single component</b> | <b>With other minerals and vitamins</b> | <b>With minerals vitamins and other substances</b> | <b>Total</b> |
|--------------------------------|-------------------------|-----------------------------------------|----------------------------------------------------|--------------|
| Cr containing food supplements | 4                       | 23                                      | 48                                                 | 75           |
| Se containing food supplements | 1                       | 30                                      | 72                                                 | 103          |

Chromium is an essential mineral found in very low concentrations in the human body. This mineral is a component of the compound, "glucose tolerance factor" (GTF) in humans. GTF is critical for the maintenance of proper serum levels of blood glucose, or insulin metabolism, and its many diverse actions. GTF plays integral roles in the metabolism of other nutrients as well. Fatty-acids and carbohydrates are metabolized by this compound. GTF is also necessary for the activation of enzymes that aid in the digestion of various food sources, and in cholesterol synthesis, which results from daily food consumption (Scwartz and Mertz, 1959; Mertz, 1993).

Chromium is found in the highest concentrations in those organs that provide for metabolic activity. These include the liver, kidney, spleen and even bone tissue(s). Many medical practitioners label chromium as an "ultra" trace mineral because it is found in such diminutive concentrations throughout the body. The amount of chromium stored in the body is in direct correlation to one's diet. It is estimated that Americans store approximately 6 milligrams of chromium in various metabolic tissues.

Chromium is commercially available in several forms including chromium nicotinate, chromium picolinate, chromium-enriched yeast, and chromium chloride. Chromium is available as part of many multivitamins or alone in tablet and capsule forms. Studies suggest that chromium supplementation may be helpful for the following conditions: diabetes , high cholesterol, obesity, strength training, high blood pressure (Lukaski, 1999; Vincent, 2003).

As to food supplements with selenium content a number of 103 products were identified. Most of them are imported from USA and other EU Member States.

Selenium is a trace mineral that is essential to health maintenance but required only in small amounts (Thomson, 2004). Selenium is incorporated into proteins to make selenoproteins, which are important antioxidant enzymes. The antioxidant properties of selenoproteins help prevent cellular damage from free

radicals. Free radicals are natural by-products of oxygen metabolism that may contribute to the development of chronic diseases such as cancer and heart disease. Other selenoproteins help regulate thyroid function and play a role in the immune system (Arthur, 1991; Combs and Grey, 1998; McKenzie, 1998; Goldhaber, 2003).

Selenium content of foods can vary according with the soil concentration of selenium. There is evidence that selenium deficiency may contribute to development of a form of heart disease, hypothyroidism, and a weakened immune system. There is also evidence that selenium deficiency does not usually cause illness by itself. Rather, it can make the body more susceptible to illnesses caused by other nutritional, biochemical or infectious stresses.

Three specific diseases have been associated with selenium deficiency: Keshan disease, which results in an enlarged heart and poor heart function, occurs in selenium deficient children; Kashin-Beck disease, which results in osteoarthropathy; Myxedematous endemic cretinism, which results in mental retardation

Studies indicate that death from cancer, including lung, colorectal, and prostate cancers, is lower among people with higher blood levels or intake of selenium. Some population surveys have suggested an association between lower antioxidant intake and a greater incidence of heart disease. Selenium is one of a group of antioxidants that may help limit the oxidation of LDL cholesterol and thereby help to prevent coronary artery disease.

Recently were established the new values of the Recommended Daily Allowances for vitamins and minerals (Table 3).

**Table 3.** Recommended Daily Allowances (RDAs) for minerals  
(established in October 2008)

| <b>Minerals</b> | <b>RDA</b> |
|-----------------|------------|
| Calcium (mg)    | 800        |
| Magnesium (mg)  | 375        |
| Iron (mg)       | 14         |
| Copper (µg)     | 1          |
| Iodine (µg)     | 150        |
| Zinc (mg)       | 10         |
| Manganese (mg)  | 2          |
| Sodium (mg)     | -          |
| Potassium (mg)  | 2000       |
| Selenium (µg)   | 55         |
| Chromium (µg)   | 40         |
| Molybdenum (µg) | 50         |
| Fluoride (mg)   | 3.5        |
| Chloride (mg)   | 800        |
| Phosphorus (mg) | 700        |

Today, minerals are used to correct subtle nutritional deficiencies in relatively healthy people; to compensate for increased mineral needs caused by health problems; to enhance exercise training etc.

## CONCLUSIONS

1. Food supplements marked at national level are mostly with complex composition, containing beside vitamins and minerals herbal extracts, proteins, probiotics etc.
2. Among the 916 identified food supplements a great number contained chromium (75 products) and selenium (103 products).
3. Most of food supplements had complex composition, meaning that in their composition contains not only vitamins and minerals but also other substances.

**Acknowledgements.** The author want to express her gratitude to all specialists from the Counties Directorate of Public Health from Roumania who participated and send data for the National Synthesis - Monitoring and Inspecting Food Supplements in Romania 2007 in the framework of the National Health Programme 1.4. This work was financially supported by the Roumanian Ministry of Public Health.

## REFERENCES

1. Arthur J.R. - The role of selenium in thyroid hormone metabolism. *Can. J. Physiol. Pharmacol.*, 1991, 69, 1648-52.
2. Berdanier C.D. - *Advanced Nutrition: Micronutrients*. CRC Press, Boca Raton, 1998.
3. Combs G.F.Jr., Gray W.P. - Chemopreventive agents: Selenium, *Pharmacol. Ther.*, 1998, 79, 179-192.
4. Combs G.F., Clark L.C., Turnbull B.W. - An analysis of cancer prevention by selenium, *BioFactors*, 2001, 14, 153-159.
5. Ellis D.R., Salt D.E. - Plants, selenium and human health. *Curr. Opin. Plant Biol.*, 2003, 6, 273-279.
6. Fleet J. C. - Dietary selenium repletion may reduce cancer incidence in people at high risk who live in areas with low soil selenium. *Nutr. Rev.* 1997, 55, 277-279.
7. Garban Gabriela – Monitorizarea și inspecția suplimentelor alimentare, Sinteză națională, Ed. Eurobit Timișoara, 2008.
8. Garban Z. – *Biochemistry: Comprehensive treatise*, Vol.I, 4<sup>th</sup> edition, Ed. Solness, Timisoara, 2008.
9. Garban Z., Garban Gabriela – *Nutriția umană*, ediția 3, Ed. Orizonturi Universitare Timișoara, 2004.
10. Levander O.A., Beck M.A. - Interacting nutritional and infectious etiologies of Keshan disease. Insights from coxsackie virus B-induced myocarditis in mice deficient in selenium or vitamin E. *Biol Trace Elem Res* 1997, 56, 5-21.
11. Lukaski H.C. - Chromium as a supplement. *Annu. Rev. Nutr.*, 1999, 19, 279-302
12. McKenzie R.C., Rafferty TS, Beckett GJ. Selenium: an essential element for immune function. *Immunol Today*, 1998, 9, 2-5.
13. Mertz W. - Chromium in human nutrition: a review. *J. Nutr.*, 1993, 123, 626-33.
14. O'Dell B.L., Sunde R.A. (Eds.) - *Handbook of nutritionally essential mineral elements*. New York, Marcel Dekker Inc., 1997.
15. Schwarz K, Mertz W. Chromium(III) and the glucose tolerance factor. *Arch. Biochem. Biophys.*, 1959, 85, 292-295.
16. Russo M.W., Murray S.C., Wurzelmann J., Woosley J.T., Sandler R.S. - Plasma selenium levels and the risk of colorectal adenomas, *Nutr. Cancer*, 1997, 28, 125-9.

17. Thomson C.D. - Assessment of requirements for selenium and adequacy of selenium status: a review, Eur. J. Clin. Nutr., 2004, 58, 91-402.
18. Vincent J.B. - The potential value and toxicity of chromium picolinate as a nutritional supplement, weight loss agent and muscle development agent. Sports Med., 2003, 33, 213-30.
19. \*\*\* - Directive 2002/46/EC
20. \*\*\* - Ordin MSP 1069/2007

# **ADDUCTS OF DEOXYRIBONUCLEIC ACID WITH METALS „AD LIMINA” OF INTERDISCIPLINARITY METALLOMICS AND PROTEOMICS**

**Zeno GARBAN \***

*Department of Biochemistry and Molecular Biology, Faculty of Food Products Technology, University of Agricultural Sciences and Veterinary Medicine of Banat, Calea Aradului nr. 119, RO-300 645, Timișoara, Roumania*

\* E-mail : zgarban@yahoo.com

## **ABSTRACT**

Biogenesis of adducts between deoxyribonucleic acid and micromolecules presents a special interest because these are studied both in biochemistry and xenobiochemistry within a dual framework: biochemistry / patobiochemistry, physiology / physiopathology, morphology / morphopathology. If we circumscribe the problem of adducts between DNA and metal ions ( $M^{n+}$ ) to the domains of major interest for life sciences, we can state that, in fact, there are approached problems of metallomics and proteomics at their very frontier.

In order to have a global image are discussed aspects related to adducts' biogenesis in biochemistry and xenobiochemistry as well as, in a succinct manner, peculiarities of metallomics and proteomics regarded as far-reaching domains that address the relation chemical structure – biological activity, evidently with implications in biochemical homeostasis, heterostasis, homeorhesis as mechanisms of the regulation and control systems in the biologic processes developed in vivo and even in vitro. Some of these aspects present interest for modern biotechnologies, too, often applied on DNA molecules.

**Key words:** DNA adducts, metallomics, proteomics

## **1. BIOGENESIS OF DNA ADDUCTS IN BIOCHEMISTRY AND XENOBIOCHEMISTRY**

The term “DNA adduct” is generally used to describe a covalent binding of a chemical compound to the macromolecule of deoxyribonucleic acid. The chemical compound can be a biological substance (e.g.: biometals) but also a xenobiotic derivative (e.g. food additives with metals, chemotherapeutic drugs etc.) .

Investigations on the compounds resulted from the interaction of the DNA macromolecule with small molecules were made for the first time in the sixth decade of the 20th century, and the results lead to the idea of appearance of some bioincompatible structures that were named for the first time “molecular complexes”.

It was also used the term “molecular associations” (Bergmann, 1968) but both denominations were improper with the characteristics of the studied chemical structures.

After years of physico-chemical investigations and mechanic-quantum calculations on the electronic density in the donor-acceptor relationship between DNA and small molecules, the term “adducts” was preferred (Grunberger and Weinstein, 1979; Stich, 1988). Therefore this denomination will be used also in this paper to describe “molecular complex” resulted from the covalent binding of a chemical substance to the DNA macromolecule.

The biogenesis of DNA adducts is important for the domain of Biochemistry and Xenobiochemistry. Biometallic ions like Mg and Zn have an important role in the stabilization of the DNA macromolecule. The negatively charged phosphate backbone of DNA (with polyanionic characteristics) prevents the formation of the duplex without the positively charged metallic ions that have the ability to neutralize the charge of the phosphate bases. The metallic ions that have this ability are usually bivalent ions and they attach to two consecutive phosphate bases on the same strand. These interactions are specific for the domain of biochemistry.

Foreign chemical substances, also called xenobiotics, or their metabolites, may react with DNA macromolecules, too. The resulting DNA-adducts can disrupt the biologic activity of DNA, leading to the perturbation of replication, transcription and even DNA-repair process. Xenobiotics that can lead to DNA adducts, and therefore have carcinogenic potential, are divided in two categories: a) indirect-acting carcinogens – xenobiotics that require metabolic activation prior to DNA adduct formation and b) direct-acting carcinogens – xenobiotics that can damage DNA directly, without needing a metabolic activation in order to generate adducts. These interactions with the DNA macromolecule make one of the object of xenobiochemistry.

## **2. METALLOMICS AND PROTEOMICS – INTERACTIVE ASPECTS**

### **2.1. Metallomics**

The term metallome has been introduced by R.J.P. Williams (2001) from Oxford University and afterwards by Hiroki Haraguchi (2004) from Nagoya University by analogy with proteome, as distribution of free metal ions in every cellular compartments. It refers to the complete complement of metals and metal moieties in a biological cell, tissue, or system. This definition is similar to that of the genome (gene), proteome (proteins), and metabolome (metabolites).

Metallomics can be defined as “the study of metals and metal species as well as of their interactions, transformations and functions in biological systems” (Koppendaal and Hieftje, 2007).

In a larger context, metallomics is defined as “comprehensive analysis of the entirety of metal and metalloid species within a cell or tissue type” (Szpunar, 2005).

In this acception “metallomics” can be considered as a branch of “metabollomics” although metals are not typically metabolites.

While traditional bioinorganic chemistry has focused on the role and interactions of a single (or few) metals in a protein or enzyme system, metallomics studies global, multi-element interactions and relationships.

Metallomics gained importance as an emerging science because about 30% of proteins contain metals, such as metalloproteins and metalloenzymes. There also

are numerous metabolites originating from carbohydrates and lipids that are bond to metals.

Metallomics is distinct from other notions (such as speciation) because it has a global role, dealing with the investigation of metals and metalloids (non-metallic elements) from nature, predilectly from biological systems.

It is necessary to mention that metallomics is a new domain, which could develop due to the progresses made in the field of atomic absorption spectroscopy

## 2.2. Proteomics

The proteome is the entire complement of proteins expressed by a genome, cell, tissue or organism. More specifically, it is the set expressing proteins at a given time under defined conditions. Taking into account the concept of proteome, the term proteomics can be defined as the study of the set of proteins produced by an organism, tissue or cell, with the aim of understanding the behaviour of these proteins under varying environments and conditions. In proteomics are studied, on a large scale, structural and functional peculiarities of the proteins. Generally, in life sciences, is considered that proteomics is the next step in the investigation of biologic systems, after genomics.

It is estimated that proteomics is more complex as compared to genomics because a proteome differs from cell to cell and changes constantly by the biochemical interactions with the genome and the environment. The protein expression differs in function of life cycle (Belhajjame et al., 2005).

At the beginning of the third millenium the evolution of knowledge about genome led to the intensification of researches of molecular biology and genetics that imposed the new term „*genomics*”.

## 2.3. Ad limina metallomics – proteomics

In the domaine of life sciences the problem of interdisciplinarity at the frontiers (ad limina) of metallomics and proteomics, can be registered and explained numerous aspects with interest for biochemistry, patobiochemistry, citostatic chemotherapy (as elective domaines discussed in this paper).

The biogenesis of compounds that can be studied by metallomics and proteomics is in fact of interest for numerous domaines such as: nutrition, pharmacology, enzymology, biotechnology a.o.

*Genomics* - is a distinct field of life sciences studying the entire genome of an organism. It also includes studies of intragenomic phenomena and other interactions between loci and alleles within the genome.

The genomics is included in the more complex problem of *proteomics*.

Genomics and proteomics are considered fundamentally important scientific fields, because genes (DNAs and RNAs) contain the genetic information codes to synthesize various proteins. However, genes and proteins cannot be synthesized without the assistance of metalloenzymes containing zinc and other metals. The term metallomics therefore was suggested to be considered at the same level of scientific significance as genomics and proteomics (Haraguchi and Matsuura, 2003).

Currently in the domaine of *genomics* one has in view (as in proteomics) the duality structure-function, knowing that: a) *structural genomics* - is the branch of genomics that determines the three-dimensional structures of proteins; b) *functional*

*genomics* - is the branch of genomics that determines the biological function of the genes and their products.

In a larger frame is remarkable the observation regarding *genetics* vs. *genomics*. The two terms are sometimes used interchangeably. However, *genetics* usually refers to the study of single genes, while *genomics* refers to the study of all the genes in a person or organism.

The goal of *genomics* is to promote the understanding of the structure, function, and evolution of genomes in all kingdoms of life and the application of genome sciences and technologies to challenging problems in biology and medicine.

In order to function properly an organism must regulate transcription, translation, assimilation and incorporation of the necessary biometals. Therefore the biochemistry of the cell must be characterised not only by its genome or its proteins content (proteome) but also by the distribution of the metals and metalloids among the different species and cell compartments (metallome). The term that would therefore refer to the totality of individual metal species in a cell including the inorganic (free and complexed) element content and the assembly of metallic complexes with biomolecules (especially with proteins) is metalloproteome (Szpunar, 2004).

Most of the biological substances (nutrients) as well as metal ions are transported inside the cell through the membrane. In general, the biotransformation of these substances is occurring inside the cell and also often in the cell membrane. The processes by which the nutrients are transported in and out of the cell and converted in biological active forms are called metabolism. Biological substances, which are usually small molecules such as amino acids, organic acids and metal ions produced in metabolism, have been called metabolomes or metabolites and are the subject of the field called metabolomics (Lindon et al., 2003).

### **3. ADDUCTS OF DNA MACROMOLECULE WITH METALLIC IONS**

#### **3.1. Characteristics of DNA adducts in biochemistry and molecular biology**

Metal ions ( $M^{n+}$ ) as chemical agents, determine specific interactions based on ionization processes occurring at different levels of the DNA macromolecule followed by the metal ions coordination. In the double-stranded and even simple-stranded structure of DNA appear changes caused by the formation of adducts ( $DNA-M^{n+}$ ).

The action of metals on biologic systems is realised during certain processes involved in chronobiochemical mechanisms. Implicitely, the reaction between DNA and metals is submitted to cronobiochemical mechanisms, fact that can be explained by the modifications of hepatic DNA synthesis conditioned by the factor time (Garban, 1984).

The interaction of DNA with metal ions is studied by various physico-chemical methods including nuclear magnetic resonance, optical rotatory dispersion, spectroscopy (UV, VIS, IR, Raman), circular dichroism.

Bindings of divalent ions depend both on the structure and configuration of the macromolecule, as well as the electronic configuration of  $M^{2+}$ . The distinctive electron provides special peculiarities to the interaction causing changes of the steric parameters followed by biological consequences (fig. 1).






**Fig. 1.** Binding of  $M^{2+}$  to DNA nucleobases: a)  $M^{2+}$ -guanine; b)  $M^{2+}$ -adenine

The electron donor groups of DNA determine the binding at the level of nitrogenous nucleobases. Guanine is often affected as it has the  $N_7$  position, being the center with the highest density of electrons

Alkaline-earth ions bind electrostatically at the phosphodiester groups of DNA. Experimentally have shown that  $Mg^{2+}$  binds more strongly to native DNA than to denaturated DNA (fig. 2)




**Fig. 2.** Binding of  $Mg^{2+}$  to phosphate groups from DNA macromolecule

The electrostatic nature of the binding is explained by the existence of a diffuse electrostatic charge effect. Binding can take place at the phosphate groups from the DNA macromolecule which has the character of macromolecular polyanion.

Transition metals form covalent bindings with the nitrogen atoms present in the nucleobases. The destabilization of the double helix depends on the molar ratio  $M^{2+}/DNA-P$ , the G-C content and the ionic force of the environment.

The effect of biometals studied *in vitro* and *in vivo*, e.g.  $Zn^{2+}$ ,  $Cu^{2+}$  can reveal, at certain concentrations (in excess), modifications of the DNA structure that can precede the biochemical injury (Garban et al., 1994).

A number of studies have shown the destabilizing effect on nucleic acids due to the interaction of divalent cations with  $\pi$ -systems of nucleobases (Garban, 2005).

The location of actions between phosphodiesteric groups is carried out by chelation of the phosphorus group with  $N_7$  of purine nucleobase of GMP.

Intercalation between complementary intrastrand nucleobases shows an intrahelical disposition of  $M^{2+}$ . There are also remarked "sandwich"-type bindings at the level of two nucleobases placed on the same strand (Garban, 2004).

Interaction of metal ions with biologic systems and mostly the interaction of nucleic acids with metal ions (and small molecules) led to numerous studies that followed the mechanism of action and the relation structure-activity (Sigel and Sigel, 1996).

A special mention refers to the fact that there are adducts of  $DNA-M^{n+}$  type used in technical domaine. Such adducts have been obtained with  $Zn^{2+}$  ion. It has to be mentioned that recent studies revealed the semiconductor properties of the  $DNA-M^{2+}$  structures or the possibility to use M-DNA as a biosensor to predict diagnosis (Lee, 1993)

### 3.2. Characteristics of DNA adducts in pathobiochemistry

Generally, the interactions produced between the DNA macromolecule and metal ions can be of interest both for biochemistry and patobiochemistry, being conditioned by the metal ion that is present in the cell and often facilitated by an environment where effects took place on the double helix leading to molecular chain clivage that can be caused by physical factors (e.g. radiations) or even biological ones (e.g. oncoviruses). In such cases, having in view the primar denaturation, the interaction with DNA becomes a biochemical injury (Garban, 2004).

Studies of the relation chemical structure-biological activity revealed also the consequences of interaction and the mechanisms of adduct biogenesis in patobiochemistry. Investigations have been extended over the biomedical, pharmacological aspects, their genotoxicity and also involvement in chemical homeostasis.

The interaction of DNA with  $M^{n+}$  ions without toxic potential (e.g.  $Hg^{2+}$ ,  $Cd^{2+}$ ) produces modifications in structure-activity relationship with undesirable biological effects. Consecutive effects lead to the appearance of chromosomal modifications as well a mutagenic, oncogenic and teratogenic effects (fig. 3).




**Fig. 3.** Internal chelation of DNA nucleobases with metallic ions  
(a)  $\text{Hg}^{2+}$  - A-T nucleobases pair; (b)  $\text{Cd}^{2+}$  - G-C nucleobases pair

Binding of metals with toxicogen potential (e.g.  $\text{Hg}^{2+}$ ) can be made too, at the guanine rests from the nucleosidmonophosphates present in pairs of nucleobases disposed on neighbouring landings in the double helix (fig. 4).




**Fig. 4.** The binding of  $\text{Hg}^{2+}$  ions to guanine adjacent nucleobases

Carcinogenic metals such as Cd, Cr, Ni inhibit DNA repair damaged by direct genotoxic agents.

### 3.3. Characteristics of DNA adducts in chemotherapy

It is known that metals interact with biologic systems representing compounds at the origin of some mutagenic and cancerigenic processes (Beyersmann, 1994). In this situations are the ions  $\text{Hg}^{2+}$ ,  $\text{Zn}^{2+}$  etc.

The interaction between cytostatic chemotherapics and biologic tissues is extremely complex and covers big domains of the chemistry of organometallic compounds (Haiduc and Silvestru, 1989, 1990). There can be mentioned the numerous aspects in connexion with the role of DNA and of the interaction with cytostatic chemotherapics among which cis-platinum plays an esential role (Lippert, 1999; Garban et al., 2000).




**Fig. 5.** Structural formula of cisplatin-dApdG adduct

There are chimiotherapic drugs whose action is based on formation of adduct with DNA macromolecule or with its precursors. There are exemplified cyclophosphamide, cis-platinum.

In case of cisplatin (cDDP) binding, the interaction can take place at one chain or at both chains.

Such an example is given in fig. 5 where the binding of cis-platinum is made at two nucleobases on neighbouring landings. The nucleobases are on the same chain. Figure 6 presents the formation of adducts on one chain - the so called monodentate binding or on both chains – the so called bidentate binding.




**Fig. 6.** Biogenesis of adducts (a) monodentate binding of cDDP to the guanine of DNA ; (b) bidentate binding of cDDP to the guanine of DNA

Interaction of DNA with cis-platinum is also affected by chronobiochemical modifications that take place in the body. It was observed the existence of homeostatic modifications in case of the interaction of cis-platinum at the level of the liver (Garban, 1989). Generally, in chemotherapy, the problem of adducts formation and that of the cytostatic effects are approached in a larger context in correlation with the cell cycle, more exactly in cytokinetics (kinetics of the cell cycle).

## CONCLUSIVE ASPECTS

Adducts' biogenesis presents a particular interest in case of their study in circumscribed domains such as: biochemistry and molecular biology, patobiochemistry, chemotherapy.

The manner in which the thematic approach on adducts focused on metallomics and proteomics has been shortly presented, can offer a more larger framework both to investigations and interpretations that can follow the more thorough defining of the mechanisms of action during the building phase of these compounds.

It is also important to remark that some information obtained in vitro could lead to predictability for in vivo effects. Also can afford a monitorization, by non-invasive means, of certain processes in vivo (e.g. testing of chemotherapeutic drugs, prediction regarding the possibility of appearance of dyshomeostatic effects).

## REFERENCES

1. Beyersmann D.: Interactions in metal carcinogenicity. *Toxicol.Lett.*, 1994, 71 (1-3), pp. 333-338
2. Belhajjame K., Embury S.M., Fan H., Goble C., Hermjakob H., Hubbard S.J., Jones D., Jones P., Martin N., Oliver S., Orengo C., Paton N.W., Poulouvasilis A., Siepen J., Stevens R.D., Taylor C., Vinod N., Zamboulis L., Zhu W. - Proteome Data Integration: Characteristics and Challenges, *Proceedings of the UK e-Science All Hands Meeting*, 2005, 418-425.

3. Bergmann E.D. - The interaction of aromatic hydrocarbons with nucleic acids and their constituents, pp.207- 216, in "Molecular Associations in Biology" (Pullman B., Ed.), Academic Press, New York-London, 1968.
4. Garban Z., Eremia Iulia, Daranyi Gabriela - Chronobiochemical aspects of the hepatic DNA biosynthesis in experimental animals under the action of some metals., J. Embryol. Exp. Morph. Suppl. Cambridge, 1984, 1, 6.
5. Garban Z., Dancău G., Daranyi Gabriela, Erdelean R., Văcărescu G., Precob V., Eremia Iulia, Udriște C. – Implication of chronobiochemistry-metabolism relationship in the induction of homeostasis changes. I. The action of cis-platinum on hepatic DNA biosynthesis and on some serum metabolites in rats, Rev. Roum. Biochim., 1989, 26(2), 107-117
6. Garban Z. - Zinc and copper effects on deoxyribonucleic acid studied in vitro and in vivo, pp.357-362, in "New perspectives in the research of hardly known trace elements" (ed. Pais I.) University Press of U.H.F.S. Budapest, 1994.
7. Garban Z., Carțiș Ioana, Avacovici Adina, Moldovan I. – Comparative aspects between the interaction of deoxyribonucleic acid with some cytostatic drugs: particularisation for the interaction with cis-platinum and cyclophosphamide. 1. Investigations in vivo on experimental animals, pp.1118-1125, in "Mengen- und Spurenelemente, 20. Arbeitstagung 1999", Friedrich Schiller Universität Jena, (Hrsg. Anke M. et al.); Verlag Harald Schubert, Leipzig, 2000.
8. Garban Z. – Influence of metals on DNA, pp. 401-414, in Elements and their compounds in the environment, occurrence, analysis and biological relevance (Eds. Merian E., Anke M., Ihnat M., Stoeppeler M), Vol. I General aspects, 2nd ed., Wiley-VCH Verlag GmbH & Co KGaA, Weinheim, 2004
9. Garban Z. – Xenobiochimie: Tratat comprehensiv, Vol. IV, Editura Eurobit, Timișoara, 2005.
10. Grunberger D., Weinstein I.B. - Biochemical effects of the modification of nucleic acids by certain polycyclic aromatic carcinogens, pp.105-149, in "Progress in nucleic acids research and molecular biology" (Cohn W.E., Editor), Academic Press, New York, 1979.
11. Haiduc I., Silvestru C. - Organometallics in cancer chemotherapy. C.R.C. Press, Boca Raton, Florida, vol.1, 1989; vol.2, 1990
12. Haraguchi H., Matsuura T. - Elemental Speciation for Metallomics, pp. 3-8, in: Proceedings of International Conference on Bio-trace Elements 2002 (BITREL 2002), ed. by S. Enomoto, Published by RIKEN (The Institute of Physics and Chemistry Research), Wako, 2003.
13. Koppenaal D.W., Hieftje G.M. - New Tools & Techniques for the Metallomics Revolution, American Geophysical Union, 2004, abstract B51D-05.
14. Lindon J.C., Holmes E., Nicholson J.K. - So what's the deal with metabolomics?, Anal. Chem., 2003, 75(17), 384A-391A
15. Lippert B.: Cisplatin: chemistry and biochemistry of a leading anticancer drug. Wiley –VCH, Weinheim-New York-Chichester-Brisbane-Singapore-Toronto,1999.
16. Sigel A., Sigel H. (eds.): Metal ions in biological systems, vol. 33, probing of nucleic acids by metal ion complexes of small molecules, Marcel Dekker, New York, 1996
17. Stich H.F., Dunn B.P. - DNA adducts, micronuclei and leukoplakias as intermediate endpoints in intervention trials, pp. 137-145, in: Methods for Detecting DNA Damaging Agents in Humans: Applications in Cancer Epidemiology and Prevention, (Bartsch H., Hemminki K., O'Neill I.K eds), IARC Sci. Publ. No. 89, International Agency for Research on Cancer, Lyon, 1988.
18. Szpunar J. – Metallomics: a new frontier in analytical chemistry, Anal. Bioanal. Chem, 2004, 378, 54-56.
19. Szpunar J. - Advances in analytical methodology for bioinorganic speciation analysis: metallomics, metalloproteomics and heteroatom-tagged proteomics and metabolomics, The Analyst, 2005, 130, 442-465.
20. Williams R.J.P. - Chemical selection of elements by cells, Coordination Chemistry Reviews, 2001, 216/217, 583-595.

# HEPATIC METALLOGRAMS IN RABBITS AFTER SODIUM NITRATE EXCESS IN DRINKING WATER

George-Daniel GHIBU<sup>1\*</sup>, Gabriela GARBAN<sup>2</sup>, Cornel BALTĂ<sup>3</sup>, Marian BURA<sup>3</sup>,  
Cristina PETRUSE<sup>3</sup>, Doina-Elena HORHOI<sup>3</sup>, Zeno GARBAN<sup>4</sup>

\* E-mail : danighibu@yahoo.com

1. Faculty of Chemistry and Environmental Engineering, University Politehnica Timișoara, Piața Victoriei Nr.2, RO-300 006 Roumania; 2. Institute of Public Health, Blvd.Dr.V. Babeș Nr.16, RO-300 226 Timișoara, Roumania; 3. Colegiul Național "Inochentie Micu Klein", str. Slmion Bărnăuțiu nr. 2, RO-515 400 Blaj, Roumania; 4. Faculty of Veterinary Medicine, Faculty of Animal Sciences and Biotechnology and Faculty of Food Processing Technology, University of Agricultural Sciences and Veterinary Medicine of Banta Timișoara, Calea Aradului Nr. 119, RO-300 645 Timișoara, Roumania

## ABSTRACT

Sodium nitrate was administrated in the tap water of rabbits in order to determine the potential effects on hepatic metallograms. The hepatic metallograms were obtained using the atomic absorption spectrophotometry technique and were used to evaluate the effects of nitrates on the biochemical homeostasis, regarding especially the variations of some macro- and trace elements in liver. Sodium nitrate was administered in high concentrations compared with the reference value equal to the maximum contaminant level (MCL) in drinking water set by Environment Protection Agency. The investigation followed the effects of sodium nitrate -  $\text{NaNO}_3$  administered on two groups of rabbits in concentrations of 20 x MCL and 40 x MCL. The analytical determinations underline the effects of nitrates on biochemical homeostasis of macro elements (Na, K, Ca, Mg) and trace elements (Fe, Zn, Cu, Mn, Ni) by determination of their concentrations in the liver of rabbits.

**Key words:** sodium nitrate, liver, metallograms

## INTRODUCTION

Nitrates are natural components of the soil derived from the demineralization of the organic matter which play an essential role in the process of plants development. The extensive agriculture which utilizes at large scale fertilizers based on nitrates and the industrial pollution often leads to the accumulation of high levels nitrates in the environment (Prakasa and Puttanna, 2002). From the environment nitrates accedes in plants and finally in the organism of animals and humans where they exert toxicogenic effects by generating nitrites with methemoglobinizing effects (Speijers et al, 1989). Also if certain conditions are met (i.e.: the presence of nitrites and amines) nitrates consumption may lead to carcinogenic nitrosamines (Vermeer et al., 1998; Rostkowska et al., 1998; Gârban, 2007). In experiments on rats with dietary sodium nitrate administration (using pellets) liver tumors were induced (Aoyagi et al, 1980).

---

„Metal Elements in Environment, Medicine and Biology”, Tome VIII, pp. 163-168, Publishing House “Eurobit” Timișoara, 2008

Proc. of 8<sup>th</sup> Int. Symp. of Roumanian Academy - Branch Cluj-Napoca and Branch Timișoara, 2008, December 5-6, Timișoara, Roumania

Nitrate administration is known to cause disturbances of kidney and liver function and also to alter the biochemical parameters of blood serum (urea, creatinine, uric acid), but the major manifestation of their toxic effects remains the generation of nitrites which react with hemoglobin and cause methemoglobinemia (Zabulyte et al., 2007; Ghibu et al, 2008a). Another aspect which is often neglected in the scientific literature is the fact that nitrates, being salts of nitric acid, also have in their chemical structure metallic ions. These ions are released inside the organism and may interact with other bioconstituents and interfere in the absorption and biodisponibility of other biometallic ions (Ghibu et al, 2008b).

Sodium nitrate is one of the most important compound from the class of nitrates due to its utilization at large scale in both agriculture and food industry. In agriculture sodium nitrate is utilized as fertilizer. Though sodium nitrate is used in agriculture, it is more expensive than ammonium-type fertilizers which are therefore preferred. Ammonium nitrate, ammonium sulfate, urea and anhydrous are the most frequently used fertilizer.

In food industry sodium nitrate is used especially due to its property to generate sodium nitrite which maintains the red color of the meat and meat products and also prevents the growth of certain pathogen bacteria (*Clostridium botulinum*) – Banu et al, 2000.

## MATERIALS AND METHODS

*Experimental model.* Sodium nitrate -  $\text{NaNO}_3$  was added in the tap water of rabbits. The water was administrated “ad libitum” and the effects of nitrates consumption on the mineral metabolism of rabbits were studied. The reference value chosen in the preparation of nitrates solutions was the maximum contaminant level (MCL) admitted in drinking water, a value established by the Environment Protection Agency (EPA) from United States Department of Agriculture (USDA) to 10 mg/L nitrogen nitrate (usually noted as  $\text{N-NO}_3$ ) and 1 mg/L nitrogen nitrite (usually noted  $\text{N-NO}_2$ ).

In this study rabbits (*Oryctolagus cuniculus*) with the age of 30 days and the average weight of  $700 \pm 25$  g were used. The rabbits were included in two experimental groups:  $E_{A(1)}$  and  $E_{A(2)}$  and a control group. All groups were comprised of 10 animals (5 males and 5 females). Animals were fed with VivaBio - a granulated fodder for rabbits produced by Freeman S.R.L (content: 14.94% protein, 2.86% fat and 8.51% cellulose and necessary minerals). Animals from control group received tap water, while animals from  $E_{A(1)}$  group received  $\text{NaNO}_3$  solution with a concentration equivalent with 20 x MCL and  $E_{A(2)}$  group received  $\text{NaNO}_3$  solution with a concentration equivalent with 40 x MCL established for nitrite in drinking water. Nitrates solutions were prepared using tap water from the same source as the control group.

Before the beginning of the experiment, a quarantine period of 10 days was kept during which the health status of rabbits was observed. Also in this period the rabbits were accommodated with the experimental laboratory environment and with the diet made of granulated fodder. After the quarantine, the experiment has started and lasted 20 days.

For the liver sample collection to be made, the subjects were killed according to the procedures recommended by O.G. nr. 37/2002. On this purpose ketamin was administered intravenously. The liver samples were then taken in concordance with the techniques of laboratory animal's necropsy and stored in 25 ml glass bottles



which were placed in a refrigerator until the biochemical investigations were performed.

**Biochemical investigations.** Tissue was prelevated from each animal from the control and experimental groups. The determinations of macro elements (Na, K, Ca, Mg) and trace elements (Fe, Zn, Cu, Mn, Ni) were performed in the “Laboratory of molecular and atomic spectroscopy” of the Faculty of Food Products Technology from Timișoara.

The liver samples were weighted and then calcinated at a 700 °C temperature for a 3 hours time. The obtained ash was mineralized with nitric acid (0,5 N) and brought in gradated flasks of 50 ml. The obtained solutions were analyzed using a spectrophotometer with continuous atomic absorption. The model of the spectrophotometer was Analytik Jena ContrAA 300. Analytical data were expressed in  $\mu\text{g/g}$  wet weight.

**Statistical analysis.** Mean values ( $\bar{X}$ ) and standard deviation (SD) were determined for each parameter obtained. Also the Student test was performed using the software Origin 6.0.

## RESULTS AND DISCUSSIONS

In order to investigate the influence of the concentration of the nitrate solution compared the results from the  $E_{A(1)}$  and  $E_{A(2)}$  groups. The comparison shows the different effects of  $\text{NaNO}_3$  at 20xMCL and 40xMCL concentrations. The results were divided in different tables for the macro elements and for trace elements.

The macro elements investigated in this paper were: Na, K, Ca and Mg. The homeostatic modifications produced by sodium nitrate solutions administrations in different concentrations (20xMCL and 40xMCL) are presented in table 1.

**Table 1.** The quantum of hepatic macro elements in rabbits after administration of  $\text{NaNO}_3$  solutions in different concentrations

| Group             | M.U.            | n  | Na<br>$\bar{X} \pm \text{DS}$ | K<br>$\bar{X} \pm \text{DS}$ | Ca<br>$\bar{X} \pm \text{DS}$ | Mg<br>$\bar{X} \pm \text{DS}$ |
|-------------------|-----------------|----|-------------------------------|------------------------------|-------------------------------|-------------------------------|
| C                 | $\mu\text{g/g}$ | 10 | 887.35<br>$\pm 130.99$        | 3018.18<br>$\pm 122.86$      | 21.14<br>$\pm 1.73$           | 193.36<br>$\pm 12.45$         |
| $E_{A(1)}$        | $\mu\text{g/g}$ | 10 | 913.76<br>$\pm 120.42$        | 2785.46<br>$\pm 116.58^*$    | 22.93<br>$\pm 2.54$           | 174.06<br>$\pm 12,73^*$       |
| $\Delta X_{A(1)}$ |                 |    | +26.41                        | -232,72                      | +1.79                         | -19.30                        |
| $E_{A(2)}$        | $\mu\text{g/g}$ | 10 | 937.72<br>$\pm 58.24$         | 2696.07<br>$\pm 131.66^*$    | 23.74<br>$\pm 2,02^{**}$      | 164.08<br>$\pm 12.26^*$       |
| $\Delta X_{A(2)}$ |                 |    | +50.37                        | -322.11                      | +2.60                         | -29.28                        |

\* $p < 0.01$ ; \*\* $p < 0.05$

The resulted analytical data revealed that sodium nitrate affects the hepatic metalograms. An increase of Na and Ca concentration followed by a decrease of K and Mg concentration shows an alteration of the extracellular ions (Na, Ca) an intracellular ions (K, Mg) whit possible implications in the function of “Na-K pumps” and “Ca-Mg pumps” at the membrane level. These modifications are probably a result of the increased sodium intake.

The liver is an organ with a known role in regulating the water and electrolytes balance in the organism, having the capacity of retaining a part of the water and some ions excess. The retention of sodium in the hepatic tissue after an excess was experimentally demonstrated (Turner and Aizman, 1984).

The sodium excess leads to the perturbation of the physiological equilibrium between extra- and intracellular electrolytes. Hepatic dysfunctions like cirrhosis are known for their ability to affect the metabolism of sodium and other electrolytes and the origin of these effects is the onset of sodium retention (Murakami et al, 1996; Yu et al., 2005).

Significant variations were found in the case of K, Mg and in the case of Ca from  $E_{A(2)}$  group. The results suggested that between the magnitude of the dyshomeostatic effects and the concentration of sodium solution there is a direct proportionality relation.

The homeostatic modifications produced by nitrates on trace elements are presented in table 2 taking as reference the same substance (sodium nitrate) in two different concentrations: 20xMCL -  $E_{A(1)}$  and 40xMCL -  $E_{A(2)}$ .

**Table 2.** The quantum of hepatic trace elements in rabbits after administration of  $\text{NaNO}_3$  solutions in different concentrations

| Group             | M.U.            | n  | Fe<br>$\bar{X} \pm \text{DS}$ | Zn<br>$\bar{X} \pm \text{DS}$ | Cu<br>$\bar{X} \pm \text{DS}$ | Mn<br>$\bar{X} \pm \text{DS}$ | Ni<br>$\bar{X} \pm \text{DS}$ |
|-------------------|-----------------|----|-------------------------------|-------------------------------|-------------------------------|-------------------------------|-------------------------------|
| C                 | $\mu\text{g/g}$ | 10 | 75.57<br>$\pm 9.10$           | 24.60<br>$\pm 4.39$           | 3.77<br>$\pm 0.85$            | 2.04<br>$\pm 0.14$            | 0.19<br>$\pm 0.05$            |
| $E_{A(1)}$        | $\mu\text{g/g}$ | 10 | 67.66<br>$\pm 6.56$           | 23.37<br>$\pm 1.93$           | 3.25<br>$\pm 0.62$            | 1.91<br>$\pm 0.21$            | 0.20<br>$\pm 0.01$            |
| $\Delta X_{A(1)}$ |                 |    | - 5.91                        | - 1.23                        | - 0.52                        | - 0.13                        | + 0.01                        |
| $E_{A(2)}$        | $\mu\text{g/g}$ | 10 | 64.79<br>$\pm 11.40$          | 22.44<br>$\pm 1.46$           | 2.83<br>$\pm 0.88^{**}$       | 1.82<br>$\pm 0.67$            | 0.22<br>$\pm 0.06$            |
| $\Delta X_{A(2)}$ |                 |    | - 8.78                        | - 2.16                        | - 0.94                        | - 0.22                        | + 0.03                        |

$^{**}p < 0.05$

The analytical results show a decrease in concentration in all the analyzed trace elements except for nickel. Significant variations are found only in the case of copper at the  $E_{A(2)}$  group. The hepatic concentrations variations of trace elements are directly proportional with the concentration of the sodium nitrate solution.

From the result shown in table 2 we can see the high concentration of iron which is specific to the liver. The liver is an organ with an important role in iron metabolism. Iron can be stocked in high concentration in liver.

Ferritin is an iron storage protein which can play the role of depositing iron at the hepatic level. After the nitrates solutions were administrated the iron level decreased probably due to the formation nitrites which interact with hemoglobin generating methemoglobin. The decrease of hepatic concentration of iron was experimentally observed after the administration of sodium nitrite to rats (Roth and Smith, 1988).

## CONCLUSIONS

1. Macro elements show in the case of sodium nitrate administration at different concentration an increase of Na and Ca (bioelements with extracellular distribution) and a decrease of K and Mg (bioelements with intracellular distribution). The variation of macro elements concentration in liver is directly proportional with the sodium nitrate solution concentration.
2. The studied trace elements concentration in liver decrease after  $\text{NaNO}_3$  administration, with the exception of Ni which shows an increase in concentration. The trace elements variation is directly proportional with the concentration of  $\text{NaNO}_3$  solutions.

**Acknowledgements:** The authors express their thanks to Assoc. Prof. Ana Driha for the support given during experimental laboratory phases.

## REFERENCES

1. Aoyagi M., Matsukura N., Uchida E., Kawachi T., Sugimura T., Takayama S., Matsui M. – Induction of liver tumors in Wistar rats by sodium nitrite given in pellet diet, *J. Natl. Cancer. Inst.*, 1980, 65(2), 411-414.
2. Banu C., Alexe P., Buțu N., Rășmeriță D., Lungu Cornelia, Vizireanu Camelia – *Aditivi și ingrediente pentru industria alimentară*, Ed. Tehnică, București, 2000
3. Bartholomew B., Hill M.J. – The pharmacology of dietary nitrate and the origin of urinary nitrate, *Food Chem. Toxicol.*, 1984, 22, 789-795
4. Boukerche S., Aouacheri W., Saka S. – Toxicological effects of nitrate : biological study in human and animal, *Ann. Biol. Clin (Paris)*, 2007, 65(4), 385-391
5. Cardenas-Navarro R., Adamowicz S., Robin P. – Nitrate accumulation in plants : a role for water, *Journal of Experimental Botany*, 1999, 50, 613-624
6. Garban Z. – *Xenobiochimie: Tratat comprehensiv*, Vol. IV, Editura Eurobit, Timișoara, 2005
7. Ghibu G.D., Garban Gabriela, Falcă Corina, Baltă C., Precob V., Garban Z. - Dyshomeostatic effects of nitrates from drinking water on non-protein nitrogen metabolites in leporides, *Chem. Bull.*, 2008a, in press.
8. Ghibu G.D., Baltă C., Gabriela Garban, Sayti L, Corina Falcă, Garban Z. - Peculiarities of hepatic metallograms in rabbits after nitrates administration, *Journal of Agroalimentary Processes and Technologies*, 2008, 14, 361-366.
9. Murakami S., Ohno T., Bernardo J.F., Pfeifer C.A., Li T., Zhang Y., Dubey R.K., Branch R.A., Sabra R. - Reduced liver function is the trigger for renal sodium retention following portal vein ligation in the rat, *J. Gastroenterol. Hepatol.*, 1996, 11(9), 850-856.
10. Prakasa Rao E.V.S., Puttanna K. - Nitrates, agriculture and environment, *Current. Sci.*, 2002, 79, 1163-1168.
11. Rostkowska K., Zwierz K., Róžański A., Moniuszko-Jakoniuk J., Roszczenko A. - Formation and Metabolism of N-Nitrosamines, *Polish Journal of Environmental Studies*, 1998 7,(6), 321-325.
12. Roth A., Smith M.K. - Nitrite-induced iron deficiency in the neonatal rat, *Toxicol. Appl. Pharmacol.*, 1988, 96, 43-51.

13. Speijers G.J.A., van Went G.F., van Appeldoorn M.E., Montizaan G.K., Janus J.A., Canton J.H., van Gestel C.A.M., van der Heijden C.A., Heijna-Merkus E., Knaap A.G.A.C., Luttik R., de Ywart D. - *Integrated criteria document nitrate* (Appendix: effects), Report No. 758473012, Research for man and environment, National Institute of Public Health and Environment Protection, Bilthoven, Holland, 1989
14. Turner A.Y., Aizman R.I. - Characteristics of the sodium accumulating capacity of the liver, *Bull. Exp. Biol. Med.*, 1984, 97(3), 230-232.
15. Vermeer I.T.M., Pachon D.M.F.A., Dallinga J.W., Kleinjans J.C.S., van Maanen J.M.S. - Volatile N-nitrosamines formation after intake of nitrate at the ADI level in combination with an amine-rich diet, *Environ. Health Perspectives*, 1998, 106, 459–463.
16. Yu Z., Serra A., Sauter D., Loffing J., Ackermann D., Frey F.J., Frey B.M., Vogt B. - Sodium retention in rats with liver cirrhosis is associated with increased renal abundance of NaCl cotransporter (NCC), *Nephrol. Dial. Transplant.*, 2005, 20(9), 1833-1841.
17. Zabulyte D., Uleckiene S., Kalibatas J., Paltanaviciene A., Jascaniniene N., Stosik M. – Experimental studies on effect of sodium fluoride and nitrate on biochemical parameters in rats, *Bull. Vet. Inst. Pulawy*, 2007, 51, 79-82.
18. \*\*\* Ordonanța de Guvern nr.37 din 30 ianuarie 2002 pentru protecția animalelor folosite în scopuri științifice sau în alte scopuri experimentale.

# SODIUM, POTASSIUM, CALCIUM AND MAGNESIUM CONTENTS – ARE INDICATORS FOR GEOGRAPHICAL ORIGIN OF BEE'S HONEY FROM BANAT COUNTY?

Monica HĂRMĂNESCU<sup>1\*</sup>, Daniel POPOVICI<sup>2</sup>, Alexandru MOISUC<sup>1</sup>,  
Iosif GERGEN<sup>3</sup>

1. Banat's University of Agricultural Sciences and Veterinary Medicine, Faculty of Agriculture, Calea Aradului nr. 119, Timisoara, RO-300 645, Romania; 2. Banat's University of Agricultural Sciences and Veterinary Medicine, Faculty of Veterinary Medicine, Calea Aradului nr. 119, Timisoara, RO-300 645, Romania; 3. Banat's University of Agricultural Sciences and Veterinary Medicine, Faculty of Food Technology, Calea Aradului nr. 119, Timisoara, RO-300 645, Romania.

\* E-mail: monicaharmanescu@yahoo.com

## ABSTRACT

*This scientific paper presents our researches regarding the usage of Na, K, Ca, Mg contents and Principal Component and Classification Analysis for differentiate the honey samples purchased from two different villages from Romania: Berini (Timis District) and Cornea (Caras-Severin District). The Na, K, Ca and Mg contents were determined by flame atomic absorption spectrometry (F-AAS) with high-resolution continuum source ContrAA 300 spectrometer.*

**Key words:** macro-minerals, honey, flame atomic absorption spectrometry, multivariate analyses

## INTRODUCTION

Honey represents a sweet natural substance produced by the bees from flowers nectar or other sweet secretions of plants [Nanda et al., 2003]. After transport and transformation processes the nectar are stored like honey in the comb cells and it is used for bee nutrition (Bura et al., 2005). The nectar processing by bees refers to invert the saccharose in fructose and glucose, and the water content decreased at 18-20% (Özcan et al., 2006).

Honey can be easy assimilating, being a valuable nutritive product for children, athletes and convalescents. It is a good source of antioxidants, macro- and micro-minerals, monosaccharides, B vitamins, and enzymes (Juszczak et al., 2009). It contains potassium in higher quantity than sodium, that's way has a diuretic effect, being indicated in kidney diseases. Other macro-minerals from honey, like magnesium (with energetic roll) and calcium (muscular stimulator) are necessary in carbohydrates metabolism, in rahitism, spasmophily, and the water repartition in human's body (Socaciu, 2003). The sources of the mineral components of honey are its raw materials (nectar, honeydew) and pollen grains (Hernández et al., 2005), not bee organisms, as some researches suggest. Also it is a biological product of the bees, obtained without human interference. Its quality is unaffected by changes

which is possible to appear during technological processes. That's way the mineral composition of honey can be used to determine its geographical origin.(Madejczyk and Baralkiewicz, 2008).

The aim of this work was to differentiate the geographical origin of honey samples purchased from two different villages from Romania (Berini and Cornea) using Principal Component and Classification Analysis and the Na, K, Ca and Mg contents.

## MATERIALS AND METHODS

The analysed samples were different types of honey from Berini (Timis District) and Cornea (Caras-Severin District). Honey samples (10 g) were dry burning in quartz capsules at 650°C for 4 hours. After complete burning a nitric acid 0.5 N solution was added up to 50 mL. The obtained solutions were used for Na, K, Ca and Mg contents determination by flame atomic absorption spectrometry (F-AAS).

The multi-elements standard solution (1000 mg/L), used to obtain the calibration curves, was purchased from Riedel de Haen (Germany). The nitric acid 65% solution used was of ultra pure grade (Merck, Germany). All solutions were prepared using deionized water. All analyses were made in triplicate and the mean values were reported in mg/kg fresh product (ppm).

The determination of Na, K, Ca and Mg contents were made with ContrAA-300, Analytik-Jena device, by flame atomic absorption spectrometry (FAAS) in air/acetylene flame.

Statistical interpretation of obtained data was made using Principal Component and Classification Analysis implemented in Statistica-6 software.

## RESULTS AND DISCUSSIONS

The Na, K, Ca and Mg contents in analysed honey samples are presented in Table 1:

**Table 1.** The Na, K, Ca and Mg contents in analysed honey samples

| No. | Honey samples                   | Na<br>ppm   | K<br>ppm      | Ca<br>ppm    | Mg<br>ppm    |
|-----|---------------------------------|-------------|---------------|--------------|--------------|
| 1   | Mixed flowers honey Berini 2005 | 34.4        | 325.1         | 12.9         | 16.3         |
| 2   | Mixed flowers honey Berini 2006 | 31.6        | 323.5         | 39.3         | 25.7         |
| 3   | Acacia honey Berini 2005        | 64.0        | 485.4         | 17.5         | 34.7         |
| 4   | Acacia honey Berini 2006        | 60.8        | 930.9         | 22.7         | 35.9         |
| 5   | Acacia honey Cornea 2006        | 52.5        | <b>1251.9</b> | 148.4        | 130.1        |
| 6   | Acacia honey Cornea 2007        | 58.6        | 1246.4        | 147.5        | 130.6        |
| 7   | Lime honey Cornea 2007          | <b>82.7</b> | 1241.9        | <b>226.4</b> | <b>132.6</b> |

The major macro-mineral in analysed honey samples was potassium (325.1 - 1251.9 ppm), followed by calcium (12.9 – 226.4 ppm), magnesium (16.3 – 132.6ppm) and sodium (31.6 – 82.7 ppm).

Between the samples from Berini the acacia honey has higher content in sodium, potassium and magnesium than mixed flowers honey (64.0 ppm, 930.9 ppm, respectively 35.9 ppm). Mixed flowers honey from Berini (2006) has the highest calcium content (39.3 ppm).

Lime honey from Cornea (2007) has the highest sodium (82.7 ppm), calcium (226.4 ppm) and magnesium (132.6 ppm) contents among all analysed samples. The highest potassium content was determined for acacia honey from Cornea (2006).

In Figure 1 was represented the distribution of analysed honey samples depending on the Na, K, Ca and Mg contents. It is possible to observe that sodium, potassium, calcium and magnesium can be used to discriminate between the honey samples from Berini than those from Cornea. The first group is formed by acacia honey (2006 and 2007) and lime honey from Cornea; which has the highest contents in potassium, calcium and magnesium between all analysed samples. In the first group it is possible to observe that sodium content can differentiate the lime honey, which has the highest content (82.7 ppm) than acacia honey samples from the same village (Berini). The second group includes the mixed flowers honey (2005 and 2006) and acacia honey (2005 and 2006) from Berini, which have smaller concentrations in potassium, calcium and magnesium than the honey samples from Cornea.




**Fig.1.** The distribution of analysed honey samples function the Na, K, Ca and Mg contents

## CONCLUSIONS

The results of this research have shown that using Principal Component and Classification Analysis and the macro-minerals contents (sodium, potassium, calcium and magnesium) it is possible to differentiate between the honeys samples purchased from Berini than those from Cornea.

## REFERENCES

1. Bura Marian, Pătruică Silvia, Bura Alexandru-Vlad, *Tehnologie apicolă*, Editura Solness, Timișoara, 2005.
2. Hernández O.M., Fraga J.M.G., Jiménez A.I., Jiménez F., Arias J.J, Characterization of honey from the Canary Islands: Determination of the mineral content by atomic absorption spectrometry, *Food Chemistry*, 2005, 93, 449–458.
3. Juszczak Lesław, Socha Robert, Jacek Roznowski, Teresa Fortuna, Karolina Nalepka, Physicochemical properties and quality parameters of herbhoney, *Food Chemistry*, 2009, 113, 538–542.
4. Madejczyk Maria, Danuta Baralkiewicz, Characterization of Polish rape and honeydew honey according to their mineral contents using ICP-MS and F-AAS/AES, *Analytica Chimica Acta*, 2008, 617, 11–17.
5. Nanda V., Sarkar B. C., Sharma H. K., & Bawa A. S., Physico-chemical properties and estimation of mineral content in honey produced from different plants in Northern India, *Journal of Food Composition and Analysis*, 2003, 16, 613–619.
6. Özcan M., Arslan D., Ceylan D. A., Effect of inverted saccharose on some properties of honey, *Food Chemistry*, 2006, 99, 24–29.
7. Socaciu Carmen, *Chimia alimentelor*, Editura Academic Pres, Cluj-Napoca, 2003, ISBN 973-8266-79-3.



# MINOR AND TRACE ELEMENTS IN PARSLEY (*PETROSELINUM SATIVUM* HOFFM.) FROM THE WEST SIDE OF ROMANIA

Nicoleta G. HĂDĂRUGĂ<sup>1\*</sup>, Daniel I. HĂDĂRUGĂ<sup>2</sup>, Adina NEGREA<sup>2</sup>,  
Mihaela E. CIOPEC<sup>2</sup>, Petru NEGREA<sup>2</sup>

1. Department of Food Quality, Faculty of Food Processing Technology, University of Agricultural Sciences and Veterinary Medicine of Banat, Calea Aradului nr. 119, RO - 300 645 Timișoara, Roumania; 2. Faculty of Industrial Chemistry, University "Politehnica" Timișoara, P-ța Victoriei nr. 2, RO-300006 Timișoara, Roumania

\* E-mail : nico\_hadaruga@yahoo.com

## ABSTRACT

The paper presents a study on the variation of the concentration of main elements from the parsley's pulp, shell and leaves harvested from the west side of Romania. The results were compared with those obtained for the similar samples harvested from the hungarian region (close to the border). The representative samples were collected from different sides of West Romania or from the East Hungary and prepared according to AOAC Official Methods of Analysis. The concentrations of main elements (Na, K, Fe, Cu, Zn, Mn, Pb, Cd, and Ni) were determined by Atomic Absorption Spectroscopy.

**Key words:** parsley, spice, metals, nonmetals, atomic absorption spectroscopy, ecology.

## INTRODUCTION

Vegetables are vital for the human diet, and particularly spices and aromatic herbs are very important for seasoning the food, providing the well-known trace elements. Some spices exhibits antioxidative properties (Zöchling S. et. al., 2002; Demo A. et. al., 1999; Belitz H.-D. et. al., 1987) or antimicrobial activity (Arora D.S. et. al., 1999; El Hamss R. et. al., 2003; Nielsen, P.V. et. al., 2000). Parsley (*Petroselinum sativum*, Hoffm.) is among the most valuable vegetables and it is a very rich source of vitamin C,  $\beta$ -carotene and mineral constituents (Lisiewska Z. et. al., 1997 ; Mohamed A.E. et. al., 2003). Minor elements such as Ca, Fe, K, Mg, and Na as well as trace elements such as Co, Cu, Mn, Ni, Pb, and Zn, were determined using the atomic absorption spectrophotometry (Mohamed A.E. et. al., 2003). A number of elements such as As(III), Al, B, Be, Cd, Co, Cr(IV), Cu, Mo, Ni, Se(IV), and Ti can be harmful to plants and humans even at quite low concentrations (Bowen H.J.M., 1979). These elements occur in variable concentrations in living tissues and are believed to be acquired by the body through environmental contamination (air and soil pollution). The aim of this work are to (1) assess the minor and trace elements in core, shell and leaves of parsley harvested from the west side of Romania for bioavailability or toxicity to humans, and (2) study of the correlational

parameters between the concentrations of these elements and the part of parsley or the harvesting region.

## MATERIALS AND METHODS

### *Sample collection.*

Five samples of parsley (*Petroselinum sativum*, Hoffm.) were collected from different region of Timiș (Cenad, Ghiroda, Moșnița Nouă, and Utvin) and from the east side of Hungary (close to Romanian boundary).

### *Sample preparation.*

The samples were washed with water several times, finally with distilled water and dried at room temperature. The core, shell (maximum 0.5 mm thickness) and leaves were collected in duplicates. These samples and a blank were put in porcelain crucibles and ashed in a furnace oven at 550°C for 48 h. The white residues were moistened with 5 ml 37% HCl (analytical grade, Merck Chemical Co.), boiled 2 minutes, evaporated to dryness on a hot plate, and heated on steam bath for 3 h. The residues were moistened with 5 ml 37% HCl, boiled 2 minutes, and then 50 ml distilled H<sub>2</sub>O were added; the clearly solutions were heated on a water bath for 5 minutes, filtered and washed with distilled water. The filtrate was diluted to 200 ml with distilled water.

### *AAS Analysis.*

Atomic absorption spectroscopic standard solutions (1-100 mg/l for Na and K, 0-15 mg/l for Fe, 0-6 mg/l for Cu and Mn, 0-1.5 mg/l for Zn, and 0-7.5 mg/l for Pb, Cd, and Ni, Merck&Co. Inc.) were used for calibration and for standard curves. A Varian-1100 atomic absorption spectrophotometer was used for the determination of element concentrations. The mean of two determinations was used for statistical analysis. Element hollow cathode lamps were used at recommended conditions.

### *Statistical analysis.*

Statistical analysis of the database includes ANOVA, cluster analysis and principal component analysis (PCA), performed with *in house* programs. The data include the elements compositions.

## RESULTS AND DISCUSSIONS

### *Minor elements.*

From the atomic absorption spectrophotometric data, the most concentrated minor elements were Na and K, with values between 748 to 17062 mg/kg raw sample for the first, and between 600 to 58657 mg/kg raw sample for the second element. The upper limits for Na appear for samples harvested from Utvin (both core and shell), and the lower ones for samples from Cenad. By far, the most concentrated sample in element K was that from Cenad (especially for core part), and that less concentrated was the sample from Ghiroda (for core and shell).

Element Fe exhibits values in the range 507 – 542 mg/kg, the most concentrated sample being from Utvin (shell and leaves parts) and the lower limit for this element was 7.3 and 7.7 ppm (sample from Cenad – core and leaves). Major samples contains element Fe in the range of 7 to 541 ppm.

The atomic absorption spectrophotometric data are indicated in Table 1, and the relative concentration of minor elements in parsley are shown in Figs 1, 2, and 3.




**Fig 1.** Minor element (Na) concentrations (ppm) in parsley




**Fig. 2.** Minor element (K) concentrations (ppm) in parsley




**Fig. 3.** Minor element (Fe) concentrations (ppm) in parsley

**Table 1a.** Minor element (Na, K and Fe) concentrations (mg/kg raw sample or ppm) in parsley from west side of Romania

| Sample name                 | Code               | Na         | K         | Fe       |
|-----------------------------|--------------------|------------|-----------|----------|
| Parsley Moşniţa Nouă (core) | <b>Mos(core)</b>   | 1578±1414  | 5317±67   | 18.3±1.4 |
| Parsley Cenad (core)        | <b>Cen(core)</b>   | 1135±71    | 58658±141 | 7.7±0.7  |
| Parsley Cenad (shell)       | <b>Cen(shell)</b>  | 748±8      | 6142±42   | 52.5±3.5 |
| Parsley Cenad (leaves)      | <b>Cen(leaves)</b> | 1747±14    | 3894±21   | 7.3±0.7  |
| Parsley Ghiroda (core)      | <b>Ghi(core)</b>   | 6965±7     | 601±2     | 13.8±2.8 |
| Parsley Ghiroda (shell)     | <b>Ghi(shell)</b>  | 6905±71    | 1571±30   | 50.5±3.5 |
| Parsley Ghiroda (leaves)    | <b>Ghi(leaves)</b> | 5371±45    | 13287±23  | 410±35   |
| Parsley Utvin (core)        | <b>Utv(core)</b>   | 17063 ±216 | 16279±424 | 53.5±2.1 |
| Parsley Utvin (shell)       | <b>Utv(shell)</b>  | 11238±424  | 13197±283 | 507±131  |
| Parsley Utvin (leaves)      | <b>Utv(leaves)</b> | 4070±21    | 9916±28   | 542±78   |
| Parsley Hungary (core)      | <b>Hun(core)</b>   | 1507±92    | 6601±4    | 11.0±1.1 |
| Parsley Hungary (shell)     | <b>Hun(shell)</b>  | 1139±63    | 2746±70   | 43.9±1.7 |

**Table 1b.** Trace element (Cu, Mn, Zn, Pb and Cd) concentrations (mg/kg raw sample or ppm) in parsley from west side of Romania.

| Sample name                 | Code               | Cu       | Mn       | Zn       | Pb       | Cd       |
|-----------------------------|--------------------|----------|----------|----------|----------|----------|
| Parsley Moşniţa Nouă (core) | <b>Mos(core)</b>   | 1.68±0.4 | 4.2±0.6  | 3.1±0.1  | 2.5±0.1  | 1.2±0.23 |
| Parsley Cenad (core)        | <b>Cen(core)</b>   | 1.44±0.3 | 3.4±0.7  | 2.5±0.6  | 1.55±0.5 | 1.67±0.3 |
| Parsley Cenad (shell)       | <b>Cen(shell)</b>  | 1.9±2.3  | 8.7±0.7  | 3.3±0.9  | 1.8±0.1  | 1.15±0.2 |
| Parsley Cenad (leaves)      | <b>Cen(leaves)</b> | 0        | 0        | 0        | 0        | 4.2±0.5  |
| Parsley Ghiroda (core)      | <b>Ghi(core)</b>   | 0.66±0.1 | 4.0±1.1  | 2.4±0.3  | 0.7±0.1  | 1.1±0.1  |
| Parsley Ghiroda (shell)     | <b>Ghi(shell)</b>  | 0.8±0.08 | 3.6±0.6  | 2.8±0.1  | 0        | 1.0±0.07 |
| Parsley Ghiroda (leaves)    | <b>Ghi(leaves)</b> | 0        | 7.8±0.2  | 17.1±0.3 | 0        | 5.06±0.1 |
| Parsley Utvin (core)        | <b>Utv(core)</b>   | 5.08±0.5 | 22±3.0   | 10.8±0.8 | 0.9±0.03 | 2.6±0.04 |
| Parsley Utvin (shell)       | <b>Utv(shell)</b>  | 5.9±0.4  | 13.1±2.3 | 14.0±1.1 | 0        | 1.3±0.1  |
| Parsley Utvin (leaves)      | <b>Utv(leaves)</b> | 18.6±1.6 | 18.7±1.9 | 41.1±3.1 | 0        | 1.4±0.5  |
| Parsley Hungary (core)      | <b>Hun(core)</b>   | 2.68±0.1 | 16.0±2.5 | 4.9±0.3  | 3.7±0.2  | 1.5±0.2  |
| Parsley Hungary (shell)     | <b>Hun(shell)</b>  | 0.71±0.7 | 4.8±0.4  | 3.1±0.04 | 0        | 0.9±0.04 |

### *Trace elements*

For the trace elements (Cu, Mn, Zn, Pb, Cd, and Ni) the concentrations are about 1000-fold lower than minor elements. The most concentrated is Cu (up to 18.6 ppm), Mn (up to a range 22 ppm), and Zn (up to 41.1 ppm). The lower concentrations were found for Cd and Pb, even if these harmful elements exhibit quite higher values. The higher concentration for Cu, Mn, and Zn were found for the samples from Utvin (especially in leaves), but the Pb concentration in these samples was lower. The core part of parsley is generally most concentrated in Pb, than the shell part. Cd was most

concentrates in leaves, in almost all cases, and Ni was not found in any part of parsley (under detection limit). The AAS data are indicated in table 1(a,b) and Figs 4 and 5.




**Fig. 4.** Trace element (Zn, Mn, Cu) concentrations (ppm) in parsley




**Fig. 5.** Trace element (Pb, Cd) concentrations (ppm) in parsley

### *Statistical analysis*

The multivariate analysis by PCA (principal component analysis), using all data (Table 1), indicate that the samples from Utvin and Ghiroda were classified in the upper side of the score plot (Figure 6), and the Moșnița, Cenad, and Hungary samples in the lower side, closely grouped (exception the Cenad core sample). The Na and K concentrations were responsible for this classification (loadings plot). The explained variance of the data was 89% for PC<sub>1</sub> and 11% for PC<sub>2</sub>.

The samples were classified by core, shell and leaves parts only by PCA analysis using the concentrations for trace elements (Mn, Zn, Pb, and Cd). The samples containing leaves were localized in the lower side of the score plot, and the samples containing core or shell were classified in the upper side (Figure 7). The explained variance of the data was 79% for the first principal component and 18% for the second one. The main elements for this classification were Mn and Zn.

From hierarchical cluster analysis of all data (minor and trace elements), the similarity of the samples was relatively good, excluding the core sample from Cenad (with a very high level of K). Very similar were the samples from Moșnița, Cenad and Hungary. The samples from Utvin and Ghiroda were also similar (Figure 9).




**Fig. 6.** Score plot for all data from PCA analysis




**Fig. 7.** Score plot for trace element data from PCA analysis




**Fig. 8.** Loading plot for trace element data from PCA analysis




**Fig. 9.** Hierarchical cluster analysis for parsley samples

## CONCLUSIONS

The concentration of trace elements in vegetables is very important for our health and from environmental ecology point of view. The samples generally exceed the maximum concentration values for almost all elements. For Cu, this maximum concentration (5 mg/kg) were exceeded for samples from Utvin, Mn and Zn are more concentrated in samples from Utvin and Ghiroda (leaves), and more concentrated in Pb are samples from Hungary. Cd concentration exceeded the maximum concentration limit (0.05 mg/kg) for all samples, more concentrated being the leaf part of parsley.

These results are good correlated with de distances from Timișoara (industrial city) to the region from where the parsley samples are harvested. Utvin and Ghiroda are more closed to Timișoara city (up to 5 km).

By multivariate analysis (principal component analysis, PCA, and hierarchical cluster analysis, HCA) the samples (harvested sites) have been classified in two main groups: Utvin - Ghiroda group (close to Timișoara city) and Cenad – Hungary group (close to Romanian-Hungary boundary).

## REFERENCES

1. Arora, D.S., Kaur, J.: "Antimicrobial Activity of Spices", *Int. J. Antimicrobial Agents* 1999, 12, 257-262.
2. Belitz H.-D., W. Grosch : *Food Chemistry*, Springer-Verlag, New York - Berlin - Heidelberg - London - Paris - Tokyo - Hong Kong - Barcelona – Budapest, 1987.
3. Bowen H.J.M.: *Environmental Chemistry of Elements*, Academic Press, London, 1979.
4. Demo A., Petrakis C., Kefalas P., Boskou D.: "Nutrient Antioxidants in some Herbs and Mediterranean Plant Leaves", *Food Res. Int.* 1999, 31, 351-354.
5. El Hamss R., Idaomar M., Alonso-Moraga A.: Muñoz Serrano, A., "Antimutagenic Properties of Bell and Black Peppers", *Food Chem. Tox.* 2003, 41, 41-47.
6. Lisiewska Z., Kmiecik W.: "Effect of Freezing and Storage on Quality Factors in Hamburg and Leafy Parsley", *Food Chemistry* 1997, 60, 633-637.
7. Mohamed A.E., Rashed M.N., Mofty A.: "Assessment of Essential and Toxic Elements in Some Kinds of Vegetables", *Ecotox.&Environm. Safety* 2003, 55, 251-260.
8. Nielsen P.V., Rios R.: "Inhibition of Fungal Growth on Bread by Volatile Components from Spices and Herbs, and the Possible Application in Active Packaging, with Special Emphasis on Mustard Essential Oil", *Int. J. Food Microbiology* 2000, 60, 219-229.
9. Zöchling S., Murkovic M., Pfannhauser, W.: "Effects of Industrially Produced Flavours with Pro- and Antioxidative Properties on the Formation of the Heterocyclic Amine PhIP in a Model System", *J. Biochem. Biophys. Methods* 2002, 53, 37-44.
10. \* \* \* *Spices. Flavor Chemistry and Antioxidant Properties*, S.J. Risch and C.-T. Ho. (eds), American Chemical Society, Washington DC.



# CONCLUSIONS CONCERNING THE SOURCE AND THE ENVIRONMENT CONCENTRATION WITH HEAVY METALS ON THE ETNO-HISTORICAL BANAT

Gheorghe IANOS

West University of Timișoara, Blv. Vasile Pârvan nr. 4, RO-300 223 Timișoara, Roumania

\* E-mail : ianos@cbg.uvt.ro

## ABSTRACT

*The Banat soil between the Carpathians, Tisa, Mureș and Danube have now a relative high content in heavy metals. Because of the lack of industrial pollutants, the causes which have favoured the accumulation of heavy metals can be put only on the mountain zone as well as the modality of their transport towards the west, towards the piedmont areas and the plain ones.*

*In the long singenetical process, under the action of the weathering solutions, both in the previous geological periods and in the present period, there have been a lot of continuous processes of delivering the constituent elements. This delivering depends in general on the solubility of the chemical compounds, their valence and the climatic and litological circumstances.*

*The researches followed the understanding on the geochemical importance of the soils from the etno-historical Banat.*

*The explication is based on the native occurrence of these elements in the rocks of south-western Romanias' Montains. The more humid climate (of oceanic influence and more warmer (the Mediteranean influence) has favoured the balteration and the Panonic fore deep which was a lake in this first phase and a swamp later, favoring the accumulation. In the last accumulative medium, the concentration of metal elements have increased also on the base of the sulfur of organic matter in an anoxic medium.*

**Key words:** heavy metals, pollution, Banat, Mureș-Tisa-Danube Euregion.

## INTRODUCTION

The geogenous and the anthropogenous abundance of heavy metals in Banats'rocks and soils (between the Carpathians-Tisa River-Mureș River and Danube) is conditioned by the paleogeographical evolution, by the type and the direction of weathering that occurred especially during the Quaternary as well as by the direction and the characteristics of the pedogenetical processes.

The Banat soils do now have a high content of heavy metals that are almost always above the background domains. In the absence of predominant industrial poluters the causes that have favoured the accumulations of heavy metals can only be explained by the high geochemical backgrounds of the rocks and of the parental materials of the eastern mountain zone and by the manner of transport

towards these areas and the modifications that occurred in the chemistry of the soils during the pedogenesis process.

The researches done took into consideration the knowledge about the geochemical endowment of Banat in order to appreciate the capacity of the soils of being able to reach or not the extraloadings of heavy metals. The identification of heavy metals on levels of content and their map materialisation is allowing the characterisation of the place, the appreciation of their origin (autochthonous, allochthonous or anthropogenous) as well as the establishing of some correlations between the total forms and the mobile ones of each studied element.

The exploration on the Serbian Banat fields and the conclusions results, made on the Serbian Banat fields, is just a hardship. Confirmation or refutation of the assumption result thoroughgoing study or reorientation of investigation.

## **MATERIALS AND METHODS**

Towards estimating the geochemical background of the ethno-historical Banat, within the surface of the Romanian Banat there have been taken about 400 soil samples from the horizon placed at different depths (from 0-150 cm), from over 100 places. The analysis methods relied on the weathering of the prevailed samples with a mixture of nitric, sulphuric and perchloric acids, in a 2 : 0,1 : 1 (I.C.P.A. - 1986) proportion and spectrometrical atomic absorption dosimetry in variant of air-acetylene flame atomisation.

The manner in which the position cartograms were made presenting the area disposition of the researched oligoelements relied in the superposition of the following maps: the map of main geomorphological units of Banat (Ianos, 1985), the geological map (Ianos, 1987), the map of the rocks and of the parental materials, the map of the intensely mineralised materials (fig.1), the map of the soil associations (Ianos, 1985), the map of Banat along XVIII and XIX century (Ianoş-1997), the map of the depth of the underground waters (Ianoş 1997), the map of the humus supplies (0-50 cm)(fig.2), the map of the soil acidity (fig.3).

Functions of the particularities of occurrence of the researched metal microelements (Cd, Co, Cu, Cr, Fe, Mo, Mn, Ni, Pb, Zn) in the subiacent and parental rocks, of the environment and the manner in which these elements are taken out of rocks by weathering, of the behaviourism of these rocks during the transportation as well as of the situation created during the sedimentation there have been delimited new areas with similar concentrations on content levels. On the basis of the above presented considerations to which there can be added lithological, diagenetic or pedogenetic particularities, in the zones where there were not gathered and analysed samples the conclusions that could be drawn were extrapolated to that the future analytical determinations to confirm or to deny the formulated suppositions

## **RESULTS AND DISCUSSIONS**

### ***1. The conditions of occurrence of the heavy metals in the rocks and soils of Banat***

In its eastern part, Banat is limited by more or less homogenous mountain massifs. Created during the alpine orogenesis, they have taken at the same time or after an intense magmatic activity generated by the Postlaramic tectonic movements after which on the fault line there were pushed towards the surface the Banatic intrusions especially formed on granodioritic type of rocks (Oncescu-1965,

Folea-1980). In paralely there have been taken place large processes of contact metamorphism and selective metasomatoza and as a result there were replaced especially by transformation of the limestones the skarns (Gheorghită-1975, Petrulian-1979).





**Fig. 1** Arels with anomale mineralizations in the mountain Banat region

**Legend:** 1. Complex sulphates; 2. Anomal areals with mineralizations; 3. Direction of spreading for the eroded materials; 4. General direction of transport and sedimentation of the intense mineralized materials.

**Fig. 2**  
Values of attentions of the heavy metals concentration in the soils of the Ethno-historical Banat



Later, the endo and exogene skarns were influenced by a hydrothermal metamorphism (Folea-1980) with the making of polymetal sulphurs (pyrite, blende,

calcopyrite, galena, aikinit, bismuthit, cerussite, crocoite, piromorphite, molybdenite etc).

A short inventory of the areas with intense mineralisation are indicating us a zone of positional accumulation in the nord-west part of Banat (Metaliferi Mountains) where there are predominating the polisulphurs of plumb, copper, zinc (fig.4). Towards west and south west Savu (1970) mentioned rich mineralisations of mollibden and manganese or iron in the eastern part of Zarand Mountains. From both zones, the Mureş River and its tributarries have taken denudated materials and put them towards west building the hills and the plains in the central and western part of Banat. There follows, on the south, the metalogenetical areals in the mountain massifs Poiana Rusca-Ruschița (lead, cooper, zinc, molybdenum), Dognecea (iron, copper, chromium), Anina-Sasca ( cobalt, nickel), Almăj-Moldova Nouă (copper) (fig.1).




**Fig. 3** Humus reseve of Banat Soils on the depth of 0-50 cm (t/ha)

Legend: 1. < 60; 2. 61-120; 3. 121-160; 4. 161-200; 5. 210-300; 6. > 301.




**Fig. 4** Soils reaction in Banat (pH units) on the depth of 0-29 cm.

Legend: 1.< 5,80 (strong medium acid);2. 5,81-6,80 (weak acid);3. 6,81-7,20 (neutral); 4.> 7.21 (alkaline).

During the geological time, but especially during the Quaternary, when the frequent climatical oscilations have been favourising an intense physical weathering in stages and a pronounced chemical weathering in the interstages, from the rocks of the mountain massifs there was released a part of the constituent chemical elements.

The positive epirogenetical movements of the Carpathians and the drainage of the Panonic lacustrine domain by Danube pearcing through the Iron Gates have been accelerating the denudational processes, those of transport of the materials resulted as well as an accelerated sedimentation in the piemountaneous zones and then on the plain zones in the east-west direction. On the basis of these considerations we have to underline the fact that not only the superficial lithological layers in the plain or piedmountain zones are loaded with heavy metals but also

thick packs of rocks placed on deep depths. This fact is explaining some overloadings with arsenium of the depth waters (60-200 m), there were registered in the northern part of the Mureş River that are also continuing in Hungary.

A last stage in the process of rocks and soil loadings with heavy metals is conditioned by the filling with alochtonous materials of the vast areas of subsidence in the eastern Banat, zones occupied by the end of the XVIII<sup>th</sup> Century by large swampy domain where there existed a varied and rich hydrophile fitocenosis that was decomposed and there resulted rich amounts of sulphur (H<sub>2</sub>S). Because the waters of the main rivers that were eroding Banat used to carry in suspension fine particles of erroded material or ions of the heavy metals in solution and as the majority heavy metals have a special affinity for the sulph, at floods these elements were bloked in the fine texturated materials loaded with sulphur compounds, fact that leads to high occurances of chemical elements in the clayed rocks, especially the smectical ones.

The tectonic and climatic stability at the beginning of the holocene marked a diwinding of the erosional processes, of the sedimentation, as well as an amplification of the solification processes function of the time lift from the beginning of the solification process of the granulometrical making of the parental rocks and of the soil, of the amount of humified organical matter, of the pH and rH. of the solutions that are circulating in the soil, heavy metals placed here together with the solification materials have been suffering concentration modifications by lateral or vertical movements ascendent or descendent.

On the basis of the analytical data there can be inferred that in the Banat soils the metal microelements that were researched are especially to be found in the structure of the primary and secondary minerals as well as kept by different physico-chemical mechanisms by the organical coloides and the mineral ones. The total supplies and the degree of mobility is conditioned by nature of the solification rock and by the processes of pedogenesys (eluvial-iluvial, bioacumulation, oxido-reducing). Generaly in reduced and moderate weathering conditions, the influence of the rock is determinant and in the conditions of advanced weathering the main role goes to the pedogenetical processes.

Generaly as regarded and correlated with the data obtained by Lăcătuşu (1992) for Romania or by Adriano (1986), Alloway (1990) for other zones of the world, the microelements determined in the Banat soils are in a concentration (mean values) of 1.2 to 3 times higher, the maximum values being characteristic especially for the lead, cadmium, cobalt and chromium.

### ***The distribution of the main heavy metals (cadmium and lead) in Banats'rocks and soils***

In order to give examples, there have been selected two chemical elements with high toxic values. Other seven elements were selected and presented in publications made in last years (Ianoş, 1999).

**The cadmium.** The high release of the cadmium from the primary rocks in the special conditions of transport towards the western accumulative areas make the Banat soils contain important amounts of cadmium (1.0-1.8 ppm) with a mean value of 1.3 ppm. In comparison with the value of the clark, the mean cadmium content in the Banat rocks is 10 times higher and in comparison with the general mean content in the soils, 4.3 times higher (Lăcătuşu-1993).



The long weathering conditions makes that in rezistants, cadmium be almost totally missing and its affinity for the sulphur determine an enriched precipitation and concentration in the hidrolizate especially in the areas that do predominantly have oxide-reducing processes (the clayed muds in the subsidence zones). The uniformity of cadmium concentrations can be owed to the values generally reduced of the cadmium in the rocks (less than 2 ppm) or on the working methodology, the determination device not being able to sustain the exactness of some determinations at low amounts.

The smallest cadmium concentrations (less than 1 ppm) are to be found in the evaluated soils on the bulk deposits on the dejection cones of Mureș, Timiș, Bârzava, Caraș, Nera Rivers (fig.5). In the remaining ones the cadmium contents are slightly higher (1.3-1.8 ppm).

The later improvement with cadmium of the soils especially of those in the podzolite series in the piedmountain areas can be explained by the absorbtion of the element in the oxidazed sediments here manganese rich.

On the genetic types of soil there can be seen a moderate degree of variation with a slight tendancy of concentration in argiluvissols. During the background domain (1.2-1.5 ppm), the cadmium is presenting a moderate degree of variation with a slight tendancy of concentration in argiluvissols and the clayed soils (vertisols). Beneath these limits there are to be found the chernozems, the cambisols and fluvisols in the Mureș, Timiș, Bârzava Plains and in the Lugoj Hills, Pogăniș Hills and the Almăj, Oravița, Moldova Nouă Depressions.

**The lead.** Released by the rocks as a result of some weathering process the plumb is oxidated and reduces its mobility especially in the presence of the suplh equalising "in situ" the lead concentrations of the eluvial layers. Taken by the groover waters and then by the running waters rich in calcium cations the lead will make new combinations especially with  $\text{CaCO}_3$  especially in the sediments and soils of the depressionary corridors (>30 ppm). In the subsidence areas with sediments rich in organic sulphurs, the lead finds storage conditions in the extremely clayed soils in the Aranca Plain in the inferiour sector of the Timiș Plain or Caraș Depression (fig. 6).




**Fig. 5** The state of assurance with cadmium (total forms) of Banat soils (depth 0-40 cm)

1. little (<1,0 ppm); 2. medium (1,1-1,5 ppm); 3. great (1,6-1,9 ppm); 4. very great (>2.0 ppm):

**Fig. 6** The state of assurance with lead (total forms) of Banat soils (depth 0-40 cm)

1. little (<20 ppm); 2. medium (21-30 ppm); 3. great (>31 ppm); 4. sampling stations.

On the background of some more reduced sulphur concentrations of the porosity and of an enriched permeability in the loessic plains, covered with chernozems, the lead will easily realise bicarbonate compounds easily removable by vertical or laterally migrations situation that in this areas presents more reduced concentrations of total lead (<20 ppm). In this conditions the lead is oscilating with a large interval of values between 15 and 40 ppm with a medium value of 28.5 ppm, concentrations that are also over the values of the clarks and of the mean contents of other similar zones in the country and in the world.

## CONCLUSIONS

1. The heavy metals in the rocks and soils of Banat come from the eastern mountain zone by weathering processes, desintegration, transport and resedimentation.

2. The enrichment with heavy metals of the soils in the Banat 'hills and plains is conditioned by the reaction of the transporting solutions, of the sedimentation background as well as of the sulphur content.

3. Function of the conjunctural conditions of releasing transport and sedimentation most of the heavy metals presented in the Banat soils have a superior abundance in comparison with the neighbouring areas or other parts of the world.

4. Areas of anomal concentrations are placed in the subsidence zones occupied by the vertisols or at the content between the piedmountain and the mountain formations.

## REFERENCES

1. Adriano D.C.: Trace elements in the terrestrial environment, Springer Verlag, New - York-Berlin-Heidelberg-Tokio, 1986.
2. Alloway B.J.: Heavy metals in soils, John Wiley and Sous, Inc., New York, 1990.
3. Fiedler H.J, Rosler H.J.: Spurenelemente in der Umwelt, Ferdinand Enke Verlag, Stuttgart, 1988
4. Gheorghiuța Ioana : 1975, Studiu mineralogic și petrografic al regiunii Moldova Nouă, St. teh. și ec., seria I, nr. 11, 1975, Institut. Geol. București .
5. Grisellini Fr.: Versuch einer politischen und natürlichen Geschichte des temeswawere Banats in Briefen an Standespersonen und Gelehrte, I, II, Vienn, 1779.
6. Folea I.: Orientarea cercetărilor geologice și tehnologice pentru valorificarea rațională a potențialului de minereuri neferoase din România, St. teh. și ec., seria A, nr. 14, 1980, Institut. Geol., București.
7. Ianoș Gh., Goian M.: Solurile Banatului-geneză și caracteristici agrochimice, Ed. Mirton, 1995, Timișoara, 292 p.
8. Ianoș Gh.: Considerations of the making and evolution of the soils in the Banat; in Geographyc Monographs of European Regions, Novisad-Timișoara-Szeged, 1997, pag. 79-87.

9. Ianoş Gh., Puşcă I., Goian M. :Solurile Banatului; condiții naturale și fertilitate, Ed. Mirton, 1998, Timișoara, 392 p.
10. Ianoş Gh.: Geogenous and anthropogenous loadings with heavy metals in the rocks and soils of etno-historical Banat. In Proceedings of the regional conference of geography "Danube-Criş-Mureş-Tisa Euregion geoeconomical space of sustainable development, Timișoara-Novisad, Szeged, Tübingen, 1999, Editura Mirton Timișoara, pag. 205-223
11. Klocke A.: Richtwerte'80: Orientierungsdaten für tolerierbare Gesamtgehalte einiger Elemente in Kulturböden, Mitt. VDLUFA, H.2, 1980.
12. Lăcătușu R., Ghelase Ileana : A supra abundenței metalelor grele din sol, Mediul înconjurător, vol. III, nr. 4, 1992, pag. 45-52, București.
13. Lăcătușu R., Răuță C., Kovacsovic Beatrice, Lungu Mihaela, Ianoş Gh., Mihăilescu A. : Abundența metalelor grele în solurile agricole din Banat, Știința solului, nr. 1-2, 1993, București.
14. Oncescu N.: Geologia României, Ed. Tehnică, 1965, București.
15. Petruțian N., Sandu D., Popescu Rodica: Contribuții la cunoașterea mineralizației polimetale din corpul Simon Iuda (Ocna de Fier), St. Teh. și ec., seria I, nr. 16, 1979, Institut. Geol. București.
16. Rankama K., Sahama T.: Geochimia, Ed. Tehnică, 1970, București.
17. Savu H., Vasiliu Cecilia, Udrescu Constanța: Geochimia și petrologia ofiolitelor din prima etapă de evoluție a magmatismului inițial alpin din Masivul Drocea, Dări de seamă ale șed., vol. LVI, 1970, Institut. Geologic, București.
18. \*\*\* Metode de analiză chimică a solului, ASAS-ICPA, 1986, București.



# METAL ION COMPOSITION OF STINGING NETTLE (*URTICA DIOICA* L.) LEAVES IN VIEW OF THE NUTRITIVE VALUE

Noémi KOCZKA <sup>1</sup>, Dóra PÉTERSZ <sup>1</sup>, Éva STEFANOVITS-BÁNYAI <sup>2</sup>

*1\** Institute of Horticultural Technology, Faculty of Agricultural and Environmental Sciences, Szent Istvan University, Pater K. u. 1, H-2103 Gödöllő, Hungary; *2.* Department of Applied Chemistry, Faculty of Food Science, Corvinus University of Budapest, Villanyi ut 29-31, H-1118 Budapest, Hungary

\* E-mail: Koczka.Noemi@mkk.szie.hu

## ABSTRACT

*The aim of our investigations was to determine the macro and micro element components in leaves of common nettle (*Urtica dioica* L.) of different origin and the total phenol content and antioxidant capacity (by FRAP method) of extracts from them. Plant materials were collected from a meadow near the city Paks (Hungary) and from along a roadway near Paks. These samples were compared to commercial teas (drug from pharmacy, and teas from conventional and organic collection). Well detectable differences were obtained for the investigated parameters between the samples. Our results showed that the regular control of element composition of teas is very important from the viewpoint of health protection because the place of growing has a big influence on the accumulation of some elements in the nettle plant. Closed relations could be found between the antioxidant capacity and concentrations of some metal elements, which influence the nutritive value of herbs*

**Key words:** nettle, *Urtica dioica* L., metal element, antioxidant capacity

## INTRODUCTION

Medicinal plants are very important for our health, in addition to active ingredients and vitamins, mineral elements also have a big influence on the human organism. Positive or negative effects of plant antioxidants depend also on the macro and micro element content since complex binding and the too low or too high uptake of elements modify the biochemical processes, not to mention the uptake of toxic elements (Szentmihályi and Then 1999, Blazovics et al. 2003).

Abiotic environmental factors have a high influence of nutritive value of the plants. Lead, cadmium, copper, zinc and nickel are metals frequently reported to have the highest impact on organisms. The metal uptake from the soil by plants is an introductory pathway for heavy metals to enter the foodchain. Plant species differ in ability to absorb and allocate nutrients. *Urtica dioica* L. is not considered as hyperaccumulator and does not exhibit heavy metal tolerance but it has been mentioned many times as a good bioindicator (Sawidis et al. 1995, Singh et al. 1995, Tack and Verloo 1996, Prasad and Freitas 2003).

Leaves of stinging nettle (*Urtica dioica* L.) are a rich source of trace elements, absorbing and accumulating them. Due to its active ingredients nettle has

---

„Metal Elements in Environment, Medicine and Biology”, Tome VIII, pp. 189-194, Publishing House “Eurobit” Timișoara, 2008

Proc. of 8<sup>th</sup> Int. Symp. of Roumanian Academy - Branch Cluj-Napoca and Branch Timișoara, 2008, December 5-6, Timișoara, Roumania

anti-arthritic, antispasmodic, diuretic, astringent, tonic and expectorant effects. Dried nettle leaves are ordinarily used in herb tea mixtures. Fresh leaves also can be consumed as a vegetable similarly to spinach, and this usage begins to spread in Hungary nowadays. (Neumayer 2000, Dudas 2008)

## MATERIALS AND METHODS

The aim of our investigations was to compare the macro and micro element composition of common nettle (*Urtica dioica* L.) leaves of different origin. Plant materials were collected from a meadow near the city Paks (Hungary) and from along a roadway near Paks. These two nettle populations were defined as the same genotype. Samples were formed from 10 subsamples each collected from a 1 m<sup>2</sup> area. Plant material was harvested in July and dried at 30°C. Samples were compared to commercial teas (drug from pharmacy, teas from conventional and organic collection).

For analysis of element composition, 0,2 g of dried and pulverized samples (duplicated) were taken. The following elements were determined by ICP -OES (Thermo Jarrell Ash Co, ICAP 61): Al, As, B, Ba, Ca, Cd, Co, Cr, Cu, Fe, Hg, K, Li, Mg, Mn, Mo, Na, Ni, P, Pb, Se, Si, Sr, Ti, V, Zn. The concentration of metal ions is given as mg/kg dry weight.

Extracts were prepared from 1 g of dried leaves which were infused with 100 ml boiling water or aqueous ethanol (25°C; water/ethanol 80/20, v/v). The ethanolic and aqueous extracts were stored at room temperature for 72 and for 24 h. After centrifugation (13000 rpm, 10 min), the supernatants were analysed.

Total phenol content was measured using the method of Singleton and Rossi (1965). The phenol content. For determining the antioxidant capacity the FRAP method was used (values are given in mmol ascorbic acid / l) (Benzie and Strain 1966).

## RESULTS AND DISCUSSIONS

Metal ion concentrations in nettle leaves vary between wide limits in the different studies (Dudas 2008). Concentrations measured by us generally were in the higher ranges.

Table 1 shows the element compositions in dry leaves of *Urtica dioica* L. The concentrations of Cd, Co, Cr, Hg, Pb and V were under the detection limit in each samples, these elements are not indicated in the table.

The concentration of Al and Ba was significantly the lowest in the sample from the meadow. In teas from conventional and organic collection relatively high concentrations of Al were found.

On the bases of previous studies it can be said that the concentration of Ca shows a high variability. In our investigations the lowest concentration was obtained for the drug from the pharmacy and the highest for the sample collected from along the roadway.

Some differences were found in the concentrations of Fe, nettle leaves from the meadow had the lowest Fe content among the investigated plant materials.

The concentration of K was almost equal, no differences could be detected between the samples.

The Mg content was significantly lower in the leaves from along the roadway than in the other samples.

Mn seems to be accumulated in a high volume in leaves of nettle. The concentration of this element was significantly higher in the obtained samples (except of sample from along the roadway) than it has been reported in the literature.

In the case of Na the two commercial teas had significantly the highest concentrations.

**Table 1:** Element concentration of *Urtica dioica* L. (mg/kg dry weight)

|           | <b>Meadow</b> | <b>Roadway</b> | <b>Pharmacy</b> | <b>Tea</b> | <b>Eco tea</b> |
|-----------|---------------|----------------|-----------------|------------|----------------|
| <b>Al</b> | 2,50          | 48,47          | 84,41           | 158,1      | 138,2          |
| <b>As</b> | 12,09         | 7,56           | 7,56            | 4,43       | 2,91           |
| <b>B</b>  | 37,84         | 36,42          | 48,84           | 46,86      | 46,75          |
| <b>Ba</b> | 16,49         | 45,72          | 35,45           | 34,58      | 42,17          |
| <b>Ca</b> | 32810         | 37690          | 27740           | 35687      | 35411          |
| <b>Cu</b> | 14,02         | 16,69          | 12,99           | 12,14      | 11,84          |
| <b>Fe</b> | 98,35         | 185,10         | 188,70          | 208,54     | 162,52         |
| <b>K</b>  | 23250         | 23750          | 24390           | 24795      | 26207          |
| <b>Li</b> | 1,202         | 1,213          | 1,282           | 0,262      | <dl            |
| <b>Mg</b> | 5231          | 3117           | 5385            | 5108       | 5435           |
| <b>Mn</b> | 116,5         | 33,6           | 224,6           | 172,5      | 177,7          |
| <b>Mo</b> | 2,29          | 4,33           | 1,42            | 4,80       | 2,17           |
| <b>Na</b> | 118,0         | 146,5          | 152,4           | 325,3      | 396,8          |
| <b>Ni</b> | <dl           | 0,53           | 2,34            | 1,42       | 1,08           |
| <b>P</b>  | 4927          | 6229           | 3877            | 5010       | 4648           |
| <b>Se</b> | 26,03         | 25,76          | 27,83           | <dl        | <dl            |
| <b>Si</b> | 140,8         | 146,3          | 137,1           | 425,5      | 427,5          |
| <b>Sr</b> | 73,41         | 125,4          | 109,8           | 107,6      | 116,7          |
| <b>Ti</b> | 1,57          | 3,65           | 2,86            | 2,35       | 1,73           |
| <b>Zn</b> | 25,75         | 27,72          | 22,04           | 27,04      | 25,11          |

<dl: under detection limit

The highest phenol content was measured in aqueous extract of commercial tea (292,07 µg/ml) and the lowest value was found in nettle leaves from the meadow (92,98 µg/ml) (Fig. 1).




**Fig. 1:** Total phenol content of extracts of *Urtica dioica* L. leaves

In the case of ethanolic extracts the highest phenol content could be detected in the sample from along the roadway (130,3 µg/ml). The lowest phenol content had in our investigations the tea from organic collection (44,56 µg/ml).




**Fig. 2.** Antioxidant capacity of extracts of *Urtica dioica* L. leaves

The antioxidant capacity was significantly higher in the aqueous extracts than in the ethanolic extracts (Fig. 2). In the case of aqueous extracts the drug from the pharmacy had the significantly highest antioxidant capacity (16,01 mM AA/l) and the sample from the meadow showed the lowest antioxidant capacity (3,98 mM AA/l).

Well detectable differences could be obtained also for the antioxidants

capacity in the ethanolic extracts: the highest value was found for the drug from the pharmacy (6,95 mM AA/l) and the lowest (1,85 mM AA/l) for the commercial tea from conventional collection.

## CONCLUSIONS

1. *Urtica dioica* L. accumulates some metal elements in excessive amount in its leaves.
2. There are big differences in element concentrations between samples of different origin. The growing place play an important role in accumulation of metal ions.
3. Antioxidant capacity was in a good accordance with the phenol content of the samples. Significant differences could be detected for the phenol content and the antioxidant capacity between the aqueous and ethanolic extracts.
4. There are closed relations between the antioxidant capacity and concentrations of some metal elements, which influence the nutritive value of herbs.
5. Place of collection of wild growing medicinal plants should be chosen cautiously and heavy metal concentration of herbs should be controlled regularly before being marketed.

## REFERENCES

1. Benzie I. I .F., Strain J. J.: The ferric reducing ability of plasma (FRAP) as a measuring of "antioxidant power": The FRAP assay. *Annal. Biochem.*, 1966, 239: 70-76.
2. Blázovics A., Szentmihályi K., Lugasi A., Balázs A., Hagymási K., Bányai É., Then M., Rapavi E., Héthelyi É.: In Vitro Analysis of the Properties of Beiqishen Tea. *Basic Nutritional Investigation. Nutrition*, 2003, 19, 869-875
3. Dudas L.: A nagy csalán (*Urtica dioica* L.) zöldségnövénykénti hasznosításának elméleti és gyakorlati alapjai. Doktori értekezés. Debreceni Egyetem, Agrár- és Műszaki Tudományok Centruma. Debrecen, 2008
4. Neumayer É.: *Urtica dioica* L. – nagy csalán. In: Bernáth, J. (ed.): Gyógy- és aromanövények. Mezőgazda Kiadó, Budapest, 2000
5. Prasad M. N. V., Freitas H. M. O.: Metal hyperaccumulation in plants – Biodiversity prospecting for phytoremediation technology. – *Electronic Journal of Biotechnology* 2003, 6(3): 285–321
6. Sawidis T., Marnasidis A., Zachariasidis G., Stratis J.: A study of air pollution with heavy metals in Thessaloniki city (Greece) using trees as biological indicators. *Arch. Environ. Contam. Toxicol.* 1995, 28:118-124
7. Singh B. R ., Narwal R . P., Almas A. R : Crop uptake and extractability of cadmium in soils naturally high in metals at different pH levels. *Commun. – Soil Sci. Plant Anal.* 1995, 26, 2123–2142.
8. Singleton V. L., Rossi J. A.: Colorimetry of total phenolics with phosphomolibdic-phosphotungstic acid reagents. *Am. J. Enol. Vit.*, 1965, 161: 144-158.
9. Szentmihályi K., Then M.: Study of the constituents on mineral elements and

- biological active substances and their extract in some plants. Olaj Szappan Kozmetika, 1999, 48,173-180
10. Tack F.M., Verloo M.G. 1996. Metal contents in stinging nettle (*Urtica dioica* L) as affected by soil characteristics. Sci. Total Environ. 192:31-39

# STUDY REGARDING THE HEAVY METALS CONTENTS IN SOME VEGETABLES CULTIVATED IN THE FIELD IN ROMANIAN WEST AREA

Aurel LĂZUREANU<sup>1</sup>, Diana MOIGRĂDEAN<sup>2\*</sup>, Constantin MARUTOIU<sup>3</sup>,  
Ioan GOGOASA<sup>2</sup>, Iosif GERGEN<sup>2</sup>

1. Faculty of Horticulture, University of Agricultural Sciences and Veterinary Medicine of Banat, Calea Aradului 119, RO-300645 Timisoara, Roumania; 2 Faculty of Food Products Technology, University of Agricultural Sciences and Veterinary Medicine of Banat, Calea Aradului 119, RO-300645 Timisoara, Roumania; 3 Department of Analytical Chemistry, Faculty of Sciences, University "Lucian Blaga" Str. Ion Rațiu nr. 7-9, RO-550 012, Str. Oituz nr. 2, RO-2400 Sibiu, Roumania

## ABSTRACT

*In this paper was analyzed the some heavy metals (Co, Ni, Pb and Cd) accumulated in some vegetables (pepper, tomatoes, cucumber and eggplants) cultivated in field, in Romanian west area. The specific climatic factors from this areal create conditions for normal uptake of the heavy metals.*

*The experimental results revealed that the distribution of the heavy metals in vegetables from this areal are in the range of normal limits concentrations.*

**Key words:** vegetables, heavy metals

## INTRODUCTION

Vegetables are a source of important vitamins, minerals and carbohydrates. Heavy metals, such as cadmium, copper, lead, chromium and mercury, are important environmental pollutants, particularly in areas with high anthropogenic pressure. Their presence in the atmosphere, soil and water, even in traces, can cause serious problems to all organisms. Heavy metal accumulation in soils is of concern in agricultural production due to the adverse effects on food quality (safety and marketability), crop growth (due to phytotoxicity) (Ma et al., 1994) and environmental health (soil flora/fauna and terrestrial animals). Metal contamination of garden soils may be widespread in urban areas due to past industrial activity and the use of fossil fuels (Sánchez-Camazano et al., 1994).

Gardeners should avoid placing a vegetable garden near an older painted building or on a site where a building once stood. If this cannot be avoided, have the soil tested for lead and other heavy metals and contaminants. Soil also may be contaminated with heavy metals from automobile and equipment exhaust and pesticide use in old orchards. Vegetables growing on contaminated, but otherwise fertile soil, are contaminated mainly by dust on the leaves. Washing vegetables before consuming them negates this potential problem. Some vegetables absorb some of the heavy metals and become contaminated, posing a threat to the health of children, the elderly and ill people who eat them ([www.hcs.ohio-state.edu](http://www.hcs.ohio-state.edu)).

---

„Metal Elements in Environment, Medicine and Biology”, Tome VIII, pp. 195-198, Publishing House „Eurobit” Timișoara, 2008

Proc. of 8<sup>th</sup> Int. Symp. of Roumanian Academy - Branch Cluj-Napoca and Branch Timișoara, 2008, December 5-6, Timișoara, Roumania

**Cobalt (Co)** is a naturally-occurring element that has properties similar to those of iron and nickel. Small amounts of these chemical compounds can be found in rocks, soil, plants, and animals. Cobalt is even found in water in dissolved or ionic form, typically in small amounts. A biochemically important cobalt compound is vitamin B<sub>12</sub> or cyanocobalamin. Vitamin B<sub>12</sub> is essential for good health in animals and humans ([www.atsdr.cdc.gov](http://www.atsdr.cdc.gov)). Cobalt is essential in trace amounts for human life. It is part of vitamin B-12, and plays a key role in the body's synthesis of this essential vitamin. Cobalt has also been used as a treatment for anemia, because it causes red blood cells to be produced. The major source of cobalt is vegetables. The toxicity of cobalt is quite low compared to many other metals in soil. Exposure to very high levels of cobalt can cause health effects. Effects on the lungs, including asthma, pneumonia, and wheezing, have been found in workers who breathed high levels of cobalt in the air (Erdogrul, O, 2006).

**Nickel (Ni)** Small nickel particles in the air settle to the ground or are taken out of the air in rain. Much of the nickel in the environment is found with soil and sediments because nickel attaches to particles that contain iron or manganese, which are often present in soil and sediments (Erdogrul, O, 2006). Nickel is a component of some enzymes, this element playing an important role in all plant species which contain these enzymes (some cucurbitaceous plants) (Barry T., 2006).

**Lead (Pb)** In general, plants do not absorb or accumulate lead. However, in soils testing high in lead, it is possible for some lead to be taken up. Higher concentrations are more likely to be found in leafy vegetables and on the surface of root. Since plants do not take up large quantities of soil lead, the lead levels in soil considered safe for plants will be much higher than soil lead levels where eating of soil is a concern. Generally, it has been considered safe to use garden produce grown in soils with total lead levels less than 300 ppm. The risk of lead poisoning through the food chain increases as the soil lead level rises above this concentration. Even at soil levels above 300 ppm, most of the risk is from lead contaminated soil or dust deposits on the plants rather than from uptake of lead by the plant (Rosen, C., 2002).

**Cadmium** is the most element among all analyzed heavy metals. Cadmium is concentrated in particular parts of plants. As a general rule, leaves contain the most, followed by storage roots and tubers, seeds or grain and fleshy fruits (Murtaza, G., 2008).

## MATERIALS AND METHODS

Were used chemicals and reagents from Merck; deionized water.

Soil samples were taken (0-25 cm depth) before vegetables cultivation.

Heavy metals determination: 20 g fresh vegetables were dried at 105<sup>0</sup>C to 3 hours, calcinated at 650<sup>0</sup>C for 3 hours; added 10 mL HNO<sub>3</sub> 0.5N solution and to run dry. The mineral residue were solubility in 25 mL HNO<sub>3</sub> 0.5N (MAIA, 1983). Determination of heavy metals were determinate by atomic absorption spectroscopy using Continuum Source Atomic Absorption Spectrometer contrAA<sup>®</sup>300 by Analytik Jena. Using standard work conditions - flame type: C<sub>2</sub>H<sub>2</sub>/air; flame height: 6mm; air flow: 568 L/h; acetylene flow: 50 L/h. The wavelength dominate ( $\lambda$ ) for each chemical elements: Cd –  $\lambda$ =228 nm, Co –  $\lambda$ = 240 nm, Ni –  $\lambda$ = 232 nm, Pb –  $\lambda$ = 217 nm.

Were used the work protocol that is stipulated in the AOAC standards.

All determination was repeated for three times.



## RESULTS AND DISSCUSIONS

The experience was done in a cambic cernosium soil, with low acidity reaction and the high natural fertility potential favorable vegetables cultivation. Experimental data of heavy metals content in soil are presented in table 1.

**Table1.** Heavy metals content in soil

| Heavy metals   | Co [mg/kg] | Ni [mg/kg] | Pb [mg/kg] | Cd [mg/kg] |
|----------------|------------|------------|------------|------------|
| Average values | 4.8        | 14.8       | 10.3       | 0          |

In comparison with official limits, the values for heavy metal content in soil are in the normal range for all elements. Cd is not present in soil. Maximum limits accept in Romanian legislation for heavy metals in soil: Pb and Ni-20.0 mg/kg dry matter, Cd - 1.0 mg/kg (Ordinance 756/1997).

In table 2 was presented the metals content in vegetables.

**Table2.** Heavy metals content in some vegetables cultivated in west area.

| Analysed product | Co [mg/kg] | Ni [mg/kg] | Pb [mg/kg] | Cd [mg/kg] |
|------------------|------------|------------|------------|------------|
| Pepper           | 0.05       | 0.22       | 0.02       | 0.02       |
| Tomatoes         | 0.05       | 0.07       | 0.00       | 0.01       |
| Cucumber         | 0.05       | 0.10       | 0.02       | 0.02       |
| Eggplants        | 0.05       | 0.11       | 0.01       | 0.02       |

Maximum limits accept in Romanian legislation for heavy metals in vegetables: Pb-0.5 mg/kg fresh matter, Cd, Co and Ni -0.1 mg/kg (Ordinance 975/1998).

The European Community legislation impose lower values for these limits: 0.1 mg/kg for vegetables (Directive 2001/22/EC).

The cobalt concentration was found in the same values (0.05mg/kg) for these vegetables.

Ni is present only in trace, in the range of 0.07-0.25 mg/kg for vegetales.

Lead is a heavy metal with a higher toxic potential, the maximum admissible limits being 0.5 mg/kg for fresh vegetables. Pb is not present in tomatoes fruit. In the study lead was identified in relatively lower concentrations, which are in the maximum admissible limits from our country, but higher than the limits imposed by the European Community. The most contaminated vegetables with lead is pepper and cucumber (0.02 mg/kg).

In all product samples the concentration of cadmium was very small, these concentrations were in the range from 0.01 to 0.02 mg/kg.

## CONCLUSIONS

The lowest concentrations were determined for lead and cadmium.

Middle concentrations were determined for cobalt, and nickel.

The studied areals do not presents any risk of pollution/contamination with this heavy metals.

## REFERENCES

1. Barry Thomas, John A. Roughan, Evelyn D. Watters, 2006, Cobalt, chromium and nickel content of some vegetable foodstuffs, *Journal of the Science of Food and Agriculture*, Volume 25 Issue 7, Pages 771 – 776
2. Erdogrul, O, Erbilir, F, Distribution of Cobalt, Lead, and Nickel in Various Vegetables from Kahramanmaras, Turkey, 2006 *Bull. Environ. Contam. Toxicol.* (2006) 77:282–288
3. Ma, QY; Traina, SJ; Logan, TJ., 1994, Effect of aqueous Al, Cd, Fe(II), Ni and Zn on Pb immobilization by hydroxyapatite. *Environ Sci Technol.* 1994;28(7):1219–1228.
4. Metodologia de analiza agrochimica a solurilor in vederea stabilirii necesarului de amendamente si de ingrasaminte (MAIA), 1983, vol. I, partea1
5. Murtaza, Ghulam, Ghafoor, Abdul, Qadir, Manzoor, 2008, Accumulation and implications of cadmium, cobalt and manganese in soils and vegetables irrigated with city effluent *Journal of the Science of Food and Agriculture*, Volume 88, Number 1, January 2008 , pp. 100-107(8)
6. Rosen J.Carl, 2002, Lead in Garden Soils and Plants Extension Research Soil Scientist Department of Soil, Water and Climate
7. Sánchez-Camazano, M; Sánchez-Martín, MJ; Lorenzo, LF., 1994, Lead and cadmium in soils and vegetables from urban gardens of Salamanca (Spain). *Sci Total Environ.* 1994;146/147:163–168
8. \*\*\* Romanian legislation, Ordinance 756/1997, The heavy metals concentrations in soil (average values)
9. \*\*\* Romanian legislation, Ordinance 975/1998, Art.97, Maximum limits for arsen and heavy metals in food
10. \*\*\* Directive 2001/22/European Community
11. <http://www.atsdr.cdc.gov/toxprofiles/phs33.html>
12. <http://www.hcs.ohio-state.edu/mg/manual/veg.htm>

# MICROELEMENT CONTENT IN FRUITS OF FIG GENOTYPES GROWN IN HUNGARY AND IN COMMERCIALY AVAILABLE DRIED FIGS

Ádám NAGY <sup>1</sup>, Éva STEFANOVITS-BÁNYAI <sup>1\*</sup>, András NEMÉNYI <sup>2</sup>,  
Attila HEGEDŰS <sup>1</sup>

*1\*.Department of Applied Chemistry, Corvinus University of Budapest, Villányi út 29, H-1118 Budapest, Hungary; 2. Institute Horticulture Technology, Szent István University of Gödöllő, H-2103 Gödöllő, Hungary*

\* E-mail : eva.banyai@uni-corvinus.hu

## ABSTRACT

*Our research was carried out to determine the antioxidant capacity, total phenolics content and nutrient element contents in the fruits of fig (*Ficus carica* L.) grown in Hungary as well as in dried fruits commercially available in the Hungarian market. In this study, the possibility of fig cultivation was investigated. The climate changing of the Carpathian Basin may favour an increase in the cultivation of this species all around this region. Our results show that fig fruits are naturally rich sources of minerals particularly as concerns elements K, Mg, Ca, Na, and P. In addition, we found a high antioxidant capacity, especially in dried figs. This might be due to a relatively gentle way of processing by preserving the heat sensitive compounds and a resultant increase in the dry mass. Our study demonstrated that fig might be considered as a perspective fruit with putative application under special diet requirements by providing high amounts of polyphenols and nutrient elements.*

**Key words:** antioxidants, *Ficus carica*, fig, macro and microelements, polyphenolics

## INTRODUCTION

The Hungarian climate is regarded continental temperate. According to the four seasons, there are huge differences in it: January is the coldest and July is the warmest month. The annual median temperature is approximately 12°C, the maximum is 38 °C, while the minimum is -10 °C. Spring starts in early April with big rainfalls. Summers are dry and hot. Autumns are chilly, misty and wet and winters are moderately cold and usually dry, but sometimes sunny. The annual rainfall is around 550-580 mm. According to the forecasts of scientists, the average temperature of the Carpathian basin is expected to increase by 1.5-8°C in the next 100 years. That kind of climate-changing would be able to make the opportunity to increase the importance of fruits, which had difficulties in the adaptation to previous ecological conditions.

Fig (*Ficus carica* L.) might be one of these plants. Figs have been known in Hungary for centuries as a fruit-bearing plant of upland historical wine-growing districts or as an ornamental and fruit-bearing plant with an exotic appearance cultivated in sheltered gardens (Reszelő, 2005). Because of its climate requirements (temperature and light), it could have been grown safely exclusively on hills in

southern exposition, because its canes might be damaged in winter below an average temperature of  $-15^{\circ}\text{C}$  (Jeszenszky, 1959, 1963). In Hungary, there are some fig clones (local cultivars) which have been adapted to the Hungarian climate. These belong to the group of common or Adriatic type figs. This group has seedless fruits and sets without fertilization.

These local cultivars have been established in Hungary since the Turkish occupation and could have been remained over time, because they can be easily propagated by layering and by softwood or hardwood cuttings (Schmidt and Tóth, 1996).

## MATERIALS AND METHODS

In this study, dried fruits of fig from different southern country (Turkey, Iran, and Syria) and fresh fruits of different Hungarian fig clones were examined. Two of the three fresh pear-shaped figs were characterised by green skin colour and the remainder was purple (Table 1).

**Table 1.** The codes used to label fresh and dried fig samples.

| Fresh figs | Genotype, extraction          | Code          |
|------------|-------------------------------|---------------|
|            | Almási fehér, 70% MeOH        | <b>FFAF A</b> |
|            | Almási fehér, water           | <b>FFAF V</b> |
|            | Almási lila, 70% MeOH         | <b>FFAL A</b> |
|            | Almási lila, water            | <b>FFAL V</b> |
|            | Házi 70%, MeOH                | <b>FFI A</b>  |
|            | Házi, water                   | <b>FFI V</b>  |
|            |                               |               |
| Dried figs | Country of origin, extraction | Code          |
|            | Assyria, 70% MeOH             | <b>AFA A</b>  |
|            | Assyria, water                | <b>AFA V</b>  |
|            | Iran, 70% MeOH                | <b>AFI A</b>  |
|            | Iran, water                   | <b>AFI V</b>  |
|            | Hungary, 70% MeOH             | <b>AFM A</b>  |
|            | Hungary, water                | <b>AFM V</b>  |
|            | Turkish, 70% MeOH             | <b>AFT V</b>  |
|            | Turkish, water                | <b>AFT A</b>  |

The element contents were determined from freeze-dried samples with ICP-OES IRIS Thermo Jarrel ASH instrument.

Total soluble phenols were determined using Folin-Ciocalteu reagent according to the method of Singleton and Rossi (1965). The content of soluble phenols was calculated from a standard curve obtained with different concentrations of gallic acid.

Antioxidant power was determined by the FRAP (Ferric Reducing Ability of Plasma) method at  $\lambda=593\text{ nm}$  (Benzie and Strain, 1996). Deionized water or 70% (V/V) methanolic extraction was performed from the homogenized samples. After centrifugation ( $4^{\circ}\text{C}$ , 35-min,  $18,750\text{ g}_n$ ) clear supernatants were utilized for redox analyses.

Dry matter content was measured to facilitate comparison between dried and fresh fruits.

## RESULTS AND DISCUSSIONS

The results of the experiments carried out on fresh and dried fig samples are presented with their codes listed in Table 1. Table 2. shows the dry matter content of each sample in percentage. According to these data, dry matter contents of dried figs are approximately 4-4.5 times higher than those of fresh fruits. Thanks to the low

water and high sugar contents due to water evaporation, dried figs can last long, and hence this technique have been utilized for thousand years.

**Table 2.** Dry matter content of fresh and dried fig samples.

| Dry matter content % | Fresh fig |       |       |  | Dried fig |       |       |       |
|----------------------|-----------|-------|-------|--|-----------|-------|-------|-------|
|                      | FFAL      | FFAF  | FFI   |  | AFA       | AFI   | AFM   | AFT   |
|                      | 23,28     | 34,01 | 19,34 |  | 96,65     | 88,31 | 90,32 | 96,29 |

Among the mineral elements, those having a value below the detection limit were not included (Table 3). Huge differences in element contents in fruits originating from Hungary or the Middle East may have several explanations. The influencing effects of different soil characteristics and a wide range of biochemical factors may be involved as it was previously shown (Stefanovits-Bányai et al., 2005, 2006; Hegedűs et al., 2008). In addition, discrepancies in genetic constitution may be also notable since genetic diversity in the Middle East region was shown to be high for several fruit species (Ercisli, 2004; Halász, 2007a,b; Ercisli et al., 2008).

**Table 3.** Nutrient element content of fresh and dried figs (µg/g).

| Elements<br>µg/g | Apple     | Fresh fig |        |        |  | Dried fig |        |       |       |
|------------------|-----------|-----------|--------|--------|--|-----------|--------|-------|-------|
|                  | (avarage) | FFAL      | FFAF   | FFI    |  | AFA       | AFI    | AFM   | AFT   |
| Al               | 5,985     | 19,91     | 10,34  | 12,48  |  | 18,07     | 15,1   | 17,68 | 24,66 |
| B                | 21        | 7,099     | 9,194  | 11,68  |  | 17,42     | 20,63  | 11,81 | 15,05 |
| Ba               | 1,8       | 9,193     | 0,8021 | 1,736  |  | 5,789     | 4,238  | 7,071 | 9,806 |
| Ca               | 58        | 1473      | 1657   | 3100   |  | 2315      | 3553   | 1979  | 3254  |
| Cu               | 12,1      | 6,916     | 6,55   | 6,557  |  | 9,164     | 7,725  | 4,581 | 11,09 |
| Fe               | 2,5       | 18,64     | 25,68  | 21,18  |  | 23,36     | 18,64  | 25,22 | 26,51 |
| Ga               | x         | x         | 2,051  | 0,8205 |  | 9,22      | 0,6154 | x     | 4,965 |
| K                | 1220      | 11977     | 10943  | 9554   |  | 11223     | 8725   | 7729  | 10888 |
| Mg               | 403       | 536       | 837,1  | 1055   |  | 892,5     | 1021   | 675,8 | 1138  |
| Mn               | x         | 1,393     | 1,393  | 3,018  |  | 4,45      | 4,992  | 5,815 | 6,164 |
| Na               | 12        | 99,76     | 127,2  | 123,2  |  | 1447      | 606,8  | 411,7 | 2828  |
| Ni               | x         | 1,384     | 1,109  | 1,274  |  | 2,171     | 1,274  | 2,153 | 2,195 |
| P                | 120       | 987,5     | 1114   | 970,7  |  | 1110      | 692,1  | 618,3 | 1103  |
| Si               | x         | 74,49     | 72,11  | 74,43  |  | 87,3      | 98,08  | 86,9  | 100,8 |
| Sr               | x         | 4,05      | 4,445  | 9,024  |  | 9,553     | 49,83  | 10,5  | 14,99 |
| Zn               | 6,1       | 20,14     | 10,85  | 9,552  |  | 12,97     | 8,152  | 5,275 | 13,85 |

Compared with apples, figs generally contain significantly higher amounts from Ca, Fe, K, Mg, and Zn. The dried fig samples originating from the Middle East have extremely high contents from such elements.

RDA% (Recommended Daily Allowance) were calculated for microelements of fresh and dried figs compared with a common apple and these data are represented on Table 4.( Belitz et al., 2005).

**Table 4.:** RDA% of microelements of the fresh and dried figs

| Elements  |              | Al     | Ca    | Cu    | Fe    | K     | Mg    | Mn    | Na   | Ni     | P    | Si    | Zn   |
|-----------|--------------|--------|-------|-------|-------|-------|-------|-------|------|--------|------|-------|------|
| Fresh fig | RDA% / 100mg | 36,39  | 5,31  | 13,64 | 3,72  | 7,00  | 5,17  | 1,41  | 0,10 | 116,66 | 2,62 | 5,62  | 2,76 |
| Dried fig |              | 175,37 | 25,78 | 60,50 | 14,51 | 22,68 | 21,64 | 14,21 | 4,24 | 658,15 | 8,18 | 25,87 | 7,48 |
| Apple     |              | 7,78   | 0,075 | 12,58 | 0,22  | 0,40  | 1,31  |       | 0,01 |        | 0,16 |       | 0,63 |

In case of both total antioxidant capacity and total soluble polyphenol content, big differences were detected between the fresh and dried fig samples. Moreover, dried figs showed 4-4.5 times higher values than fresh fruits (Fig. 1., Fig. 2.).

However, the differences among dried figs may come from the different drying technologies used and from the distinct features of specific plant genotypes. The correlation between total soluble polyphenolics content and FRAP values seems to be close ( $R=0.89284$ ), so it might be supposed that in case of figs total soluble polyphenolics content has the most decisive contribution to the total antioxidant capacity.




**Fig. 1.** The total antioxidant capacity of water and methanolic extracion from fresh and dried figs (mg AS/100g).




**Fig. 2.** The total polyphenolics content of water and methanolic extracion from fresh and dried figs (mg GS/100g).

## CONCLUSIONS

This study demonstrated that fig might be considered as a perspective fruit with putative application under special diet requirements since it provides high amounts of polyphenols and nutrient elements. Combining the results for ICP and redox analyses, we can conclude that higher element contents of dried figs may not only be attributed to the different soil conditions in foreign countries, but also a putative higher chelating capacity in these fruit samples due to their high polyphenolic contents. Even if flavonoids are devoted to metal chelation they retain their radical scavenging capacity (Lugasi et al., 2003) and can have a crucial contribution to the antioxidant capacity of fruits.

## REFERENCES

1. Benzie I.F., Strain J.J. (1996): The ferric reducing ability of plasma (FRAP) as a measuring of "antioxidant power": The FRAP assay. *Anal. Biochem.* 239: 70–76.
2. Ercisli S. (2004): A short review of the fruit germplasm resources of Turkey. *Genet. Resour. Crop Ev.*, 51: 419–435.
3. Ercisli S., Akbulut M., Ozdemir O. (2008): Phenolic and antioxidant diversity among persimmon (*Diospyrus kaki* L.) genotypes in Turkey. *Int. J. Food Sci. Nutr.*, 59(6):477–482.
4. Belitz H.-D., Grosch W., Schieberle P. (2005): *Food Chemistry* 3rd revised edition, Springer.
5. Halász J. (2007a): Review of self-(in)compatibility in apricot (*Prunus armeniaca* L.). *World J. Agric. Sci.*, 3(5): 670–677.
6. Halász J. (2007b) A kajszai önmeddőségét meghatározó S-allél-rendszer molekuláris háttere. PhD thesis. Budapest, Budapesti Corvinus Egyetem
7. Hegedűs A., Balogh E., Engel R., Sipos B.Z., Papp J., Blázovics A., Stefanovits-Bányai, É. (2008): Comparative nutrient element and antioxidant characterization of berry fruit species and cultivars grown in Hungary. *HortScience*, 43(6): 1711–1715.
8. Jeszenszky Á., Kárpáti I. (1963): A füge (*Ficus carica* L.). Magyarország kultúrflórája, VII. kötet, 12. füzet, Akadémiai Kiadó, Budapest, 68 o.
9. Lugasi A., J. Hóvári K.V. Sági, L. Bíró. (2003): The role of antioxidant phytonutrients in the prevention of diseases. *Acta Biol. Szeged.* 47:119–125.
10. Reszelő D. (2005): A füge és hasznosítása, diplomamunka, SZIE, Gödöllő.
11. Schmidt G., Tóth I. (ed.)(1996): *Díszfaiskola*. Mezőgazda Kiadó, Budapest, 669.
12. Singleton V.L., Rossi J.A. (1965): Colorimetry of total phenolics with phosphomolybdic phosphotungstic acid „reagents”. *Am. J. Enol. Vitic.* 16: 144–158.
13. Stefanovits-Bányai É., Engel R., Hermán R., Blázovics A., Hegedűs A. (2005): Antioxidant characterization of apricot (*Prunus armeniaca* L.) cultivars and hybrids. *International Journal of Horticultural Science*, 11(4): 47–51.

14. Stefanovits-Bányai É., Szentmihályi K., Hegedűs A., Koczka N., Váli L. Taba G., Blázovics A. (2006): Metal ion and antioxidant alterations in leaves between different sexes of *Ginkgo biloba* L. Life Sci., 78(10): 1049–1056.



# BODY ZINC LEVEL, EATING HABITS AND SOME NUTRITIONAL PARAMETERS IN AN ELDERLY GROUP

Victoria PAPADOPOLO\*, Elena POPA, Gabriela ALBU, Cristina CÎMPEANU,  
Nicoleta FLORESCU, Iliana PALAMARU

*Department of Food Hygiene and Nutrition, Institute of Public Health, 14 Victor Babes, RO-700465 Iasi, Roumania*

\* E-mail : vicpapadopol@yahoo.com

## ABSTRACT

*Elderly people is an important population group affected by the zinc deficiency. One of the cause of this deficiency may be the nutrition. The aim of the study was to search the relation between the zinc level and some eating habits and between the zinc level and some nutritional parameters in a group of elderly (60 – 85 years) comparing with a group of adults (20 – 59 years). We found in all subjects an insufficient energy intake and some nutritional imbalance. In elderly serum zinc positively correlated with ingested energy and negatively with ingested animal fats. The frequency of food consumption generally did not correlate with zinc level. Investigated biochemical parameters correlated with serum or erythrocyte zinc level both in older and younger subjects, emphasizing the implication of the metal in protein, lipid and glucose metabolism.*

**Key words:** zinc, elderly, eating habits, biochemical parameters

## INTRODUCTION

Zinc plays an essential role in living bodies, being indispensable for cellular functions. Zinc deficiency affects at least 1 in 5 people worldwide (Maret & Sandstead, 2006) and among the implicated population groups elderly people are an important one. It is assumed that zinc deficit gradually occurs; its bioavailability for intracellular enzymes decreases, and the ageing process is initiated. It was suggested that intracellular zinc deficiency may be the first biochemical event that starts the whole spectrum of functional changes that lead to the loss of division capacity and to the cellular death in cells culture in vitro (Hayflick, 1982). There are many causes of zinc deficit in elderly, one of them being nutrition. Today, the elderly tend to avoid meat and other high zinc content food due to fears of cholesterol. Rather, they increase consumption of refined wheat products. Much more important than the quantity of ingested zinc are factors that condition zinc absorption. The proportion of the absorbed zinc may vary from 20 to 80 % of the ingested zinc. Phytic acid, vegetal fibres, calcium and copper decrease zinc absorption, while animal proteins, picolinic acid, some amino acids and peptides increase zinc absorption (Bulboacă, 2005).

The aim of the study was to search the relation between the zinc level and some eating habits and between the zinc level and some nutritional parameters in a group of elderly comparing with a group of adults.

## MATERIALS AND METHODS

Study was performed in a group of persons selected from the consulting rooms of two general practitioner physicians from the Iasi city. Initially, the elderly group (60 - 85 years) comprised 139 subjects that were compared with a group of 104 younger subjects (20 – 59 years). After removing the subjects suffering of diseases (stroke, allergy) and taking medicines (diuretics) that showed associations with the zinc level, the elderly group had 65 subjects and the younger group had 60 subjects.

The body zinc level was determined by assessing serum, erythrocyte and urine zinc using atomic absorption spectrophotometry. Serum glucose, total proteins, albumin, total cholesterol, HDL-cholesterol, triglycerides were determined. LDL-cholesterol was calculated. Eating habits were investigated by applying a questionnaire including weekly frequency of food consumption (14 items) and 24h recall of food intake. Total energy and the main nutrients were estimated. Anthropometric variable (weight and high) were registered and Body Mass Index (BMI) was calculated. Statistical analysis was carried out using Epi Info programme (Pearson correlation index, one way ANOVA test, "t" test).

## RESULTS AND DISCUSSIONS

Some characteristics of the elderly were compared with those of the adults (Table 1).

**Table 1.** Characteristics of the study sample

| Variable                           | ≥ 60 years (n = 65) |               | < 60 years (n = 60) |               |
|------------------------------------|---------------------|---------------|---------------------|---------------|
|                                    | %                   | Means +/- SD  | %                   | Means +/- SD  |
| Age (years)                        |                     | 69.4 +/- 6.4  |                     | 49.0 +/- 9.1  |
| Weight (kg)                        |                     | 74.4 +/- 13.4 |                     | 80.4 +/- 21.2 |
| BMI (kg/m <sup>2</sup> )           |                     | 28.6 +/- 5.0  |                     | 28.6 +/- 6.3  |
| Income (low, medium, high)         | 45, 45, 10          |               | 39, 44, 17          |               |
| Education (low, medium, high)      | 43, 28, 29          |               | 17, 43, 40          |               |
| Living alone                       | 20                  |               | 2                   |               |
| Smoking (never, previous, current) | 65, 24, 11          |               | 60, 13, 27          |               |

Table 2 shows the zinc levels of the two groups. One can see a significant difference between sex in elderly regarding serum and erythrocyte zinc. Subjects ≥ 60 years had serum and erythrocyte zinc levels nonsignificantly lower.

Table 3 shows an ingested energy under the recommended values (≥ 2100 kcal for adults in Romania) (Ionuț, 2004), a carbohydrate intake closed to low limit (50 – 60 % recommended) and a total lipid intake over the upper limit (24 – 34 % recommended) for the younger women. One can see a decrease of energy intake in

old women, reflected especially by the decreasing of the proportion of ingested total lipids and the amount (g) of carbohydrates.

**Table 2.** Mean values of zinc

| Specification               |       | >= 60 years                                       |                           | < 60 years    |               |
|-----------------------------|-------|---------------------------------------------------|---------------------------|---------------|---------------|
|                             |       | Women                                             | Men                       | Women         | Men           |
| Serum zinc                  | µg/dl | 75.3 +/- 12.6<br>a: <b>p&lt;0.05</b><br>b: p>0.05 | 86.3+/- 13.7<br>c: p>0.05 | 80.3 +/- 16.7 | 92.3 +/- 16.3 |
| Erythrocyte zinc            | µg/ml | 11.7 +/- 1.4<br>a: <b>p&lt;0.05</b><br>b: p>0.05  | 13.6 +/- 1.1<br>c: p>0.05 | 12.3 +/- 2.0  | 14.0 +/- 2.1  |
| Urinary zinc                | µg/l  | 544 +/- 425<br>a: p>0.05<br>b: p>0.05             | 701 +/- 561<br>c: p>0.05  | 472 +/- 287   | 842 +/- 650   |
| Urinary zinc/<br>creatinine | µg/g  | 687 +/- 599<br>a: p>0.05<br>b: p>0.05             | 562 +/- 420<br>c: p>0.05  | 469 +/- 286   | 561 +/- 346   |

a: comparing with the older men; b: comparing with the younger women; c: comparing with the younger men

**Table 3.** Estimated daily energy and nutrients intake

| Specification   |                       | >= 60 years    |                 | < 60 years     |                 |
|-----------------|-----------------------|----------------|-----------------|----------------|-----------------|
|                 |                       | Women          | Men             | Women          | Men             |
| Energy          | kcal                  | 1545 +/- 474   | 2065 +/- 697    | 1836 +/- 730   | 1923 +/- 749    |
| Total proteins  | g                     | 60.5 +/- 25.0  | 69.9 +/- 29.7   | 62.4 +/- 33.7  | 65.9 +/- 34.0   |
|                 | % from energy         | 15.9 +/- 4.1   | 13.9 +/- 3.6    | 13.8 +/- 4.3   | 14.0 +/- 3.7    |
| Animal proteins | g                     | 35.0 +/- 25.9  | 34.7 +/- 28.0   | 33.4 +/- 25.8  | 35.5 +/- 27.8   |
|                 | % from total proteins | 50.0 +/- 30.7  | 45.0 +/- 26.9   | 47.7 +/- 22.8  | 49.1 +/- 18.9   |
| Total lipids    | g                     | 57.0 +/- 30.5  | 81.1 +/- 40.4   | 82.0 +/- 66.7  | 68.0 +/- 38.7   |
|                 | % from energy         | 32.9 +/- 10.5  | 35.7 +/- 10.8   | 42.2 +/- 39.3  | 31.8 +/- 11.6   |
| Animal lipids   | g                     | 26.3 +/- 22.8  | 30.8 +/- 27.3   | 31.8 +/- 24.9  | 31.6 +/- 23.7   |
|                 | % from total lipids   | 44.1 +/- 31.1  | 36.3 +/- 28.2   | 42.1 +/- 24.5  | 49.2 +/- 27.3   |
| Carbohydrates   | g                     | 186.8 +/- 61.4 | 252.1 +/- 102.0 | 220.8 +/- 88.2 | 247.6 +/- 100.3 |
|                 | % from energy         | 51.0 +/- 12.6  | 50.2 +/- 11.2   | 50.4 +/- 11.3  | 54.2 +/- 13.4   |

Table 4 shows the frequency consumption of food. Older women ate less meat and vegetables than the younger ones and than the men. Younger men ate less dairy products and sweets than older men and than the women. The men consumed fewer eggs than women and the elderly people less than the young people. The old women ate more fish, pasta and pastry products comparing with the other groups. Older men eat less dry legumes than women and than younger people. Older women consumed fewer vegetables than the younger women and than the men. The consumption of vegetal fats and potatoes was comparable in all the groups. The consumption of animal fats was much higher in young men comparing with the other

groups. The women drank alcohol beverages less than men and the older people less than younger people. The women drank coffee more than men and the younger people more than the older ones. One can note generally a reduced consumption of fish and vegetables and a high consumption of animal fats and coffee.

**Table 4.** Frequency of food consumption for a week (%)

| Specification          | Frequency   | >= 60 years |     | <60 years |     |
|------------------------|-------------|-------------|-----|-----------|-----|
|                        |             | Women       | Men | Women     | Men |
| Meat                   | <= once     | 24          | 18  | 15        | 16  |
|                        | 2 – 3 times | 38          | 25  | 22        | 32  |
|                        | >= 4 times  | 38          | 57  | 63        | 53  |
| Dairy products         | <= once     | 18          | 11  | 25        | 42  |
|                        | 2 – 3 times | 38          | 36  | 23        | 16  |
|                        | >= 4 times  | 54          | 54  | 52        | 42  |
| Eggs                   | <= once     | 51          | 64  | 40        | 47  |
|                        | 2 – 3 times | 35          | 18  | 50        | 16  |
|                        | >= 4 times  | 14          | 18  | 10        | 37  |
| Fish                   | <= once     | 78          | 86  | 87        | 89  |
|                        | 2 – 3 times | 22          | 14  | 13        | 11  |
| Potatoes               | <= once     | 30          | 10  | 25        | 26  |
|                        | 2 – 3 times | 40          | 54  | 48        | 42  |
|                        | >= 4 times  | 30          | 36  | 27        | 32  |
| Dry vegetables         | <= once     | 78          | 89  | 85        | 79  |
|                        | 2 – 3 times | 22          | 11  | 15        | 21  |
| Vegetables             | <= once     | 30          | 11  | 22        | 11  |
|                        | 2 – 3 times | 22          | 11  | 5         | 21  |
|                        | >= 4 times  | 48          | 78  | 73        | 69  |
| Fruits                 | <= once     | 13          | 7   | 17        | 16  |
|                        | 2 – 3 times | 14          | 7   | 10        | 21  |
|                        | >= 4 times  | 73          | 86  | 73        | 63  |
| Sweets                 | <= once     | 59          | 56  | 48        | 68  |
|                        | 2 – 3 times | 16          | 21  | 12        | 16  |
|                        | >= 4 times  | 24          | 32  | 40        | 16  |
| Pasta, pastry products | <= once     | 13          | 68  | 60        | 58  |
|                        | 2 – 3 times | 65          | 25  | 28        | 37  |
|                        | >= 4 times  | 22          | 7   | 12        | 5   |
| Vegetal fats           | <= once     | 78          | 64  | 70        | 74  |
|                        | 2 – 3 times | 13          | 18  | 20        | 21  |
|                        | >= 4 times  | 8           | 18  | 10        | 5   |
| Animal fats            | <= once     | 68          | 57  | 57        | 16  |
|                        | 2 – 3 times | 24          | 32  | 28        | 63  |
|                        | >= 4 times  | 8           | 11  | 15        | 21  |
| Alcohol beverages      | <= once     | 84          | 65  | 70        | 58  |
|                        | >=2 times   | 16          | 36  | 30        | 42  |
| Coffee                 | <= once     | 35          | 50  | 22        | 42  |
|                        | >=2 times   | 65          | 50  | 78        | 58  |

Total proteins and albumin values did not significantly vary between sex and groups (Table 5). HDL-cholesterol slightly decreased with age being lower in men. Triglycerides and glucose were slightly increased in elderly.

**Table 5.** Mean values of biochemical parameters

| Specification     | Years $\geq 60$ |            | Years $< 60$ |            |
|-------------------|-----------------|------------|--------------|------------|
|                   | Women           | Men        | Women        | Men        |
| Glucose           | 90 +/- 31       | 92 +/- 26  | 86 +/- 25    | 91 +/- 38  |
| Total proteins    | 83 +/- 4        | 85 +/- 4   | 84 +/- 4     | 85 +/- 4   |
| Albumin           | 48 +/- 5        | 48 +/- 5   | 48 +/- 5     | 48 +/- 6   |
| Total cholesterol | 232 +/- 33      | 224 +/- 37 | 228 +/- 32   | 230 +/- 38 |
| HDL -cholesterol  | 55 +/- 13       | 50 +/- 12  | 58 +/- 11    | 55 +/- 12  |
| LDL-cholesterol   | 150 +/- 34      | 144 +/- 39 | 147 +/- 32   | 147 +/- 42 |
| Triglycerides     | 136 +/- 46      | 148 +/- 54 | 124 +/- 52   | 140 +/- 43 |

The following of the relationships between zinc and ingested nutrients in elderly, showed a positive correlation between serum zinc and ingested energy, a negative correlation between serum zinc and animal lipids (% from total ingested lipids) and a positive correlation between urinary zinc/creatinine and animal proteins (%) (Table 6). A similar tendency of correlation urinary zinc/creatinine - animal proteins was emphasized in the younger subjects.

**Table 6.** Correlations between zinc level and ingested nutrients

| Age (years) |                         |                    | n  | r      | p            |
|-------------|-------------------------|--------------------|----|--------|--------------|
| $\geq 60$   | Serum zinc              | Energy             | 64 | + 0.23 | <b>0.017</b> |
|             | Serum zinc              | Animal lipids (%)  | 62 | - 0.29 | <b>0.024</b> |
|             | Urinary zinc/creatinine | Animal protein (%) | 51 | + 0.28 | <b>0.047</b> |
| $< 60$      | Urinary zinc/creatinine | Animal protein (%) | 46 | + 0.28 | 0.057        |

Serum and erythrocyte zinc did not associated with the frequencies of food consumption probably because of many factors that determine zinc status. Only urinary zinc/creatinine positively associated in elderly with eggs consumption ( $p=0.040$ ).

Serum zinc level positively correlated with serum total proteins and LDL-cholesterol and negatively with HDL-cholesterol in elderly (Table 7). Also, erythrocyte zinc level positively correlated with total proteins and albumin and urinary zinc/creatinine positively correlated with glucose.

In younger subjects, we found a positive correlation between serum zinc and total proteins, between erythrocyte zinc and total proteins and triglycerides and a negative correlation with HDL-cholesterol. Urinary zinc/creatinine positively correlated with glucose and negatively correlated with total cholesterol and LDL-cholesterol.

Our results are in agreement with those of Hermann et al. (1993) who found that both plasma total cholesterol and LDL-cholesterol were elevated with lower zinc intakes. Recently Mazzatti et al. (2007) found that many genes involved in lipid and cholesterol homeostasis are differentially regulated by zinc in an age-dependent manner.

**Table 7.** Relationships between zinc level and biochemical parameters

| Specification           | Parameter         | ≥ 60 years |        |              | < 60 years |        |              |
|-------------------------|-------------------|------------|--------|--------------|------------|--------|--------------|
|                         |                   | n          | r      | P            | n          | r      | P            |
| Serum zinc              | Total proteins    | 65         | + 0.35 | <b>0.005</b> | 59         | + 0.25 | 0.055        |
|                         | HDL-Cholesterol   | 65         | - 0.39 | <b>0.000</b> |            |        |              |
|                         | LDL-Cholesterol   | 65         | + 0.26 | <b>0.036</b> |            |        |              |
| Erythrocyte zinc        | Total proteins    | 63         | + 0.44 | <b>0.000</b> | 58         | + 0.30 | <b>0.022</b> |
|                         | Albumin           | 60         | + 0.34 | <b>0.009</b> |            |        |              |
|                         | HDL-Cholesterol   |            |        |              | 53         | - 0.30 | <b>0.030</b> |
|                         | Triglycerides     |            |        |              | 59         | + 0.37 | <b>0.005</b> |
| Urinary zinc/creatinine | Glucose           | 53         | + 0.26 | 0.061        | 49         | + 0.48 | <b>0.000</b> |
|                         | Total cholesterol |            |        |              | 49         | - 0.29 | <b>0.045</b> |
|                         | LDL cholesterol   |            |        |              | 45         | - 0.29 | 0.055        |

## CONCLUSIONS

1. We found in all subjects an insufficient energy intake and some nutritional imbalance.
2. Serum zinc positively correlated with ingested energy and negatively with ingested animal fats.
3. The frequency of food consumption generally did not correlate with zinc level.
4. Investigated biochemical parameters correlated with serum or erythrocyte zinc level both in older and younger subjects, emphasizing the implication of the metal in protein, lipid and glucose metabolism.

## REFERENCES

1. Bulboaca A., Rolul zincului intracelular în senescență, Clujul medical, 2005, vol LXXVIII, 2: 281 - 7
2. Hayflick L. Biological aspects of human aging, British J of Hospital Medicine, 1982, 27: 366-75
3. Herman J.R., Arquitt A., Hanson C.: Relationships between dietary minerals and plasma lipids and glucose among older adults, J. Nutr. Elderly, 1993, 12 (3)
4. Ionuț C. (sub redacția) :Compendiu de igienă, Edit.Med. Univ. „Iuliu Hațieganu” Cluj-Napoca, 2004
5. Maret W., Sandstead H.H., Zinc requirements and the risks and benefits of zinc supplementation, J Trace Elem. Med. Biol., 2006, 20: 3-18
6. Mazzatti D. J., Mocchegiani E., Powell J.R.: Age-specific modulation of genes involved in lipid and cholesterol homeostasis by dietary zinc, Rejuvenation Res., 2007, Dec.26

# EFFECT OF FOLIAR MAGNESIUM FERTILIZATION ON LEAF AND FRUIT MINERAL COMPOSITIONS OF GREENHOUSE GROWN CUCUMBERS

Katalin SLEZÁK<sup>1</sup>, Sandor KISS<sup>1</sup>, Uzuni Hanna NÉMETHY<sup>1</sup>, Ferenc OROSZ<sup>1</sup>,  
Éva STEFANOVITS-BÁNYAI<sup>2</sup>

*1\* Department of Vegetable and Mushroom Growing, Faculty of Horticulture, Corvinus University of Budapest, Villányi út 29-43., H-1118 Budapest, Hungary; 2 Department of Applied Chemistry, Faculty of Food Science, Corvinus University of Budapest, Villányi út 29-43., H-1118 Budapest, Hungary*

\* E-mail : katalin.slezak@uni-corvinus.hu

## ABSTRACT

*In our experiment we studied the effect of foliar magnesium application (Hydromag, in concentrations of 0.2 and 0.4%, a total of 300 ml solution per plant) on leaf and fruit mineral composition with cucumber plants (*Cucumis sativus* cv. Mustang) grown in peat in containers under greenhouse conditions. Foliar applications were carried out 5 times, 10 days apart. Plants receiving no foliar treatment were also grown as control. For the purpose of laboratory analysis fruit and leaf samples were collected twice. Fruit mineral composition was determined using the apparatus ICP-OES IRIS Thermo Jarrel ASH, Corp., Franklin, MA, USA. The Mg content of the fresh cucumber fruits did not show any increase in response to the foliar treatments. At the first occasion of measurement, the Mg content of the leaves of the treatments with foliar magnesium application were higher than the value registered for the control, but on the second occasion we obtained results that were just the opposite. The foliar Mg fertilization changed the macro- and microelements concentration of leaves in cucumber fruits.*

**Key words:** cucumber, magnesium, mineral composition, foliar fertilization

## INTRODUCTION

Peat is one of the most commonly used root media in the ever more popular container vegetable growing in greenhouse. In the case of this growing method roots have a much smaller volume available as compared to traditional growing in real soil. Therefore a fertilization with a daily frequency is applied using complex chemical fertilizers (Göhler – Drews, 1990). On the other hand, based on the experiences from production, it can be seen that yields and quality can be improved further by foliar fertilization and this way the harmful effect of the nutrient antagonisms occurring in the root medium can also be eliminated.

40-50% of the magnesium absorbed by plants is assimilated by the foliage (Terbe, 1989). The beneficial effects of the application of magnesium as a foliar fertiliser have been demonstrated for several plants (saffron: Takácsné Hájos et al., 2001; vine: Moretti, 2002; tomato: Takács Hájos et al, 2003; pear: Gastol - Domagala-Swiatkiewicz, 2006). Magnesium applied in the form of foliar fertilizer will remain in the leaves and will not move together with plant saps. The young leaves of

a healthy cucumber plant contain 0.5-0.8% magnesium, while the elder ones 0.8-2.0%. The efficiency of the fertilizer spraying is highly dependent on the time of application. In general, the best result can be achieved during the period of intensive development (Fazekas et al., 1992).

The importance of magnesium is manifested in the first place in protein synthesis and photosynthesis, as well as in the enzyme reactions of starch formation (Marschner, 1986).

Occurrence of deficiency symptoms can be expected under 0.3-0.4%. As a result of an inadequate supply, magnesium will be mobilized from the older plant organs to the young ones, therefore the chlorosis of older leaves is observable first (Terbe, 1992; Ushijima et al., 2004).

## MATERIALS AND METHODS

The experiment was set up at the Experimental Farm of the Faculty of Horticulture, Corvinus University of Budapest in a plastic tunnel greenhouse of Filclair type, under unheated conditions.

The following 3 treatments were applied in 4 repetitions (5 plants/plot):

Mg1 = 0.4 g/l Mg as foliar fertilizer

Mg2 = 0.8 g/l Mg as foliar fertilizer

Control = no treatment (receiving only fertigation)

For the purpose of the foliar fertilizer we used the foliar fertilizer product Hydromag (Phosyn fertilizer family) (Mg concentration: 200 g/l) and in the determination of the concentration we followed the manufacturer's recommendation for the bottom and top limits (0.2 and 0.4%).

The variety Mustang (*Cucumis sativus* L. cv. Mustang) was chosen as test plant.

Growing was carried out in black buckets of 12 litres having rigid walls in which 10 litre medium was filled, respectively. For the purpose of root medium we used a 1:1 volume mixture of fen and raised bog peat which was given 1.5 kg/m<sup>3</sup> fodder lime in order to set the pH value.

Seeds were sown in 9 cm pots on March 3, in a mixture of peat and sand. Seedlings were transplanted on April 7. 1.95 plants per square meter were arranged at a spacing of 160 x 32 cm. Shoots were trained on a string trellis up to the horizontal wires stretched at 240 cm high. Plants were pruned to the main stem.

After transplanting plants received nutrient solution on a regular basis according to Table 1. The nutrient solution was applied always in the morning hours and if it was necessary some irrigation was done in the afternoon hours using either plain water or nutrient solution and taking care not to leach the nutrient solution out of the medium.



**Table 1:** Frequencies and concentrations of drip fertigation

| Period              | Frequency      | Amount of Nutrient Solution (l/day/plant) | Fertiliser composition (g/1000l) |                               |                  |    |    |
|---------------------|----------------|-------------------------------------------|----------------------------------|-------------------------------|------------------|----|----|
|                     |                |                                           | N                                | P <sub>2</sub> O <sub>5</sub> | K <sub>2</sub> O | Mg | Ca |
| April 7-8*          | Daily          | 0.5                                       | 300                              | 600                           | 300              | -  | -  |
| April 11-19*        | Every two days | 1                                         | 433                              | 410                           | 260              | 10 | 47 |
| April 20 – June 8** | Daily          | 1                                         | 253                              | 105                           | 315              | 10 | 47 |
| June 9 – July 7*    | Daily          | 1-1.5                                     | 443                              | 245                           | 280              | 10 | 47 |

\* 1,000 litre nutrient solution contained also: Fe: 2.6g; Mn: 0.6g; Zn: 0.4g; Cu:0.26g; B: 0.06g; Mo: 0.04g active ingredient.

\*\* 1,000 litre nutrient solution contained also: Fe: 1.95g; Mn: 0.45g; Zn: 0.3g; Cu:0.195g; B: 0.045g; Mo: 0.03g active ingredient.

Foliar fertilization was applied at every 10-14 days, 5 times in all (for the first time on May 13 and for the last on July 1). Using a hand pump sprayer the nutrient solution was applied evenly to every plant leaf in an amount to provide a thin uniform coating over them. The amount applied in this way over the five occasions was a total of 300 ml/plant, i.e. 585 ml/m<sup>2</sup>.

For the purpose of the laboratory analysis, leaf and fruit samples were collected twice (May 23 and June 27). The leaves came from the middle region of the plant. The mineral composition of the fruits, after their washing, cutting, homogenisation in a blender and drying and pulverization, was determined using an ICP-OES IRIS Thermo Jarrel ASH, Corp., Franklin, MA, USA. At the same time, we also determined the dry matter content of the homogenized pulp, this way we were able to perform calculations to convert the data registered in the dry samples to fresh weight.

For the statistical evaluation we used the single-factor analysis of variance at 95% significance level.

## RESULTS AND DISCUSSIONS

Measurement results relative to the fruit and leaf magnesium contents are illustrated by Table 2. The magnesium content of the fresh cucumber fruits did not show any increase in response to the foliar fertilizer applications. It can be ascribed to the fact that the fruits contained the nutrient examined in an amount that was over the value given for the cucumber in general (9 mg / 100 g) even in the case of no foliar fertilization and this way probably the plants did not assimilate any more magnesium in the fruits.

At the first measurement time, the Mg content of the leaves of the treatments with foliar magnesium application were higher than the value registered for the control, but on the second occasion we obtained results that were just the opposite.

**Table 2:** Fruit and leaf magnesium contents

| Plant part          | 1 <sup>st</sup> sampling (May 23) |                          |                          | 2 <sup>nd</sup> sampling (June 27) |                          |                          |
|---------------------|-----------------------------------|--------------------------|--------------------------|------------------------------------|--------------------------|--------------------------|
|                     | Control<br>( $\mu\text{g}$ )      | Mg1<br>( $\mu\text{g}$ ) | Mg2<br>( $\mu\text{g}$ ) | Control<br>( $\mu\text{g}$ )       | Mg1<br>( $\mu\text{g}$ ) | Mg2<br>( $\mu\text{g}$ ) |
| Leaf (1 g dry)      | 8227                              | 9475                     | 8435                     | 12447                              | 12169                    | 11045                    |
| Fruit (1 g dry)     | 2951                              | 2477                     | 2675                     | 2157                               | 2212                     | 2201                     |
| Fruit (100 g fresh) | 13052                             | 10846                    | 11198                    | 10075                              | 9413                     | 9345                     |

The effect of the treatments on the different nutrients examined is summarized in Tables 3 and 4.

According to the results of our statistical analyses:

1. **leaf Ga content** at the first sampling time was diminished by the foliar application of magnesium, but was increased at the end of the growing season, particularly in response to the treatment Mg2, though there was no significant difference between the two foliar fertilization treatments;
2. **leaf Mn content** was increased by the foliar fertilization, in particular in the case of the Mg treatment with the lower concentration;

**Table 3:** Changes in leaf and fruit nutrient compositions in response to magnesium fertilization (1<sup>st</sup> sampling)

| Treatment effect                                    | Result of statistical calculations (SD 95%) | Leaf                                 | Dry fruit                               | Fresh fruit                                 |
|-----------------------------------------------------|---------------------------------------------|--------------------------------------|-----------------------------------------|---------------------------------------------|
| increase in response to treatment Mg1 (Mg1>Control) | SD exists                                   | Zn                                   |                                         |                                             |
|                                                     | no SD                                       | Al, B, Li, Mg, Mn, Mo                | Al, B                                   | Al                                          |
| decrease in response to treatment Mg1 (Mg1<Control) | SD exists                                   |                                      | Ba, Ca, Fe, K, Mg, Na, Si, Sr           | Ba, Ca, K, Mg, Na, Si, Sr, Zn               |
|                                                     | no SD                                       | Ba, Ca, Fe, Ga, K, Na, P, Si, Sr     | Mn, Ni, P, Zn                           | B, Fe, Mn, Ni, P                            |
| increase in response to treatment Mg2 (Mg2>Control) | SD exists                                   |                                      | Fe                                      |                                             |
|                                                     | no SD                                       | Al, Li, Mg, Mn, Mo, Na, Si, Ti, Zn   | Al, B, Ba, Li, Mn, Zn                   | Al, B, Fe, Li, Zn                           |
| decrease in response to treatment Mg2 (Mg2<Control) | SD exists                                   |                                      | Ca, K, Na, Sr                           | Ca, K, Mg, Na, Si, Sr                       |
|                                                     | no SD                                       | B, Ba, Ca, Fe, Ga, K, P, Sr          | Mg, Ni, P, Si                           | Ba, Mn, Ni, P                               |
| increase in response to concentration (Mg1<Mg2)     | SD exists                                   |                                      | Ba, Fe                                  | Ba                                          |
|                                                     | no SD                                       | Ga, K, Mo, Na, Ni, Si                | B, Ca, K, Mg, Mn, Mo, Na, P, Si, Sr, Zn | B, Ca, Fe, K, Mg, Mn, Mo, Na, P, Si, Sr, Zn |
| decrease in response to concentration (Mg1>Mg2)     | SD exists                                   |                                      |                                         |                                             |
|                                                     | no SD                                       | Al, B, Ba, Ca, Fe, Mg, Mn, P, Sr, Zn | Al, Ni                                  | Al, Ni                                      |

3. **leaf Si content** was increased by the foliar fertilization at the second sampling time, in the case of the Mg1 treatment we observed significant increase as compared to the control; at the first measuring time, in response of the lower magnesium doses, the value was lower as compared to the control.
4. **leaf Ba content** at the second sampling time was significantly diminished in the Mg2 treatment as compared to the control and to the treatment Mg1;
5. **fruit Ca, K, Mg, Na, Si, Sr and Zn contents** at the first measuring time were significantly diminished in both magnesium treatments as compared to the control, while at the second measuring time we continued to observe a decrease for these elements (except for Ca and Mg) , but already there was no significant difference between the treatments;
6. **fruit Fe content** was significantly increased at the first measurement time by the treatment Mg2, but this was not observed at the second measurement time and a decrease was observed in this parameter at both measurement times in response to the lower dose;

**Table 4:** Changes in leaf and fruit nutrient compositions in response to magnesium fertilization (2<sup>nd</sup> sampling)

| Treatment effect                                    | Result of statistical calculations (SD 95%) | Leaf                                 | Dry fruit                     | Fresh fruit                          |
|-----------------------------------------------------|---------------------------------------------|--------------------------------------|-------------------------------|--------------------------------------|
| increase in response to treatment Mg1 (Mg1>Control) | SD exists                                   | Ga, Mn, Si                           |                               |                                      |
|                                                     | no SD                                       | B, Ca, Fe, K, Mo, Na, P, Ti, Zn      | Al, B, Ba, Ca, Mg, Sr         | Al                                   |
| decrease in response to treatment Mg1 (Mg1<Control) | SD exists                                   |                                      |                               | P, Zn                                |
|                                                     | no SD                                       | Al, Ba, Li, Mg, Sr                   | Fe, K, Mn, Na, P, Si, Zn      | B, Ba, Ca, Fe, K, Mg, Mn, Na, Si, Sr |
| increase in response to treatment Mg2 (Mg2>Control) | SD exists                                   | Ga                                   |                               |                                      |
|                                                     | no SD                                       | Al, Fe, K, Mn, Mo, Na, P, Si         | Al, B, Ba, Ca, K, Mg, Sr      | Al                                   |
| decrease in response to treatment Mg2 (Mg2<Control) | SD exists                                   | Ba                                   |                               | P, Zn                                |
|                                                     | no SD                                       | B, Ca, Li, Mg, Sr, Ti, Zn            | Fe, Mn, Na, P, Si, Zn         | B, Ba, Ca, Fe, K, Mg, Mn, Na, Si, Sr |
| increase in response to concentration (Mg1<Mg2)     | SD exists                                   |                                      |                               |                                      |
|                                                     | no SD                                       | Al, Ga, K, Li, Mo, Na                | Al, B, Fe, K, Na              | Al, B, Fe, K, Na                     |
| decrease in response to concentration (Mg1>Mg2)     | SD exists                                   | Ba                                   |                               |                                      |
|                                                     | no SD                                       | B, Ca, Fe, Mg, Mn, P, Si, Sr, Ti, Zn | Ba, Ca, Mg, Mn, P, Si, Sr, Zn | Ba, Ca, Mg, Mn, P, Si, Sr, Zn        |

7. at the second measurement time we observed that **fruit Zn and P contents** per unit of fresh weight were decreased compared to the control in both treatments receiving foliar application of magnesium.

8. **fruit Ba content** was significantly increased at the first measurement time with the increase in the concentration of the foliar fertilizer, while a slight decrease was observed during the second measurement period.

## REFERENCES

1. Fazekas T., Selmeczi B., Stefanovits P. (ed.): A magnézium forrásai és jelentősége az élővilágban. Akadémia Kiadó, Budapest, 1992.
2. Gastol, M., Domagala-Swiatkiewicz, I.: Effect of foliar sprays on potassium, magnesium and calcium distribution in fruits of the pear. *J. Fruit Ornament. Plant Res.* vol. 14 (Suppl. 176 2), 2006, 169-176
3. Göhler, F., Drews, M.: Fortschritte bei der Produktion von Gewächshausgurken durch Anwendung des Nährfließverfahrens. *Gartenbau*, 1990, 37(2), 38-40.
4. Marschner, J.D.: Mineral Nutrition of Higher Plants. Academic Press, London, 1986.
5. Moretti, G.: Effect of foliar treatments of magnesium, manganese, and zinc on grafted vines in the nursery. *Acta Hort.*, 2002, (ISHS) 594, 647-652.
6. Takács Hájos M., Kiss A. S., Csikkel Szolnoki A. and Eperjes Z. (2001): Evolution of Ca, Mg, Na and K contents in tomato fruit as affected by Mg foliage sprays. *Magnesium Research*. 2001, 14(4), 322.
7. Takácsné Hájos M., Stefanovitsné Bányai É., Kiss A. S.: Effect of Mg leaf fertilisation on morphological and economic parameters in saffron (*Carthamus tinctorius* L.). *Magnesium Research*. 2003, 16, 338.
8. Terbe I.: A hajtattott zöldségnövények magnéziumellátottsága és trágyázása. *Hajtattás, korai termesztés*, 1989, 20(3), 7-8.
9. Terbe I.: A hajtattott zöldségnövények magnéziumigénye és trágyázása. *Hajtattás, korai termesztés*, 1992. 23(1), 6-10.
10. Ushijima, T., Kaneko, A., Yamamoto, T.: Magnesium excess-mediated chlorosis in leaves of cucumber grown in soils of granodiorite origin under a plastic house. *Journal of the Japanese Society for Horticultural Science*, 2004, 73(5), 476-483

# BIODIVERSITY AND XENOBIOTICS RELATED TO FOODS OF ANIMAL ORIGIN

Alexandru T. BOGDAN<sup>1</sup>, Stelian ACATINCAI<sup>2\*</sup>, Adina-Elena AVACOVICI<sup>3</sup>,  
Zeno GÂRBAN<sup>2</sup>

1. Center of Studies and Research for Agroforestry Biodiversity of the Roumanian Academy-București, Calea 13 Septembrie Nr.13, RO-050 071 București, Roumania; 2. Faculty of Animal Science and Biotechnology and Faculty of Food Products Technology, University of Agricultural Sciences and Veterinary Medicine of Banat, Calea Aradului Nr. 119, RO-300 645, Timișoara, Roumania; 3. Faculty of Chemistry and Environmental Engineering, University Politehnica Timișoara, Pta Victoriei, Nr.2, RO-300 006 Timișoara, Roumania

\* E-mail : sacatincai@yahoo.com

## ABSTRACT

*Biodiversity is a notion that refers to the variability among living organisms from all sources, including terrestrial, marine and other aquatic ecosystems, and the ecological complexes of which they are part. This also includes diversity within species (genetic diversity), between species and of ecosystems. Biodiversity can be influenced by a number of natural and anthropogenic factors. The chemical composition of the environment and foods can be one of the factors influencing biodiversity. Xenobiotics are a class of chemical that can affect the living organisms and in some cases may interact with the deoxyribonucleic acid macromolecule generating cancerigenic, teratogenic or mutagenic effects. By their action at the level of the DNA macromolecule and also by their presence in the environment in high concentrations as anthropogenic pollutants xenobiotics can dramatically affect the ecosystem and may influence biodiversity.*

**Key words:** biodiversity, animal food products, xenobiotics

## 1. CONSIDERATIONS REGARDING BIODIVERSITY

Biodiversity is a concept that defines the variation of life forms within a given ecosystem, biome, or the entire Earth. A more concise definition is given by Gaston and Spicer (2004) which define biodiversity as the variation of life at all levels of biological organisation.

This concept regards the world in its complexity, from genes to species, through the broad scale of ecosystems, thus summarizing the problem of "life on Earth" (Gaston, 1996). In this acception the definition of biodiversity is of interest not only for "living organisms" (biochemistry) and their complex interactions but also for the interactions with the environment (geochemistry).

The term „natural diversity” was first used in “The Preservation of Natural Diversity” a study elaborated in 1975 by “The Science Division of The Nature Conservancy”. Later in 1985 Rosen uses for the first time the term “biodiversity” in a planning for the “National Forum on Biological Diversity” organized by the National Research Council (NRC) which was to be held in 1986, and first appeared in a

publication in 1988 when entomologist Wilson used it as the title of the proceedings of that forum. After 1986 the term and the concept of biodiversity was used more often by biologists, ecologists, political leaders and peoples involved in the domain.

In 1993, a statistic following the keywords found in Biological Abstracts revealed the fact that the term biodiversity was used 72 times and the term biological diversity was used 19 times.

The first book on biodiversity appeared in 1997 with the title "Biodiversity II" (Reaka-Kudla et al. 1997). The book documents the rapid rise of the term "biodiversity" regarding its theoretical and applicative importance and influence. Also the book traces the study of aspects of biodiversity back to Aristoteles in the Antiquity

After the year 2000 the concept of biodiversity is more often found in the scientific publications, in the thematic of various scientific manifestations, and also appeared an increasing number of work groups and research centers regarding this domain (Bogdan et al., 2008).

## **2. CHARACTERISTICS OF BIODIVERSITY**

### **2.1. Natural variations**

The natural variations are conditioned by the evolution in time and can be classified in 3 subcategories: a) spatial variations; b) functional variations; c) morphological variations.

Spatial variations of biodiversity - may appear due to geological processes or to climacteric changes. The main geological processes that may influence biodiversity are the tectonical processes (tectonically movements, orogenesis, soil corrosion) and the marine processes (marine transgression and regression).

Functional variations - are time related and may refer to: circadian, circamensual, seasonal and circannual variations.

Morphological variations - refer to natural selection associated to geological and climacterical changes

The natural process of new species formation, which is the motor of biodiversity, usually takes place in two ways. The first way is common to plants and consists in the process of polyploidy. This is a simple multiplication in the number of gene-bearing chromosomes - sometimes within a preexisting species and sometimes in hybrids between two species. A second major process is geographic speciation and takes much longer. It starts when a single population (or series of populations) is divided by some barrier extrinsic to the organisms. The isolated populations then diverge from each other in evolution because of the inevitable differences of the environments in which they find themselves. Since all populations evolve when given enough time, divergence between all extrinsically isolated populations must eventually occur (Wilson, 1988).

In order to have an image of the magnitude of natural biodiversity one must mention that in the year 2000 there were around 1,75 million species named by taxonomists (UNEP-WCMC, 2000).

### **2.2. Artificial variations**

The artificial variations are conditioned by the human intervention on: a) the genome; b) eukaryotic organisms (animals and plants) and prokaryotic organism (microorganisms); c) products of vegetal and animal organisms .

Variations triggered by human interventions on the genome are of interest for medical biology (genic therapy) and for agricultural biology (genetically modified organism).

Those variations conditioned by human interventions on eukaryotic and prokaryotic organisms are regarding the growth-development systems (e.g.: farm animals), the cultivation systems related to hydric regime, temperature and minerals from soil

Variations due to human intervention on products of vegetal and animal origin have as destination: food industry, light industry, pharmaceutical industry, etc

An aspect of the modern science that can affect biodiversity is the genetic engineering that is increasingly used for the genetic improvement of crops. The organisms produced are referred to as genetically modified organisms (GMOs) or living modified organisms (LMOs) and efforts have focused on such crops as tomatoes, grains, cassava, maize and soybeans. Genes from any species of plant, animal, or microorganism can now be moved into a particular plant. Genetic engineering is not without risks and concerns have been raised about potential risks to biodiversity (Committee on Noneconomic and Economic Value of Biodiversity, 1999). In response to this concern, a subsidiary agreement to the Convention on Biological Diversity was negotiated to address the potential risks posed by cross-border trade and accidental releases of GMOs. Adopted in January 2000, the Cartagena Protocol on Biosafety was developed to ensure that recipient countries have both the opportunity and the capacity to assess risks relating to GMOs, and to ensure their safe transfer, handling and use.

### **3. BIOCONSTITUENTS OF ANIMAL PRODUCTS OF NUTRITIONAL INTEREST**

The components of biodiversity (e.g.: plants, animals) are the source for foods, drugs, fibers, biofuels, and industrial products. About 150 species of plants have entered world commerce, and 103 species account for 90% of the supply of food plants by weight, calories, protein, and fat for most of the world's countries (Prescott-Allen and Prescott-Allen 1990). Just three crops - wheat, rice, and maize - account for roughly 60% of the calories and 56% of the protein consumed directly from plants (Yeatman et al, 1985).

General data regarding the composition of natural food products (unprocessed) are given in Table 1.

#### **3.1. General aspects regarding nutrients**

Nutrients are the chemical constituents of foods which, after the process of metabolism, ensure the morphophysiological status of the organism and the energy necessary for all vital processes. As a generally accepted classification, nutrients are grouped in: a) macronutrients – carbohydrates, lipids, proteins; b) micronutrients – biominerals, vitamins; other nutrients – water, alimentary fibers, bioactive substances.

Literature presents the complex problems of the mechanisms specific for toxicogenesis with implications on public health (Goudez-Perriere et al., 2003).

#### **3.2. Particularizations for nutrients in foods of animal origin**

In foods of animal origin predominate the lipids (acylglycerols, sterides, phospholipids, fatty acids etc.), proteins (aminated acids, holo- and heteroproteides).

Also there are significant quantities of biominerals and water. In more reduced amounts are carbohydrates, vitamins. There are no dietary fibers. Numerous papers present in detail this problem (Watt et al., 1975; Moll and Moll, 2003).

## **4. BIOCHEMISTRY AND XENOBIOCHEMISTRY IN RELATION WITH BIODIVERSITY**

### **4.1. Comparative aspects**

Xenobiochemistry or is a domain of Biochemistry which regards both molecular biology and pathobiochemistry. In order to better understand the field of work of xenobiochemistry, we must first define the term of xenobiotic and make the distinction between xenobiotics and nutrients (Garban, 2005).

Nutrients, as base constituents of aliments, accede into the organism through the digestive way (enteral) or parenteral (in therapeutic conditions). In the organism, the nutrients follow the metabolization processes generating various metabolites with a great importance in all the functions of the body. The nutrients are involved especially in morphogenetic and energogenetic processes (Dumitrache, 1996; Kastori, 2000).

In the case of nutrients, which are studied by biochemistry, the concept of metabolisation defines their physiologic role in the organism. Metabolisation is characterized by two phases: a) catabolism (the totality of biodegradation reactions peculiar to nutrients); b) anabolism (the totality of biosynthesis reactions peculiar to nutrients).

Xenobiotics (xenos – foreign; bios – life), as non-nutritive substances accede in the organism usually along with the other nutrients and follow, gradually, the biotransformation process.

In the case of xenobiotics, which are studied in xenobiochemistry, the definitory concept is biotransformation. Similar to metabolism, in biotransformation there are also two phases:

a) xenobiodegradation – defined by the totality of biodegradation reactions peculiar to xenobiotics (i.e.: the oxydo-reduction reaction and hydrolyses reaction). After these reactions, the non-polar compounds which are biologically inactive are brought to a polar state and thus activated.

b) xenobiosynthesis – including the totality of biosynthesis reactions which are specific to xenobiotics. The xenobiosynthesis reactions contain conjugation reactions and reaction with adduct formation. The conjugation reactions are specific for the binding of xenobiotics to some endogenous substances e.g.: glucuronic acid, glycolol, acetylated compounds, sulphonic compounds, etc. The reactions that lead to adducts biogenesis take place between various organic and inorganic compounds and deoxyribonucleic acid (DNA).

### **4.2. Xenobiotics of alimentary interest as disturbants of biodiversity**

Chemical xenobiotics of alimentary interest accede in the organism usually along with the nutrients. Once inside the body xenobiotics become part of the biotransformation process yielding various residual compounds which in some cases are electrophyl radicals that can interact inside the cells with the nucleophylic deoxyribonucleic acid (DNA) macromolecule. In the case of xenobiotics - DNA interactions a new complex arises called DNA adduct.

Among the xenobiotics that can lead to DNA adducts formation there are compounds from the following classes: a) metallic ions (e.g.: Hg, Cd); b) nitrosamines, polycyclic aromatic hydrocarbons, steroidal compounds, micotoxins (e.g.: aflatoxines),



**Table 1.** General chemical composition of the main food products of animal origin

| Foods<br>(name and description) |                  | Water<br>(%) | Proteids<br>(g) | Lipids<br>(g) | Carbohydrates (g) |       | Biominerals (mg) |                               |                  |                 |                |
|---------------------------------|------------------|--------------|-----------------|---------------|-------------------|-------|------------------|-------------------------------|------------------|-----------------|----------------|
|                                 |                  |              |                 |               | Total             | Fibre | Ca <sup>2+</sup> | PO <sub>4</sub> <sup>3-</sup> | Fe <sup>2+</sup> | Na <sup>+</sup> | K <sup>+</sup> |
| B                               |                  | C            | E               | F             | G                 | H     | J                | K                             | L                | M               | N              |
| ovine                           | Beef meat        | 80.3         | 14.7            | 4.0           | -                 | -     | 8.0              | 127                           | 2.0              | 228             | 183            |
|                                 | Veal meat        | 80.7         | 15.5            | 2.7           | -                 | -     | 10               | 145                           | 1.7              | 226             | 214            |
|                                 | Organs           | Hearth       | 77.5            | 17.1          | 3.6               | 0.7   | 5                | 195                           | 4.0              | 86              | 193            |
|                                 |                  | Liver        | 69.7            | 19.9          | 3.8               | 5.3   | 8                | 352                           | 6.5              | 136             | 281            |
|                                 |                  | Kidney       | 75.9            | 15.4          | 6.7               | 0.9   | 11               | 219                           | 7.4              | 176             | 224            |
| swine                           | Potted meat      | 60.7         | 17.5            | 19.2          | -                 | -     | -                | -                             | -                | -               | -              |
|                                 | Fat class        | 33.4         | 9.1             | 57.0          | -                 | -     | 5                | 88                            | 1.4              | 70              | 281            |
|                                 | Medium fat class | 37.3         | 10.2            | 52.0          | -                 | -     | 6                | 103                           | 1.5              | 70              | 281            |
|                                 | Thin class       | 41.1         | 11.2            | 47.0          | -                 | -     | 6                | 116                           | 1.7              | 70              | 281            |
| vine                            | Lamb             | 56.3         | 15.4            | 27.1          | -                 | -     | 9                | 135                           | 1.1              | 75              | 291            |
|                                 | Organs           | Hearth       | 71.6            | 16.8          | 9.6               | 1.0   | 11               | 249                           | -                | -               | -              |
|                                 |                  | Liver        | 70.8            | 21.0          | 3.9               | 2.9   | 10               | 349                           | 10.9             | 52              | 203            |
|                                 |                  | Kidney       | 77.7            | 16.8          | 3.3               | 0.9   | 13               | 218                           | 7.6              | 200             | 231            |
| oultry                          | Turkey           | 64.2         | 20.1            | 14.7          | -                 | -     | -                | -                             | -                | -               | -              |
|                                 | Hen              | 56.9         | 17.4            | 24.8          | -                 | -     | 10               | 167                           | 1.4              | -               | -              |
|                                 | Chicken          | 73.7         | 23.4            | 1.9           | -                 | -     | 11               | 218                           | 1.1              | 50              | 321            |
|                                 | Goose            | 51.1         | 16.4            | 31.5          | -                 | -     | 10               | 176                           | 1.6              | -               | -              |
|                                 | Quail            | 65.9         | 25.0            | 6.8           | -                 | -     | -                | -                             | -                | -               | -              |
|                                 | Duck             | 54.3         | 16.0            | 28.6          | -                 | -     | 10               | 176                           | 1.6              | -               | -              |
| Fish                            | Cod              | 81.2         | 17.6            | 0.3           | -                 | -     | 10               | 194                           | 0.4              | 70              | 381            |
|                                 | Crap             | 77.8         | 18.0            | 4.2           | -                 | -     | 50               | 253                           | 0.9              | 50              | 281            |
|                                 | Herring          | 79.4         | 17.5            | 2.6           | -                 | -     | -                | 225                           | 1.3              | 74              | 421            |
|                                 | Salmon           | 63.6         | 22.5            | 13.4          | -                 | -     | 79               | 186                           | 0.9              | -               | -              |
|                                 | Pike             | 78.3         | 19.3            | 1.2           | -                 | -     | -                | 214                           | 0.4              | 51              | 311            |

etc. (Landis and Yu, 1995; Hodgson and Smart, 2001). These compounds are often found as food pollutants and some of them may lead to carcinogenic, mutagenic and teratogenic effects, thus influencing the genome and the biodiversity (Garban, 2008).

Referring at the contamination induced by metals there are important scientific papers that present the problem of their mutagenous and carcinogenous effects (Kazantsis and Lorna, 1979).

Genetic diversity provides the raw material for plant breeding, which is responsible for much of the increases in productivity in modern agricultural systems. Much of the genetic diversity available for crop breeding is now stored in a network of national and international genebanks administered by the UN Food and Agriculture Organization, the Consultative Group on International Agricultural Research, and various national agricultural research programs. The important contribution of genebanks to agricultural productivity has been recognized by government since the 18th century. It led to the rise of botanical gardens and expeditions in search of new plant varieties (Fowler 1994).

Another important aspect of influence of xenobiotics is the use of nitrates in agriculture. Besides their effects as xenobiotics of nutritional interest that range from interactions with hemoglobin up to formation of nitrosamines with cancerigene, teratogene and mutagene effects, nitrites affects the nitrogen cycle with unpredictable outcomes for biodiversity. Nitrogen accumulation in the environment has become a major cause of biodiversity loss. It has increased substantially in recent decades, primarily as a result of an increase in the use of fertilizer and the burning of fossil fuels. Increased nitrogen in soil and water can lead to loss of species and shifts in the species composition of plant communities (Wedin and Tilman, 1996).

## **5. SUSTAINABLE AGRICULTURE IN RELATION WITH THE PROBLEMS OF BIODIVERSITY AND XENOBIOCHEMISTRY**

Sustainable agriculture is a concept defined by Perlas (1983) which in time led to the founding of International Alliance for Sustainable Agriculture (IISA).

The spreading of the studies and the broadening of the knowledge related to sustainable agriculture has lead to the elaboration of a generic model published in the book "Seven Dimensions of Sustainable Agriculture"(Perlas, 1993).

These seven dimensions are correlated with Agricultural biodiversity and Agricultural xenobiology (the xenobiochemistry applied in this field).

In this context one must take into account the following:

- 1) instead of pesticides, is recommended the use of ecological management principles;
- 2) replacement of chemical fertilizers with a integrated management of soil fertilization;
- 3) the use of monocultures is recommended to replace policultures, thus developing the biodiversity;
- 4) replacement of chemical treated seeds with alternatives strategies of reproduction, strategies that are adopted in the ecological practice
- 5) instead of soil erosion and water depletion, soil and water conservation

- 6) replacement of animal growing based on large farms with a zooculture based on growing animals in households or small farms where the ecological measures can be better implemented and supervised;
- 7) instead of applying techniques based on genes and chemical substances, using the formative energies from nature (i.e. biodynamic preparatives) is recommended

These percepts, even if without a rigorous explanation and without elucidated methods, can be constituted as “operating means” in sustainable agriculture.

Regarding foods, providing an adequate composition of nutrition, without contaminant metals, assures foods safety for the consumer. There are also preoccupations and papers elaborated on this topics (Lee, 1972; Moll and Moll, 2003).

## CONCLUSIVE ASPECTS

Biodiversity in the animal reign, for interest in farm animal breeding and valorisation of their products, presents a special interest because almost all products are destined for foods. One can observe the existence of a biodiversity of composition of foods originating from animals, conditioned by biogeographical conditions, ecologic frame, breeding technologies and evidently, animals feeding.

It must be mentioned the specificity of chemical composition interesting the proteins, lipids, carbohydrates and biominerals present in meat, organs at diverse species but also in animal products.

The paper presents aspects referring to meat and organs, showing data regarding their natural composition. From the point of view of xenobiochemistry this composition can be denaturated by the presence of diverse contaminants of organic nature (polycyclic aromatic hydrocarbons, micotoxins, steroids) or of inorganic nature (heavy metals, nitrates, nitrites).

Beside the contamination that appears in the existential environment of animals, there can also be remarked a contamination produced during processing of animal products in nutritional purpose (e.g. additivation with preservatives, sweeteners etc.).

In this context studies on biodiversity from the last decade approach more and more the problem of contaminants exclusion and the orientation toward an ecological nutrition.

These problems are possible applying the principles of a sustainable agriculture.

**Acknowledgements:** The authors express thanks to George-Daniel Ghibu – PhD student in chemistry and Cornel Baltă – PhD student in Veterinary Medicine for their participation at documentation in the domaine of xenobiotics of nutritional interest, some mentioned in the present paper.

## REFERENCES

1. Bogdan A.T. – Intedisciplinaritatea în investigarea biodiversității agro-silvice – unpublished data (2008)
2. Dumitrache S. - Igiena alimentației, pp.244-358, în “*Igiena*” (Mănescu S., Tănăsescu G., Dumitrache S., Cucu M., Eds.), Ed. Medicală, București, 1996
3. Garban Z. – Xenobiochimie: Tratat comprehensiv, Vol. IV, Editura Eurobit, Timișoara, 2005.
4. Garban Z. – Xenobiotice chimice de interes alimentar cu impact asupra calității cărnii și produselor de carne. Comun. La Primul Simpozion al Centrului de Studii și Cercetări pentru Biodiversitate Agro-Silvică “Acad. David Davidescu” cu tema “Biodiversitatea raselor de taurine de carne”, București 17 iulie 2008.

5. Gaston K.J. (ed). - Biodiversity: a biology of numbers and difference, Oxford: Blackwell, 1996.
6. Gaston K.J., Spicer J.I. - Biodiversity: an introduction, 2nd Ed, Blackwell Publishing, 2004.
7. Goudez-Perrière F., Bon C., Meney A., Puiseux-Dao S. - *Toxicogenese*, Elsevier-Lavoisier, Paris, 2003
8. Hodgson E., Smart R.C. (Eds.) - *Introduction to biochemical Toxicology*, 3rd ed., New York,, Wiley, 2001.
9. Kastori R., Sovljanski Radmila, Monarov Elizabeta, Maletin S., Pujin V. - *Safe Food*, Vol.I - II, Eko-Konferencija, September 27-30, 2000, Novi Sad 2000.
10. Kazantzis G., Lorna J. Lilly – Mutagenic and carcinogenic effects of metals, Chap.14, pp.1-36, in *Handbook on the toxicology of metals* (Friebert L. et al., Eds.), Elsevier, North-Holland Biomedical Press, 1979.
11. Landis W.G., Yu M.H. – *Introduction to Environmental Toxicology*, Lewis Publishers, Boca Raton, Florida, 1995.
12. Lee D.N. (Ed.) – *Metallic Contaminants and Human Health*, Academic Press, New York, 1972.
13. Moll M., Moll Nicole (Eds.) - *Sécurité alimentaire du consommateur*, Editions Lavoisier, Collection Science et Technologie, Paris, 2003.
14. Perlas N. – The seven dimensions of sustainable agriculture, Publ. by Center for Alternative Development Initiatives, Quezon City, Philippines, 1993
15. Prescott-Allen R, Prescott-Allen C. – How many plants feed the world?, Cons. Biol., 1990, 4, 365-374.
16. Watt K., Merrill A.L., Pecot K. Rebecca, Adams F. Catherine, Orr L. Martha, Miller D.F. – *Handbook of the Nutritional contents of foods*, United States Department of Agriculture, Dover Publ. Inc., New York, 1975.
17. Wedin D., Tilman, D. - Influence of nitrogen loading and species composition on carbon balance of grasslands, Science, 1996, 274, 1720-1723
18. Wilson E.O. (ed), Peter F.M. (assoc. ed.) - Biodiversity, National Academy Press, 1988.
19. Young T., Burton M.P. - *Agricultural Sustainability: definition and implications for agricultural and trade policy*, FAO, Rome, 1992.
20. Yeatman C.W, Kafton D., Wilkes G. (eds) - Plant genetic resources: a conservation imperative, AAAS Selected Symposium #87, Westview Pr. Co., 1985.
21. \*\*\* Committee on Noneconomic and Economic Value of Biodiversity - Perspectives on Biodiversity: Valuing Its Role in an Everchanging World, National Academy Press Washington D.C., 1999.
22. \*\*\* United Nations Environment Programme - World Conservation Monitoring Centre (UNEP-WCMC) - Global Biodiversity: Earth's living resources in the 21st century, World Conservation Press, Cambridge, 2000

# SNUTRIENT ELEMENT OF STONE FRUIT SPECIES AND CULTIVARS

Blanka SZILVÁSSY<sup>1</sup>, Nóra PAPP<sup>1</sup>, Zoltán SZABÓ<sup>2</sup>, József NYÉKI<sup>2</sup>,  
Enikő TORDAI<sup>3</sup>, Andrzej PEDRYC<sup>3</sup>, Éva STEFANOVITS-BÁNYAI<sup>1</sup>,  
Attila HEGEDŰS<sup>1,3\*</sup>

*1\*. Department of Applied Chemistry, Faculty of Food Science, Corvinus University of Budapest, Villányi út 29., H-1118 Budapest, Hungary; 2. Centre of Agricultural Sciences, Institute for Extension and Development, University of Debrecen, Böszörményi út 138, H-4032 Debrecen, Hungary; 3. Department of Genetics and Plant Breeding, Faculty of Horticultural Science, Corvinus University of Budapest, Ménesi út 44, H-1118 Budapest, Hungary*

\* E-mail: hegedus.attila@uni-corvinus.hu

## ABSTRACT

Several epidemiological studies revealed that the consumption of antioxidant compounds and the risk of atherosclerosis, increased blood pressure or cancer are inversely proportional. The individual amounts and relative proportions of macro- and micro elements present in food is also of great consideration since these are involved in a wide range of physiological processes including the influence of the redox homeostasis. This study was carried out to survey the antioxidant power and mineral element content in fresh fruits of four stone fruit cultivars. The elemental analysis of fruits was carried out by ICP-OES (Inductively Coupled Plasma Optical Emission Spectrometry). Considerable differences in element contents between cultivars of the same species occurred most often in sour cherries and apricots. Such differences could be primarily attributed to their different genetic constitutions. Japanese plums contained the smallest quantities from most of the elements. It points to their potential use in the treatment of some neurodegenerative diseases, where the restriction of the exposure to Al, Ba, Fe, Na, Mn, Zn is crucial. A detailed knowledge on the nutrient element content of fruits may offer the possibility to construct individual diets in response to the varied requirements arising from specific health conditions and promote the medical treatment and help the patients to get better in a natural way.

**Key words:** healthy diet, macro and microelements, minerals, Prunus, stone fruit

## INTRODUCTION

Hungary has been witnessed a slight increase in the production of main stone fruit crops (eg. cherries, apricot, peaches and nectarines and plums) (Faostat, 2007). Probably the consumption of stone fruits will also rise, because people pay more and more attention for their health. Therefore, natural foods having enhanced functional properties may be very popular in the future.

The increased consumption of fresh fruits has several health benefits. Because of their high antioxidant capacity, they inhibit the harmful oxidative processes, and reduce the risk of several degenerative diseases (i.a. cancer, stroke, cardiovascular diseases) (Liu, 2003; Scalzo et al., 2005; Dauchet and Dallongeville,

2008). In addition, minerals have also considerable health effects (Stefanovits-Bányai et al., 2006; Hegedűs et al., 2008). Stone fruits are rich sources of the most important macro- (Na, K, Ca, Mg) and microelements (Fe, Cu, Zn, Mn) that have a considerable contribution to the human redox homeostasis through several mechanisms, e.g. they are indispensable cofactors of antioxidant enzymes. The transient metals, Fe and Cu ions may induce oxidative hazard because they can easily donate electrons to other molecules (Toyokuni, 2002; Valko et al., 2005). The Zn helps to build biomembranes, in absence of this metal element biomembranes may be easily disorganized. Knowledge on the Na/K ratio is indicative, because of the high Na levels in the human diet. Ca and Mg are required for the healthy bones and teeth, normal blood circulation, as well as for the efficient working of nerves and muscles.

The aim of this study was to evaluate the differences in the element contents of stone fruit species and cultivars.

## MATERIALS AND METHODS

**Fruit samples.** Experiments were carried out by using 4 sour cherry ('Érdi bőtermő', 'VN-7', 'Oblacsinszka', 'Csengődi'), 4 sweet cherry ('Karina', 'Celeste', 'Firm Red', 'Sylvia', 'Van'), 5 Japanese plum ('Super Giant', 'Black Amber', 'Black King', 'Fortune') and 3 apricot ('Zard', 'Gönci Magyar', 'Auróra') cultivars.

**Sample preparation.** 100 g fruit were homogenized (peel and flesh together) and freeze-dried.

**Biochemical assays.** 0.2 g of the dried fruit samples were weighted in teflon bomb (PTFE). 2.0 cm<sup>3</sup> nitric acid and 2.0 cm<sup>3</sup> hydrogen-peroxide were added to the samples and they were left standing for a night. Next day the closed teflons were boiled in hot water for half an hour. The digested samples were transfused in volumetric flask and supplemented to the volume of 10 ml with Milli-Q water. This solution was filtered through filter-paper into test tubes. The following elements were determined by ICP-OES (Thermo Jarrell Ash Co, ICAP 61): Al, As, B, Ba, Ca, Cd, Co, Cr, Cu, Fe, K, Li, Mg, Mn, Na, P, Pb, and Zn.

## RESULTS AND DISCUSSIONS

Sodium contents in the analysed fruits showed great discrepancies. In average, Na contents in sweet cherries and apricots were 2-3-fold higher than the levels that sour cherries and Japanese plums had. 'Érdi bőtermő' proved to be an exceptional sour cherry cultivar since its fruits have accumulated the highest quantities of Na. The average potassium content of all tested fruits was similar with only some outstanding values measured for the sour cherry cultivar 'Oblacsinszka' and the three apricots.




**Fig. 1.** Na and K contents in stone fruits. The grey, striped, V mottled columns and those with black diamonds refer to sour cherry, sweet cherry, Japanese plum and apricot cultivars, respectively.

From a nutritional aspect, the Na/K ratio may be of crucial importance. In the values there are fairly large margins. The lowest and hence the most favourable ratio was determined Japanese plums and in three of the sour cherry cultivars. These fruits contained the lowest levels of Na and the highest amounts of K. The sour cherry cultivar, 'Érdi bőtermő' proved to be an exception again with an outstandingly high Na/K ratio. Sweet cherries were also characterized by high values. The ratios were also high in case of two apricot cultivars, but were a little bit lower than those for sweet cherries.




**Fig. 2.** Na/K ratio in stone fruits. The grey, striped, V mottled columns and those with black diamonds refer to sour cherry, sweet cherry, Japanese plum and apricot cultivars, respectively.

The apricots, sour and sweet cherries contained similar levels of Ca. Japanese plums had significantly lower levels from Ca, while the highest values could have been determined for the sour cherry 'Oblacsinszka', which possessed 22,1 mg Ca/100g fresh weight. Japanese plums also had the lowest quantities of Mg; without notable fluctuations among the tested cultivars. The rest of the fruits showed 2- to 3-fold higher Mg concentrations than plum cultivars. In our study, the 'Oblacsinszka' sour cherry had the highest Mg concentration (14,8 mg/100 g fresh weight).




**Fig. 3.** Ca and Mg content of stone fruits. The grey, striped, V mottled columns and those with black diamonds refer to sour cherry, sweet cherry, Japanese plum and apricot cultivars, respectively.

In average, apricots showed the highest Cu contents, but concentrations were quite similar in the samples. The results ranged from 0.09 to 0.15 mg/100 g fresh weight. Only three cultivars could be identified as accumulating significantly lower values of Cu in their fruits. These include 'Csengődi' sour cherry, 'Celeste' sweet cherry and 'Gönci magyar' apricot. The highest amount of Fe was determined for

'Karina' sweet cherry (0.37 mg/100 g fresh weight). Japanese plums showed the lowest values of Fe, its concentration ranged from 0.15 to 0.19 mg/100 g fresh weight in plums.




**Fig. 4.** Cu and Fe content of stone fruits. The grey, striped, V mottled columns and those with black diamonds refer to sour cherry, sweet cherry, Japanese plum and apricot cultivars, respectively.

As considers the Mn concentrations there could be observed only one outstanding value: in average, the 'Celeste' sweet cherry showed 2-fold higher element content than the rest of the fruits. From sour cherry cultivars, the 'VN7' had the highest values and approached the level determined for 'Celeste'. The average Zn-content in Japanese plums and the sour cherries 'Csengődi' and 'VN7' was somewhat lower than the rest of the fruits, while the apricot cultivar 'Aurora' accumulate outstanding Zn levels.




**Fig.5.** Mn and Zn content of stone fruits. The grey, striped, V mottled columns and those with black diamonds refer to sour cherry, sweet cherry, Japanese plum and apricot cultivars, respectively.

## CONCLUSIONS

From the tested macro-elements relevant differences were obtained for the Na/K ratio where obviously the Na- concentration defines the value. For example, foods with high Na-content are not recommended for people who suffer from kidney- or nervous diseases. In contrast, fruits with low Na/K ratio (like Japanese plums) might be useful to compensate for high Na levels in a typical human diet (John et al., 2002). The Ca and Mg-contents were the lowest in the Japanese plums while the other three fruit showed nearly identical values.



The micro-element contents were also different among the species. In average, the Japanese plums accumulated the lowest levels from all four elements. It might point to their use in the treatment of some neurodegenerative diseases, where the decrease of the exposure to Al, Ba, Fe, Na, Mn, Zn is important (Stefanovits-Bányai et al., 2006). The results of the sour and sweet cherries were similar. Their values were in the middle of the range. The element content of the apricots revealed the most interesting tendencies. Their Cu and Zn-concentrations were the highest among all fruits, while their Fe and Mn levels were almost as low as for the Japanese plums.

Considerable differences in element contents between cultivars occurred most often in case of the species, sour cherries and apricots. Since all trees were grown under identical climatic and soil conditions, such differences among cultivars within each species may be attributed to their different origins and genetic constitutions (Halász, 2007a,b; Pedryc et al., 2005; Szabó, 2007; Szilvássy et al., 2008). It might offer the possibility to construct individual diets in response to the varied requirements arising from specific health conditions and promote the medical treatment and help the patients to get better in a natural way.

Stone fruits are characterized by seasonal availability in the markets. The food processing industry should find alternative ways to process these fruits while preserving their beneficial compounds, antioxidants, vitamins and mineral elements. Vegetable and fruit consumption are very important factors of a health-promoting diet, and statistic data show that Hungarian people do not eat enough from these foods. New products, which are rich in health protecting components, together with the required marketing strategy may help to alleviate this problem.

**Acknowledgements.** This work was co-financed by the Ányos Jedlik programme NKFP06A2-BCETKA06 and the NKTH-OTKA K68921 grant. Attila Hegedűs is grateful for receiving a János Bolyai Scholarship.

## REFERENCES

1. Dauchet L., Dallongeville J. : Fruit and vegetables and cardiovascular disease: epidemiological evidence from the non-Western world, *Brit. J. Nutr.*, 2008, 99, 219–220.
2. Halász J. : Review of self-(in) compatibility in apricot (*Prunus armeniaca* L.). *World J. Agric. Sci.*, 2007, 3(5), 670–677.
3. Halász J. : A kajszi önmeddőségét meghatározó S-allél-rendszer molekuláris háttere. Doktori (PhD) értekezés, Budapesti Corvinus Egyetem, Budapest, 2007
4. Hegedűs A., Balogh E., Engel R., Sipos B.Z., Papp J., Blázovics A., Stefanovits-Bányai É. : Comparative nutrient element and antioxidant characterization of berry fruit species and cultivars grown in Hungary, *HortScience*, 2008, 43(6): 1711–1715.
5. John J.H., Ziebland S., Yudkin P., Roe L.S., Neil H.A.W. : Effects of fruit and vegetable consumption on plasma antioxidant concentrations and blood pressure: A randomised controlled trial, *Lancet*, 2002, 359, 1969–1974.
6. Liu R.H. : Health benefits of fruit and vegetables are from additive and synergistic combinations of phytochemicals, *Amer. J. Clin. Nutr.*, 2003, 78, 517S-520S.
7. Pedryc A., Hermán R., Halász J. : A Budapesti Corvinus Egyetem KTK Genetika és Növénynevelés Tanszéken előállított új kajsziarackhibridek értékelése. XI.

- Növénynemesítési Tudományos Napok, 2005. március 3-4., Összefoglalók (Szerk.: Kertész Z.), MTA, Budapest, 2005
8. Scalzo J., Politi A., Pellegrini N., Mezzetti B., Battino M. : Plant genotype affects total antioxidant capacity and phenolic contents in fruit, *Nutrition*, 2005, 21, 207–213.
  9. Stefanovits-Bányai É., Szentmihályi K., Hegedűs A., Koczka N., Váli L. Taba G., Blázovics A. : Metal ion and antioxidant alterations in leaves between different sexes of *Ginkgo biloba* L., *Life Sci.*, 2006, 78(10), 1049–1056.
  10. Szabó, T. : Az északkelet-magyarországi meggy tájfajta-szelekció eredményei és gazdasági jelentősége. Doktori (PhD) értekezés, Budapesti Corvinus Egyetem, Budapest, 2007.
  11. Szilvássy B., Papp N., Szabó Z., Nyéki J., Stefanovits-Bányai É., Hegedűs A. : Csonthéjas gyümölcsök mikroelem-tartalmának szerepe a táplálkozásban, *Acta Agronomica Óváriensis*, 2008, 50(1), 87–94.
  12. Toyokuni S. : Iron and carcinogenesis: from Fenton reaction to target genes, *Redox. Rep.*, 2002, 7(4), 189–197.
  13. Valko M., Morris H., Cronin : M.T.D. - Metals, toxicity and oxidative stress, *Curr. Med. Chem.*, 2005, 12(10), 1161–1208.
  14. \*\*\*Faostat, 2007. Crops data. 20 Nov 2008. <http://faostat.fao.org>.

# INTESTINE AND LIVER HISTOPATHOLOGICAL CHANGES INDUCED IN FED AND STARVATED GOLDFISH (*Carassius auratus auratus*, *Cyprinidae*, *Pisces*) DURING CHRONIC EXPOSURE TO MANGANESE

Andrea-Cristina STAICU\*, Carmen POSTOLACHE, Maria-Cristina MUNTEANU, Marieta COSTACHE, Anca DINISCHIOTU

*Multiple Users Research Center in Molecular Biology, Faculty of Biology, University of Bucharest, Splaiul Independentei, No 91-95, Bucharest, RO- 050 095, Romania,*

\* E-mail: crista@bio.unibuc.ro

## ABSTRACT

*Two lots of *Carassius auratus auratus* were exposed to sublethal concentration of manganese for 28 days. A lot of individuals were fed daily to satiation, the other one starved during manganese intoxication. Intestine and liver tissue fragments were used in order to evaluate manganese concentration and structural changes. Metal exposure stress caused no mortality. Nutrition status, fish size and growth rate as well as tissue heavy metal concentration are important data for monitoring assessment. After 28 days of exposure to the pollutant fish exhibited statistically insignificant decrease in their weights. Manganese concentration was higher in intestine compared to liver. Intestine manganese concentration in fed fish was lower than in starved fish. Both stress factors induced histological changes to diminish absorption surface of the intestine. Injuries of epithelial layer, tip of the villi changes, villi fusion, macrophages accumulation and sinusoids capillaries dilatation were frequent.*

**Key words:** manganese pollution, intestine, liver histopathological changes

## INTRODUCTION

Manganese was regarded as one of the least toxic elements but more recent experimental and epidemiological studies have shown that exposure to manganese can indeed lead deleterious changes (European Commission, Health and Consumer Protection Directorate General, 2002).

Manganese accumulates in mitochondria, a major source of superoxide, which can oxidize  $Mn^{2+}$  to the powerful oxidizing agent  $Mn^{3+}$ . Oxidation of important cell components by  $Mn^{3+}$  has been suggested as a cause of the toxic effects of manganese.

Teleost intestine surface represents a selectively permeable barrier that allows nutrient absorption, but excludes most toxic substances and pathogenic organisms. Additionally, the intestine is involved in water and electrolyte balance, active transport of salts and water, as part of hydromineral homeostasis, immunity, and regulation of

digestion and metabolism, the absorptive transport of nutrients, Intestinal structure responds rapidly and reversibly to changes in dietary inputs.

In order to assess manganese effects on fish digestive system, intestine and liver histological changes and the influences of nutritional status on metal toxicity and accumulation were analyzed.

## MATERIALS AND METHODS

Two years old individuals of *Carassius auratus gibelio* were kept in 150 liter tanks for two weeks before metal exposure. Pathological symptoms did not appear and fish were in good state of health. After acclimation period, manganese chloride was added in tap water to a final manganese concentration of 75 µg/L. Ten aquaria of 40 l capacity were used and each one was occupied by 10 individuals. The lots of individuals from five of aquaria were not fed, the other received food during the entire period of the experiment. Two aquaria were occupied by fed control individuals and starved control individuals respectively. Each 7 days, the water was replaced and manganese concentration of 75 µg/ml adjusted. We used a static water system with continuous aeration, 21±2.15 °C temperature, pH 8.23±0.24, total hardness 1.75 nmol/ in CaCO<sub>3</sub>, the amount of dissolved oxygen was 6 mg/l. The metal ions content was measured by atomic absorption spectrometry, using a Perkin Elmer AAS 600 graphite oven spectrometer. For manganese ions analysis, manganese solutions of 0 – 40 µg/l concentration were prepared. The following parameters were chosen: λ = 279.5 nm, slot=0.7 nm; two matrix modifiers: Pd and Mg(NO<sub>3</sub>); atomization temperature: 1900° C, atomization period: 3 sec. For each sample three replicates were made, repeating the test until standard deviation was lower than 3%. Liver and intestine fragments were removed after 7, 14, 21, and 28 days of exposure to the metal, then fixed in buffered formaldehyde 24 hours long and further prepared for light microscopy and Hematoxylin-Eosin stained. The tissues samples were investigated by Olympus CX 40 light microscope. Weight, standard length, total length were recorded for all individuals

## RESULTS AND DISCUSSIONS

Water samples analysis showed that manganese content ranged between 61 and 80 µg/ml. Chronic exposure of fish to 75µg/L manganese caused no mortality. No weight, standard and total length changes occurred, excepting the lots of fed and starved individuals exposed for 28 days to the pollutant. This group showed an insignificant weight decrease as Javed (2006) noticed during nickel and manganese intoxication.

The main tissues that stored manganese were the intestine (42%) and liver (24.61%). It is known that liver plays an important role in metal sequestration because heavy metals may be bound to metallothioneins or other ligands (Dallinger, 1995). Manganese accumulation was higher in starved individual compared to the fed individuals (Fig.1). A higher metabolism of fed fish and a great efficiency in metal remove could explain the results. Our hypothesis is supported by a slow raise in liver metal concentration after 7 day exposure followed by a decrease due to detoxification processes (Fig. 2) .




**Fig. 1.** Intestine manganese concentrations in fed (red line) and starved individuals (blue line)




**Fig. 2.** Liver manganese concentrations in starved individuals

The digestive tract and its associated glands are the first organs impaired by food deprivation (Theilacker, 1978). Following 7 day exposure to manganese higher vacuolization of epithelial cells and epithelial lifting were noticed in starved individuals (Fig. 3B). In both fed and starved fish following 14 day exposure to the pollutant, epithelial layer disorganization in the apical region of intestinal villi, cellular and nuclear hypertrophy, flattening towards the tip of the villi appeared (Fig. 3 C, D).

The 28 day manganese intoxication induced intestinal villi fusion in order to decrease the absorption surface and to reduce the pollutant absorption (Fig. 4C, D). Additionally epithelial layer multiplied atypically (Figure 4 B). Reduced nutrient supply in the gut lumen is responsible for significant histopathological changes: mucosal atrophy (McLesse-Moon, 1989), absorptive surface reductions by regression or distortion of villi or microvilli (Gas-Noailac-Depeyre, 1976, Segner et al., 1987; Hall-Bellwood, 1995), reduction of mucosal thickness due to decrease heights of epithelial cells (Hall-Bellwood, 1995), widening of intercellular spaces and loosening of cell contacts (O'Connell, 1976; Theilacker, 1978; Peters, 1982; Nonotte et al, 1986; Segner et al., 1987), cell dissociation and disruption in mucosa (Uriarte-Balbotin, 1987) vacuolization (Peters, 1982; Eckman, 1985; Segner et al., 1987), autophagic process (Gas-Noailac-Depeyre, 1976; Peters, 1982; Segner et al., 1987), villi flattening, hypertrophy of epithelial cells,

swelling or oedema of *lamina propria*, and fusion of villi due to excessive hypertrophy, flattening and rupture of villi at tip (Bhatnagar et al., 2007). The liver is known as an integrator of physiological and biochemical functions and alterations in its structure is expected under toxic circumstances (Hinton-Lauren, 1990). After 14 day exposure to the pollutant rare hypertrophied nuclei of hepatocytes were observed (Fig. 5 B) and macrophages aggregates appeared (Fig. 5A). Additionally after 21 days of chronic intoxication intercellular corridors and sinusoid capillaries dilatation were noticed. No other liver histological changes appeared after 21 days (Fig. 5B).




**Fig. 3** *Carassius auratus auratus* -Intestine sections; A-Control (H-E, x 20); B-7 day exposure to manganese of starved individuals; cytoplasm vacuolization and hyperactivity of goblet cells, intestinal epithelium lifting in the apical region (arrow) (H-E, x100); C-14 day exposure of starved individuals (H-E; x20); D – 14 day exposure of fed individuals flattening at the tip of villi; nuclear hypertrophy; E, F-21 day exposure to manganese; severe disorganization of intestinal epithelium both in fed and starved fish






**Fig. 4** *Carassius auratus auratus* -Intestine sections; 28 day exposure to the pollutant of the fed fish A-intestinal epithelium rupture, hyperactive goblet cells, blood vessels dilatated at the tip of villi (x100); B, flattening and dilatation of the apical area, goblet cells hypersecretion (x 100); C –fusion of intestinal villi (x 100); D- starved fish, intestine villi fusion (x 100)




**Fig. 5** *Carassius auratus auratus*; Liver sections; A- Control Liver (x 40); B 14 day intoxicated starved fish (x100); nuclear hypertrophy, large corridors between hepatocytes




**Fig. 6.** *Carassius auratus auratus*; Liver sections; A- 14 day intoxication of fed fish ; macrophages aggregates (arrow) (x100; H-E); B- 21 day intoxicated starved fish (x 100)

## CONCLUSIONS

1. Nutrition status, fish size and growth rate as well as tissue heavy metal concentration are important data for monitoring assessment.
2. No significant weight, standard and total length changes occurred, excepting the lots of fed and starved individuals exposed for 28 days to the pollutant where an insignificant weight decrease occurred.
3. Manganese concentration was higher in intestine compared to liver.
4. Intestine manganese concentration in fed fish was lower than in starved fish.



5. Both intestine and liver histological changes appeared due to manganese exposure and starvation. Both stress factors induced modifications to diminish absorption surface of the intestine.

## REFERENCES

1. Bhatnagar C., Bhatnagar M., Bhag C.R., Udaipur B.C.R. - Fluoride-induced histopathological changes in gill, kidney and intestine of freshwater teleost, *Labeo rohita*, Research report Fluoride, 2007, 40 (1) , 55-61.
2. Dallinger R. - Metabolism and toxicity of metals: metallothioneins and metal elimination, pp. 171-190, In: Cell biology in Environmental Toxicology. (ed Cajaraville M.P.), Univ. del Pais Vasco, Bilbao, 1995.
3. Eckmann R. - Histopathological alterations in the intestine of whitefish (*Coregonus* sp.) larvae reared on zooplankton from Lake Constance, Dis. Aquat. Org., 1985, 1 ,11-17.
4. Gas N., Noaillac-Depeyre J. - Studies on intestinal epithelium involution during prolonged fasting, J. Ultrast. Res., 1976, 56,137-151.
5. Hall K.C., Bellwood D.R. - Histological effects of cyanide, stress and starvation on the intestinal mucosa of *Pomacentrus coelestis*, a marine aquarium fish species, J.Fish.Biol., 1995, 47, 438-454.
6. Hinton D.E., Lauren D.J. - Liver structural alterations accompanying chronic toxicity in fishes: potential biomarkers of exposure, pp. 17-57, In: Biomarkers of Environmental Contamination (ed. J.F McCarthy/L.R.Shugart), Florida, Lewis, 1990.
7. Javed M. - Studies on growth responses of fish during chronic exposures of nickel and manganese, Pakistan Journal of Biological Sciences, 2006, 9 (2), 318-322.
8. McLesse, J.M, Moon T.W. - Seasonal changes in the intestinal mucosa of winter flounder, *Pseudopleuronectes americanus* (Walbaum) from Passamaquoddy Bay, New Brunswick, J.Fish Biol., 1989, 35, 381-393.
9. Nonotte L., Nonotte G., Leray G. - Morphological changes in the middle intestine of the rainbow trout, *Salmo gairdneri*, induced by a hiperosmotic environment, Cell Tiss Res., 1986, 243, 619-628.
10. O'Connell C.P. - Histological criteria for diagnosing the starving conditions in early post yolk sac larvae of the northern anchovy, *Engraulis mordax* Girard, J.Exp. Mar. Biol. Ecol., 1976, 25, 285-312.
11. Peters G. - The effect of stress on the stomach of the European eel, *Anguilla Anguilla*, L. J.Fish Biol., 1982, 21, 497-512.
12. Segner H., Burkhardt P., Avila E.M., Juario J.,V., Storch V. - Nutrition-related histopathology of the intestine of milkfish *Chanos chanos* fry, Dis. Aquat.Org., 1987, 2, 99-107.
13. Theilacker G.H. - Effect of starvation on the histological and morphological characteristics of jack mackerel, *Trachurus symmetricus* larvae, Fish. Bull. (US), 1978, 76, 403-414.
14. Uriarte I, Balbotin F. - Caracterizacion del estado de hambruna en las larvas de sardina *Sardinops sagax musca* (Pisces, Clupeiformes) mediante criterios morfometricos y histologicos, Rev. Bio. Mar. (Valparaiso), 1987, 23 , 77-106.

15. \*\*\*European Commision, Health and Consumer Protection Directorate General, Report of the Scientific Committee for Animal Nutrition on the use of manganomanganic oxide in Feedingstuffs, 2002.

# ACUTE ELECTROPHYSIOLOGICAL EFFECTS OF A METAL AND A NON-METAL MITOCHONDRIAL TOXIN

Szabolcs TAKÁCS \*, Sarolta BANKÓ, Andras PAPP

Department of Public Health, Faculty of Medicine. University of Szeged, Dóm tér 10. H-6720 Szeged, Hungary

\* E-mail : takszabi@gmail.com

## ABSTRACT

*Mitochondrial toxins are related both to causation and modelling of central nervous damage. Manganese, often present in occupational settings and as environmental pollutant, inhibits complex 2 and 3 of the mitochondria and is known to cause a Parkinson-like CNS damage. 3-nitropropionic acid (3NP), a neurotoxin of plant and microfungal origin, inhibits succinate dehydrogenase leading to various biochemical and morphological alterations in the brain. Experimental animals treated by 3NP are used to model Huntington's disease. In this work, adult male Wistar rats were acutely prepared for recording spontaneous and stimulus-evoked activity from the projection area of the whiskers and the tail. It was tested whether the dependence of the evoked response on the parameters of stimulation was influenced by the mitochondrial toxins and what correlation existed between the cortical and peripheral responses. Mn intensified frequency-dependent fatigue. 3-NP caused latency decrease with minimal amplitude change, and had less effect on frequency dependence. The observed effects were apparently limited to the brain.*

**Key words:** manganese, 3-nitropropionic acid, evoked activity, nerve action potential, stimulation parameters

## INTRODUCTION

Mitochondrial toxins diminish oxidative energy production by blocking the mitochondrial enzyme complexes. This causes a decline in the cells' reserve energy that can result in acute symptoms and contribute to various diseases.

Manganese is essential for living organisms in small amounts but toxic when overdosed. The presence of Mn in the general environment is partly due to human activity (petrol additive, fungicide, dry cells etc). Inorganic Mn can deposit in the brain (Aschner et al., 1999), causing functional (Shinotoh et al., 1997) and structural (Yamada et al., 1986) damage. The toxin 3-nitropropionic acid (3NP) is naturally present in leguminous plants, occasionally poisoning grazing livestock. Human poisoning may come from consumption of foodstuffs infested with certain moulds producing 3NP (Brouillet et al., 1999). A common effect of both Mn and 3NP is the inhibition of the enzyme complexes responsible for energy production. Mn decreases the activity of all complexes, particularly complex 2 and 3 (Zhang et al. 2003) whereas 3NP irreversibly inhibits succinate dehydrogenase in complex 2 (Coles et al., 1979).

Although Mn and 3-NP both are mitochondrial blockers, there are many differences in their intra- and extra-mitochondrial mode of action which leads to completely dissimilar clinical pictures in chronic intoxications. The symptoms of excess Mn intake are, on the whole, called manganism, a state that can be very similar to Parkinson's disease, while 3NP poisoning resembles Huntington's disease and is used in an animal model of that (Brouillet et al., 1999).

Since the function of the nervous system requires lots of energy, the mitochondrial damage is probably reflected in the electrical activity of the brain. Beside damage in energy production, the two substances studied have also more direct effects in the nervous system, so their excessive presence should be detectable by electrophysiology. To study this, we examined the changes of evoked activity in acute experiments in rats.

## MATERIALS AND METHODS

Adult male Wistar rats (ca. 300 g) were given urethane (1000 mg/kg b.w.) ip. for anaesthesia. The left hemisphere was exposed by removing a part of the bony skull. Following a short recovery period, silver recording electrodes were placed on the surface of the cortex at the primary somatosensory projection area of the tail and the whiskers for measuring electrical activity. Evoked potentials (EPs) were triggered by 0.05 ms square electric stimuli that were delivered by pairs of needle electrodes placed in the contralateral whisker pad and in the tail base. To record the peripheral nerve responses another pair of needles were inserted in the tail 50 mm distally from the base. See Lukács and Szabó, 2007, for the recording techniques.

First, in 8 rats for both substances, the dependence of the obtained cortical EP on parameters of the whisker stimulation was tested. The just-supramaximal stimulus strength (around 3 - 5 V) was determined first, then trains of 50 stimuli were given at frequency of 1, 2, 5 and 10 Hz. This sequence was done 3 times with 30 min intervals for control. Then the appropriate solution was given in an intraperitoneal injection, and further 4 records were taken. 3NP was given in a dose of 20 mg/kg, or  $\text{Mn}^{2+}$  (in form of  $\text{MnCl}_2$ ) in 50 mg/kg, both dissolved in distilled water. In another experiment both whiskers and tail of 8 rats was stimulated at 1 Hz. Most conditions were the same as in the previous set except that beside the corresponding cortical EPs, the tail nerve response was recorded also. The records were averaged, and their onset latency and peak-to-peak amplitude was measured.

To eliminate individual variation, the latency and amplitude values were normalized to the mean of the control period (first 3 series). From these normalized data, group mean was calculated and plotted against the frequency (1 to 10 Hz) of stimulation, and any difference between the resulting curves before vs. after administration of the toxicant was sought for. In the second experiment, cortical and peripheral response parameters were plotted in the same graph to see their similar or dissimilar time trend, and their correlation was tested.

## RESULTS AND DISCUSSION

Before application of either Mn or 3NP, or in the untreated parallel controls, the latency data normalized to control mean and plotted against frequency of stimulation gave a horizontal line, showing that stimulation with 10 Hz vs. 1 Hz had no effect on the latency (Fig. 1). After injection of Mn, the latency started to increase and developed nonlinear frequency dependence with maximum at 2 Hz. Treatment with 3NP induced latency decrease which increased in time (curve 4 to curve 7) and diminished with increasing stimulation frequency.

The time trend of the latency and amplitude of EPs obtained by whisker or tail base stimulation was investigated together and was compared to the trend of the peripheral response (nerve action potential) evoked by tail stimulation.




**Fig. 1.** Frequency dependence of the EP latency before and after administration of Mn (left) and 3NP (right). Curves 1 to 3 show values before, and 4 to 7 after, application. Group means of normalized individual data (see Methods).

In the Mn-treated rats, the latency of the cortical EPs evoked by whisker or tail stimulation had a clear decreasing trend (Fig. 2, top left), while the amplitude of the EPs increased (Fig. 2, top right), but neither parameter of the tail nerve response had a noteworthy change. The correlation diagrams of Fig. 2 show that the two cortical responses changed together but independently of the peripheral response.





**Fig. 2.** Top: Relative change (compared to control mean) of the latency (left) and amplitude (right) of the cortical and peripheral evoked responses in Mn-treated rats. Record N° 1 to 3 was taken before, and 4 to 7 after, Mn application. Bottom: correlation diagrams of the same data sets. Inserts: equations and correlation coefficients of the fitted straight lines.

In the rats treated with 3NP the effect was highly similar. Latency decrease and amplitude increase of the cortical responses evolved in parallel, shown by the linear trend line standing at about 45 degrees inclination; but there was no change in the tail nerve response, as shown by the nearly horizontal trend lines. (Fig. 3).



Mitochondrial inhibition is the obvious common point in the action of Mn and 3NP, and the cortical activity changes in inherited mitochondrial encephalopathy are known (Smith and Harding, 1993). The correlation diagrams suggest, however, that a cortex- or at least CNS-specific mechanism played a role, most probably the disturbance of glutamate turnover (Mn: Hazell and Norenberg, 1997; 3NP: Tavares et al., 2001). Electrophysiological recording may be of use in the study of mitochondrial toxins.

## REFERENCES

1. Aschner M., Vrana K.E., Zheng W. : Manganese uptake and distribution in the central nervous system (CNS), *NeuroToxicol*, 1999, 20, 173-180.
2. Brouillet E., Conde F., Beal M.F., Hantraye P. : Replicating Huntington's disease phenotype in experimental animals, *Prog. Neurobiol.*, 1999, 59, 427-468.
3. Coles C.J., Edmondson D.E., Singer T.P. : Inactivation of succinate dehydrogenase by 3-nitropropionate, *J. Biol. Chem*, 1979, 254, 5161-5167.
4. Hazell A.S., Norenberg M.D. : Manganese decreased glutamate uptake in cultured astrocytes, *Neurochemistry*, 1997, 22, 1443-1447.
5. Lukács A., Szabó A.: Functional neurotoxic effects of heavy metal combinations given to rats during pre- and postnatal development, *Centr. Eur. J. Occup. Environ. Med.*, 2007, 13, 299-304.
6. Shinotoh H., Snow B.J., Chu N.S., Huang C.C., Lu C.S., Lee C., Takahashi H., Calne D.B. : MRI and PET studies of manganese intoxicated monkeys, *Neurology* 1997, 45, 1199-1204.
7. Smith S.J., Harding A.E. : EEG and evoked potential findings in mitochondrial myopathies. *J. Neurol.* 1993, 240, 367-372.
8. Tavares R. G., Santos C. E., Tasca C. I., Wajner M., Souza D. O., Dutra-Filho C.S. : Inhibition of glutamate uptake into synaptic vesicles from rat brain by 3-nitropropionic acid in vitro, *Exp. Neurol.*, 2001, 172, 250-254.
9. Yamada M., Ohno S., Okayasu I., Okeda R., Hatakeyama S., Watanabene H., Ushio K., Tsukagoshi H. : Chronic manganese poisoning: a neuropathological study with determination of manganese distribution, *Acta Neuropathol.*, 1986, 70, 273-278.
10. Zhang S., Zhou Z., Fu J. : Effect of manganese chloride exposure on liver and brain mitochondria function in rats, *Environ. Res.*, 2003, 93, 149-157.

# IMPACT OF SOME MINERAL ELEMENTS FROM INDUSTRIAL WASTE ON WHEAT PLANTS

Lidia TAUBERT <sup>1\*</sup>, Hortensia RĂDULESCU <sup>2</sup>, Ecaterina PRINCZ <sup>1</sup>,  
Éva STEFANOVITS-BÁNYAI <sup>3</sup>, Sandor KISS <sup>4</sup>

1\*. Department of Inorganic Chemistry, Chemistry Institute of the Romanian Academy, 24, Mihai Viteazu Blvd., RO- 300223 Timisoara, Roumania; 2. Department of Ecology and Environment, Faculty of Agriculture, University of Agricultural Sciences and Veterinary Medicine of Banat 119, Calea Aradului, RO-300645 Timisoara, Roumania; 3. Department of Applied Chemistry, Faculty of Food Science, Corvinus University of Budapest, 29-43, Villányi Street, H-1118 Budapest, Hungary; 4. Hungarian Magnesium Society, 73A/2, Fő fasor, H-6726 Szeged, Hungary

\* E-mail : lidiat@acad-icht.tm.edu.ro

## ABSTRACT

*The present paper follows the cultivation of green wheat plants in low acid soils, which were previously treated by different doses of mineral waste from the magnesium products industry. The two experimented industrial addings, one from the industrial process (A) and the second resulted as crusts deposited on the equipment walls (B), can be used in certain doses as soil amendment for low fertile acid soil and fertilizer for plant nutrition. The presence of magnesium and calcium in the waste composition represents a source of macroelements for plant nutrition. The enhance of the trace elements in soil is representative for iron, manganese and zinc. At harvest time, the size of green wheat plants treated with addings doses was taller comparative to the control alternative. The nitrogen supplementation increases their dry weight and dry matter, especially for the alternatives with waste B.*

**Key words:** mineral supplementation, waste, soil, wheat plant, essential and trace elements

## INTRODUCTION

At all times, the life of human beings was connected with the soil in which grow and develop plants. The soil composition has a decisive influence on the yield and the crops quality. Therefore, it is obvious that the mineral deficiencies of soil can generate diseases of animals and men. Besides the geochemical problems existing in certain areas, the man intervention could also contribute to the disturbance of nutritive elements equilibrium, even without intention; for example, by inadequate fertilization is possible to attend lacks of mineral elements in soil content (Kiss, 1978).

In this context, extensive studies had put into evidence the influence of mineral elements, magnesium especially, on the wheat characteristics and fall resistance of plant stalk (Kiss, 1984).

The magnesium products industry, magnesium carbonate and oxide obtained by carbon dioxide leaching of dolomites, generates a waste with an important mineral content. The composition of this waste includes calcium carbonate, precipitated magnesium carbonate (in ratio 3:1) and as impurities certain compounds of iron,

manganese, copper, zinc (Radulescu et al., 2006; Radulescu et al, 2007; Taubert, 2001; Taubert, 2002). This waste was tested in order to be valued in agriculture as soil amendment and tested for green oat cultivation (Taubert et al., 2008).

The present paper follows the cultivation of green wheat plants in low acid soils, which were previously treated by different waste doses, without or with nitrogen contribution.

## MATERIALS AND METHODS

The soil of luvosoil type, having a  $\text{pH}_{\text{H}_2\text{O}}$  of 6.85 and  $\text{pH}_{\text{KCl}}$  of 5.60 and a rather low fertility was collected, air-dried, crushed, mixed and put into pots, each containing 1 kilogram soil. Two types of addings were experimented, one from the industrial process (A) and the second resulted as crusts deposited on the equipment walls (B), having the composition presented in Table 1. The soil was treated with the two types of industrial addings in different amounts.

**Table 1:** Composition of the two experimented industrial addings

| Specification | Waste A | Crusts B |
|---------------|---------|----------|
| Ca, %         | 28      | 19       |
| Mg, %         | 7       | 14       |
| Na, %         | 0.77    |          |
| Fe, mg/kg     | 1850    | 880      |
| Cu, mg/kg     | 1.9     | 51       |
| Mn, mg/kg     | 136     | 51       |
| Zn, mg/kg     | 2.6     | 50       |

The experimental alternatives pursued by this research consist of four different doses for each adding ( $A_1$ - $A_4$ ,  $B_1$ - $B_4$ ) and also a control alternative ( $C_0$ ) representing untreated soil.

**Table 2:** Description of the experimental alternatives

| Experimental alternative |       | N contr., mg/kg | Mineral supplement / soil |           |           |           |           |                      |                      |                      |
|--------------------------|-------|-----------------|---------------------------|-----------|-----------|-----------|-----------|----------------------|----------------------|----------------------|
|                          |       |                 | Dose, mg/kg               | Ca, mg/kg | Mg, mg/kg | Na, mg/kg | Fe, mg/kg | Mn, $\mu\text{g/kg}$ | Zn, $\mu\text{g/kg}$ | Cu, $\mu\text{g/kg}$ |
| $A_1$                    | R     | -               | 179                       | 50        | 13        | 1.38      | 0.33      | 24.3                 | 0.47                 | 0.34                 |
|                          | $R_N$ | 134             | 179                       | 50        | 13        | 1.38      | 0.33      | 24.3                 | 0.47                 | 0.34                 |
| $A_2$                    | R     | -               | 357                       | 100       | 25        | 2.76      | 0.66      | 48.7                 | 0.93                 | 0.68                 |
|                          | $R_N$ | 134             | 357                       | 100       | 25        | 2.76      | 0.66      | 48.7                 | 0.93                 | 0.68                 |
| $A_3$                    | R     | -               | 714                       | 200       | 50        | 5.51      | 1.32      | 97.4                 | 1.86                 | 1.36                 |
|                          | $R_N$ | 134             | 714                       | 200       | 50        | 5.51      | 1.32      | 97.4                 | 1.86                 | 1.36                 |
| $A_4$                    | R     | -               | 1429                      | 400       | 100       | 11.02     | 2.64      | 194.7                | 3.72                 | 2.72                 |
|                          | $R_N$ | 134             | 1429                      | 400       | 100       | 11.02     | 2.64      | 194.7                | 3.72                 | 2.72                 |
| $B_1$                    | R     | -               | 263                       | 50        | 37        |           | 0.23      | 13.4                 | 13.2                 | 13.4                 |
|                          | $R_N$ | 134             | 263                       | 50        | 37        |           | 0.23      | 13.4                 | 13.2                 | 13.4                 |
| $B_2$                    | R     | -               | 526                       | 100       | 74        |           | 0.46      | 26.8                 | 26.4                 | 26.8                 |
|                          | $R_N$ | 134             | 526                       | 100       | 74        |           | 0.46      | 26.8                 | 26.4                 | 26.8                 |
| $B_3$                    | R     | -               | 1053                      | 200       | 147       |           | 0.93      | 53.6                 | 52.6                 | 53.6                 |
|                          | $R_N$ | 134             | 1053                      | 200       | 147       |           | 0.93      | 53.6                 | 52.6                 | 53.6                 |
| $B_4$                    | R     | -               | 2105                      | 400       | 295       |           | 1.85      | 107.3                | 105.2                | 107.3                |
|                          | $R_N$ | 134             | 2105                      | 400       | 295       |           | 1.85      | 107.3                | 105.2                | 107.3                |



All the experimental alternatives took place in two replicates, one being without nitrogen treatment (R) and the other (R<sub>N</sub>) - treated with 134mg N/kg soil as ammonium nitrate (see Table 2).

All the pots were sown with thirty wheat grains. The vegetation period was that of green plant, pursued for 6 weeks. Along the vegetation period, some morphological parameters, like number of risen plants, plant size, fresh and dry plant weight were pursued. At harvest time, soil samples were collected in order to establish the impact of waste treatment on soil fertility. Essential and trace elements were analyzed; the metal element content in soil at harvest time was established by AAS - ICP method.

## RESULTS AND DISCUSSIONS

The impact of soil treatment by different waste doses on the macroelements content is shown in Table 3.

**Table 3:** Impact of waste treatment on soil macroelements content

| Experimental alternative |                | Ca content |     | Mg content |     | K content |     | Na content |     |
|--------------------------|----------------|------------|-----|------------|-----|-----------|-----|------------|-----|
|                          |                | g/kg       | %   | g/kg       | %   | g/kg      | %   | g/kg       | %   |
| C <sub>0</sub>           | R              | 2.434      | 100 | 0.418      | 100 | 0.125     | 100 | 0.149      | 100 |
|                          | R <sub>N</sub> | 2.295      | 100 | 0.381      | 100 | 0.091     | 100 | 0.145      | 100 |
| A <sub>1</sub>           | R              | 2.475      | 102 | 0.422      | 101 | 0.115     | 92  | 0.138      | 93  |
|                          | R <sub>N</sub> | 2.390      | 104 | 0.401      | 105 | 0.095     | 104 | 0.135      | 93  |
| A <sub>2</sub>           | R              | 2.572      | 106 | 0.436      | 104 | 0.114     | 91  | 0.136      | 91  |
|                          | R <sub>N</sub> | 2.476      | 108 | 0.418      | 110 | 0.088     | 97  | 0.141      | 97  |
| A <sub>3</sub>           | R              | 2.111      | 87  | 0.365      | 87  | 0.097     | 78  | 0.132      | 89  |
|                          | R <sub>N</sub> | 2.151      | 94  | 0.339      | 89  | 0.079     | 87  | 0.137      | 92  |
| A <sub>4</sub>           | R              | 2.126      | 87  | 0.367      | 88  | 0.081     | 65  | 0.122      | 82  |
|                          | R <sub>N</sub> | 1.732      | 75  | 0.315      | 83  | 0.061     | 67  | 0.119      | 82  |
| B <sub>1</sub>           | R              | 1.979      | 81  | 0.346      | 83  | 0.086     | 69  | 0.096      | 64  |
|                          | R <sub>N</sub> | 1.532      | 67  | 0.284      | 75  | 0.065     | 71  | 0.087      | 60  |
| B <sub>2</sub>           | R              | 2.381      | 98  | 0.413      | 99  | 0.105     | 84  | 0.116      | 78  |
|                          | R <sub>N</sub> | 2.339      | 102 | 0.387      | 102 | 0.081     | 89  | 0.121      | 83  |
| B <sub>3</sub>           | R              | 2.451      | 101 | 0.425      | 102 | 0.101     | 81  | 0.115      | 77  |
|                          | R <sub>N</sub> | 2.319      | 101 | 0.409      | 107 | 0.090     | 99  | 0.114      | 79  |
| B <sub>4</sub>           | R              | 2.024      | 83  | 0.403      | 96  | 0.080     | 64  | 0.101      | 68  |
|                          | R <sub>N</sub> | 2.056      | 90  | 0.423      | 111 | 0.076     | 84  | 0.114      | 79  |

Analysing the calcium content, an increase was established only for the two lowest doses of waste A and B. Nitrogen contribution enriches calcium soil content once with the growth of waste A and crusts B doses.

A growth of the magnesium soil content once with the increase of the waste doses was more evident for crusts B, which contains more magnesium in its composition. The supply of nitrogen generates for both waste apports an evident increase of magnesium soil content.

The potassium soil content established for all experimental alternatives is lower as that of the control alternative. The nitrogen contribution generates an evident increase of the potassium soil content for both waste apports.

The sodium soil content is lower for all alternatives in comparison with the control alternative. The sodium values are lower for crusts B and nitrogen contribution does not enrich evidently the sodium soil content.

**Table 4:** Influence of waste type and doses on soil trace elements content

| Experimental alternative |                | Fe content |     | Mn content |     | Zn content |     | Cu content |     |
|--------------------------|----------------|------------|-----|------------|-----|------------|-----|------------|-----|
|                          |                | µg/g       | %   | µg/g       | %   | µg/g       | %   | µg/g       | %   |
| C <sub>0</sub>           | R              | 379.6      | 100 | 52.07      | 100 | 35.04      | 100 | 3.768      | 100 |
|                          | R <sub>N</sub> | 381.2      | 100 | 43.55      | 100 | 48.30      | 100 | 3.543      | 100 |
| A <sub>1</sub>           | R              | 391.8      | 103 | 54.74      | 105 | 37.15      | 106 | 3.730      | 99  |
|                          | R <sub>N</sub> | 410.8      | 108 | 45.45      | 104 | 28.76      | 60  | 3.786      | 107 |
| A <sub>2</sub>           | R              | 422.3      | 111 | 48.26      | 93  | 35.82      | 102 | 3.908      | 104 |
|                          | R <sub>N</sub> | 424.9      | 111 | 49.14      | 113 | 38.26      | 79  | 4.010      | 113 |
| A <sub>3</sub>           | R              | 393.8      | 104 | 50.60      | 97  | 30.20      | 86  | 3.871      | 103 |
|                          | R <sub>N</sub> | 384.3      | 101 | 46.75      | 107 | 40.84      | 85  | 3.750      | 106 |
| A <sub>4</sub>           | R              | 336.4      | 89  | 39.13      | 75  | 33.87      | 97  | 3.186      | 85  |
|                          | R <sub>N</sub> | 410.6      | 108 | 48.98      | 112 | 33.70      | 70  | 3.740      | 106 |
| B <sub>1</sub>           | R              | 385.9      | 102 | 63.36      | 122 | 44.47      | 127 | 3.701      | 98  |
|                          | R <sub>N</sub> | 374.0      | 98  | 42.71      | 98  | 47.16      | 98  | 3.570      | 101 |
| B <sub>2</sub>           | R              | 403.8      | 106 | 50.64      | 97  | 32.68      | 93  | 3.554      | 94  |
|                          | R <sub>N</sub> | 416.8      | 109 | 55.21      | 127 | 47.74      | 99  | 3.671      | 104 |
| B <sub>3</sub>           | R              | 422.0      | 111 | 52.51      | 101 | 30.36      | 87  | 3.738      | 99  |
|                          | R <sub>N</sub> | 395.1      | 104 | 42.84      | 98  | 35.94      | 74  | 3.358      | 95  |
| B <sub>4</sub>           | R              | 384.5      | 101 | 51.21      | 98  | 32.56      | 93  | 3.459      | 92  |
|                          | R <sub>N</sub> | 395.4      | 104 | 53.61      | 123 | 35.94      | 74  | 3.336      | 94  |

Because of their trace elements content, the soil treatment with waste A and B generates a different trace element level in comparison with the control alternative. Results are given in Table 4.

A proportional increase of the iron soil content once with the growth of the waste dose was established only for the alternatives with waste B. The apport of nitrogen generates an increase of the iron soil content only for the alternatives with waste A.

An evident increase of the manganese soil content took place only for the lowest dose of waste B. However, the supplementary apport of nitrogen increases in all cases the manganese content.

For all alternatives the zinc soil content is low. There is an exception for the lowest dose of waste B. Nitrogen contribution decreases for all alternatives the zinc soil content.

The copper content established for all the experimental alternatives is low. However, the apport of nitrogen increases the copper content in comparison with the untreated alternatives.

The effects of soil treatment with waste A and B show an evident influence on the development of green wheat plants by analysing some vegetation characteristics (Table 5). Adding waste A and B in soil, a beneficial effect was established for plant germination and a higher number of risen plants was determined. Adding nitrogen to soil, the number of risen plants increased especially for the alternatives with crusts B.

**Table 5:** Influence of addings on vegetation characteristics of wheat

| Experimental alternative |                | Risen plants |      | Size green pl. |       | Fresh weight |       | Dry weight |       | Dry matter |
|--------------------------|----------------|--------------|------|----------------|-------|--------------|-------|------------|-------|------------|
|                          |                | nb.          | %    | cm             | %     | mg/p.        | %     | mg/p.      | %     | %          |
| C <sub>0</sub>           | R              | 28           | 93.3 | 31.2           | 100   | 212          | 100   | 30.7       | 100   | 14.5       |
|                          | R <sub>N</sub> | 30           | 100  | 32.6           | 100   | 207          | 100   | 40.8       | 100   | 19.7       |
| A <sub>1</sub>           | R              | 30           | 100  | 31.6           | 101.3 | 205          | 96.7  | 33.0       | 107.5 | 16.1       |
|                          | R <sub>N</sub> | 30           | 100  | 33.2           | 101.8 | 171          | 82.6  | 42.8       | 104.9 | 25.0       |
| A <sub>2</sub>           | R              | 30           | 100  | 30.6           | 98.1  | 182          | 85.9  | 31.1       | 101.3 | 17.1       |
|                          | R <sub>N</sub> | 30           | 100  | 36.2           | 111.0 | 251          | 121.3 | 42.4       | 103.9 | 16.9       |
| A <sub>3</sub>           | R              | 28           | 93.3 | 32.3           | 103.5 | 139          | 65.6  | 32.0       | 104.2 | 23.0       |
|                          | R <sub>N</sub> | 28           | 93.3 | 38.8           | 119.0 | 241          | 116.4 | 53.3       | 103.6 | 22.1       |
| A <sub>4</sub>           | R              | 29           | 96.6 | 33.6           | 107.7 | 233          | 109.9 | 35.6       | 116.0 | 15.3       |
|                          | R <sub>N</sub> | 30           | 100  | 38.3           | 117.5 | 237          | 114.5 | 53.6       | 131.4 | 22.6       |
| B <sub>1</sub>           | R              | 29           | 96.6 | 31.2           | 100   | 163          | 78.9  | 39.1       | 127.4 | 24.0       |
|                          | R <sub>N</sub> | 29           | 96.6 | 36.6           | 112.3 | 156          | 75.4  | 51.8       | 127.0 | 33.2       |
| B <sub>2</sub>           | R              | 27           | 90.0 | 31.0           | 99.4  | 143          | 67.5  | 35.3       | 115.0 | 24.7       |
|                          | R <sub>N</sub> | 30           | 100  | 35.3           | 108.3 | 155          | 74.9  | 48.5       | 118.9 | 31.3       |
| B <sub>3</sub>           | R              | 28           | 93.3 | 32.6           | 104.5 | 152          | 71.7  | 36.3       | 118.2 | 23.9       |
|                          | R <sub>N</sub> | 30           | 100  | 34.6           | 106.1 | 169          | 81.6  | 48.5       | 118.9 | 28.7       |
| B <sub>4</sub>           | R              | 29           | 96.6 | 31.3           | 100.3 | 241          | 113.7 | 37.8       | 123.1 | 15.7       |
|                          | R <sub>N</sub> | 29           | 96.6 | 34.5           | 105.8 | 203          | 98.1  | 55.2       | 135.3 | 27.2       |

Adding waste A and B in soil, a beneficial effect was established for plant germination and a higher number of risen plants was determined. Adding nitrogen to soil, the number of risen plants increased especially for the alternatives with crusts B.

Green wheat plants grew taller once with the increase of the waste dose for both waste. Adding nitrogen an increase of the plant size was established, especially for the alternatives with waste A.

At harvest time, only the green wheat plants treated with the highest doses of waste A and B had a higher fresh weight towards the control alternatives. Nitrogen contribution improved the values of weight only for the alternatives with waste A.

Nitrogen contribution improves the values of dry weight and dry matter especially for soil treated with waste B.

## CONCLUSIONS

1. Considering the obtained results, the two experimented industrial addings can be used in certain doses as soil amendment for low fertile acid soil and fertilizer for plant nutrition.
2. The presence of magnesium and calcium in the waste composition represents a source of macroelements for plant nutrition.
3. The enhance of the trace elements in soil is representative for iron, manganese and zinc (without nitrogen contribution).
4. At harvest time, the size of green wheat plants treated with addings doses was taller comparative to the control alternative. Nitrogen contribution increase their size.
5. The nitrogen supplementation increases their dry weight and dry matter, especially for the alternatives with waste B.

## REFERENCES

1. Kiss A.S.: A buza magneziumtartalma (Magnesium content of wheat), Med. Univ. Mg. Melléklet, 1978, 11, 37-40
2. Kiss A.S.: A buza megdőlésének gátlása magnéziumkezeléstől a szárerősség mérése és a terméselemekre való hatás (Stopping of resistance to fall for wheat plant by magnesium treatment; measurement of the falling resistance and the influence on crops) Magyar Mezőgazdaság; Agrokémiái Tájékoztató (Hungarian Agriculture; Agrochemical Informations), 1984, 9(II), 27-28
3. Radulescu H., Taubert L., Kiss A. S., Stefanovits-Bányai É., Princz E.: Metal trace elements supplementation in oat plant nutrition, Proceedings of 7th International Symposium on Metal elements in Environment, Medicine and Biology, 2006, 291-294
4. Radulescu H., Taubert L., Kiss A. S., Princz E., Stefanovits-Bányai É.: Effect of an industrial chemical waste on the uptake of cations by green oat, J. Serb. Chem. Soc., 2007, 72(6), 629-633
5. Taubert L.: Utilization of calcium carbonate waste from the magnesium products industry, 12<sup>th</sup> International Conference on Chemistry and Chemical Engineering, 2001, 201-206
6. Taubert L.: Utilization of waste from the magnesium products industry, Proceedings of the 9<sup>th</sup> Symposium on Analytical and Environmental Problems, 2002, 41-45
7. Taubert L., Radulescu H., Kiss A. S., Kastori R., Princz E., Stefanovits-Bányai É.: Practical application in agriculture of the magnesium products industry waste, Chem. Listy, 2008, 102, 482-486

# LACTATE DEHYDROGENASE ACTIVITY AND HISTOPATHOLOGICAL CHANGES IN $\text{Cu}^{2+}$ INTOXICATION OF *CARASSIUS AURATUS GIBELIO* LIVER AFTER SHORT AND LONG TERM EXPOSURE

Daniela TEODORESCU <sup>1\*</sup>, Maria-Cristina MUNTEANU <sup>2</sup>, Andreea-Cristina STAICU <sup>1</sup>, Elena-Lotus MEȘTER <sup>1</sup>, Anca DINISCHIOTU <sup>2</sup>

1. Department of Animal Biology, Faculty of Biology, University of Bucharest, Splaiul Independenței, no. 91-95, Sector 5, RO-050 095, Bucharest, Roumania; 2. Molecular Biology Center, Faculty of Biology, University of Bucharest, Splaiul Independenței, no. 91-95, Sector 5, RO-050 095, Bucharest, Roumania.

\* E-mail: dana\_teodorescu06@yahoo.com

## ABSTRACT

The pathological effects of two sublethal concentrations (100  $\mu\text{g/l}$  and 250  $\mu\text{g/l}$ ) of copper ( $\text{CuSO}_4 \times 5\text{H}_2\text{O}$ ) on goldfish *Carassius auratus gibelio* were studied for short (24, 48, 72 hours) and long term exposure (7, 14, 21 days). The specific activity of LDH in liver, the first organ exposed to toxic compounds, were assayed. In both experiments we registered increased LDH specific activity compared to the control. This increase induced a hypoxic condition but also stimulated the glycolysis rate. The elevation in LDH activity may reflect a high dependence on anaerobic carbohydrate metabolism of the liver exposed to copper. Generally, the histopathological changes were related to the toxicant dose and exposure time.

**Key words:** *Carassius auratus gibelio*, copper toxicity, LDH, histopathology.

## INTRODUCTION

The aquatic environment has suffered due to the growing number of xenobiotics such as heavy metals and pesticides. Copper ( $\text{Cu}^{2+}$ ) is a trace metal, classified as an essential element for most living organisms but, in high concentrations, it can be a toxic pollutant. It reaches the aquatic environment through wet or dry deposition, mining activities, domestic and agricultural waste disposal. The toxic effect of Cu is related to its capacity for catalyzing oxidative reactions, leading to the production of reactive oxygen species (Lopes et al., 2001). These highly reactive compounds may also include tissue alterations and physiological derangement in fish (Varanka et al., 2001).

The application of environmental toxicology studies among nonmammalian vertebrates is rapidly expanding, and for aquatic systems, fish have become indicators for the evaluation of the effects of noxious compounds.

Lactate dehydrogenase is an oxydoreductase that catalyzes the oxidation of L-lactate to pyruvate with the mediation of  $\text{NAD}^+$  as hydrogen acceptor. This enzyme activity is present in all cells of the body and is invariably found only in the cytoplasm of the cells. In toxicology and clinical chemistry, LDH is widely used as marker of

organ or tissue lesions. Therefore, LDH has also been used as an indicator of hypoxic conditions in organisms and plays an important role in glycolysis (Das et al., 2004).

In this study liver histopathology and LDH enzymatic activity were investigated in *Carassius auratus gibelio* exposed to copper intoxication.

## MATERIALS AND METHODS

Freshwater goldfish *Carassius auratus gibelio* of length 13.5-16.5 cm and weight 20.0-30.0 g were obtained from Nucet Fishery Research Station and kept in three glass tanks (each of 60 l) at 20-21°C. Prior to exposure, fish were held for 15 days for acclimatization and evaluation of overall health under laboratory conditions. The water from the control and experimental tanks was changed every 3 days. For the intoxication experiments the fish were divided in three groups: one as control and the others two as experimental groups. In both experimental tanks, concentrations of 100 µg Cu<sup>2+</sup>/l or 250 µg Cu<sup>2+</sup>/l were used. In the short term exposure the fish were intoxicated for 24, 48 and 72 hours, while in the long term exposure they were intoxicated for 7, 14 and 21 days. During the experiment the fish were not fed. After the selected periods, the liver was excised and prepared for biochemical and histological analysis.

Enzymatic assay. LDH activity was determined by the rate of oxidation of NADH measured by the change in absorbance at 340 nm (Bergmeyer and Bernet, 1974).

Protein concentration was determined according the method of Lowry (1951) with bovine serum albumin (BSA) as standard.

Histological procedures. Small pieces of gills and intestine were immersed in Bouin's solution (overnight), dehydrated in ethanol, cleared in xylene and embedded in paraffin. The sections were stained with hematoxylin-eosin. The photos were made with an Olympus B061 Microscope.

## RESULTS AND DISCUSSIONS

Liver is the central organ of metabolism and acts as an organ of storage. Metabolic action by the hepatic parenchyma cells has been regarded as an important defense system against toxicants. The great susceptibility of liver to be damaged by chemical agents is presumably a consequence of its primary role in metabolism of xenobiotic substances.

Several liver enzymes are generally associated with cellular metabolic activity, which is inhibited or elevated under oxidative stress, especially, after exposure to heavy metals. LDH is used as a biochemical marker for assessing liver function (Das et al., 2004).

In our short term experiment, in the case of the first concentration (100 µg Cu<sup>2+</sup>/l) LDH specific activity increased by 72% after 24 h, by 151% after 48 h, respectively by 162% compared to control (Fig. 1A). This increase could be due the Cu<sup>2+</sup> induced mitochondrial dysfunction which has generated hypoxia and stimulated glycolysis rate. LDH acts as a pivotal enzyme between the glycolytic pathway and the tricarboxylic acid cycle. Thus, the nonavailability of oxygen and simultaneous elevation of LDH may suggest a bias towards the anaerobic glycolytic pathway in order to mitigate the energy crisis for survival. The same type of increase was recorded by Das et al. (2004) in the liver of three species of Indian major carp (*Catla catla*, *Labeo rohita* and *Cirrhinus mrigala*) due to nitrite toxicity.




**Fig. 1.** The variation of LDH specific activity in liver of *Carassius auratus gibelio* exposed to 100 µg Cu<sup>2+</sup>/l (A) and 250 µg Cu<sup>2+</sup>/l (B).

At the second concentration (250 µg Cu<sup>2+</sup>/l) the LDH enzymatic activity increased very significantly after 24 h, by 523%, compared to the control, but decreased after 48 and 72 h, by 58.6%, respectively 61.3%, compared to the first period of exposure (Fig. 1B). This decrease can be due to the copper induced oxidative stress in the fish, which could generate an oxidative damage at proteins level (Stadtman and Levine, 2001). Sastry and Gupta (1980) postulated that inhibition of LDH activity may be due to ion imbalance or to intracellular action of metal subsequent to initial cell membrane damage.

In our long term exposure in the case of the first concentration (100 µg Cu<sup>2+</sup>/l) the specific activity of LDH increased, by 103%, after 7 days, compared to the control, inducing a hypoxic condition. After 14 and especially 21 days the activity decreased, by 17.7%, respectively 52.6%, compared to 7 days intoxicated liver and as a consequence the glycolysis rate could be diminished (fig. 2A).




**Fig. 2.** The variation of LDH specific activity in liver of *Carassius auratus gibelio* exposed to 100 µg Cu<sup>2+</sup>/l (A) and 250 µg Cu<sup>2+</sup>/l (B).

In consistence with our results Mishra and Shukla (1997) reported that endosulfan (an insecticide) inhibited the LDH activity in the liver of *Clarias batrachus*,

while Elumalai et al. (2002) recorded inhibition in LDH activity in the crab *Carcinus moenas* due to exposure to Cu, Cr or a mixture of both. In the same way Sastry and Rao (1984) and Almeida et al. (2001) recorded inhibition of LDH activity in *Channa punctatus* due to exposure to mercuric chloride and in Nile tilapia *Oreochromis niloticus* due to exposure to Cd.

It can be seen that in the second concentration ( $250 \mu\text{g Cu}^{2+}/\text{l}$ ) the enzymatic activity decreased after 7 days, by 67%, compared to the control, but increased after 14 and 21 days, by 101%, respectively 568%, compared to the first period of exposure (Fig. 2B). The lower LDH specific activity after 7 and 14 days of exposure to  $250 \mu\text{g Cu}^{2+}/\text{l}$  could be explained by the marked inhibition of hexokinase and moderate one for pyruvate kinase (Lai and Blass, 1984). The important increase of this activity after 21 days treatment it is possible to be due to the high need of ATP for survival in the oxidative stress conditions.

These sublethal levels of Cu affect also the morphology and cause pathological changes in the liver, comparatively with liver control (Fig. 3C). So, in our short term exposure in the case of the second concentration ( $250 \mu\text{g Cu}^{2+}/\text{l}$ ), after 72 h of treatment, the main alterations found in the liver were: irregular-shaped nuclei, nuclear hypertrophy, or nucleus in a lateral position, close to the cell membrane. Anomalies such as cytoplasmic vacuolation and nucleus in a lateral position were also described in the siluriform *Corydoras paleatus* contaminated by organophosphate pesticides (Fanta et al., 2003). At the same concentration we observed melanomacrophage centres (MMG). Several studies have demonstrated that fish livers from contaminated areas show occasional MMG centres and granulomas (Teh et al., 1997).




**Fig. 3** - Liver **control (C)** – hepatocytes (red arrow), nucleus (green arrow), blood sinusoids (black arrows), x 20; **(A)** *Carassius auratus gibelio* liver intoxicated with  $250 \mu\text{g Cu}^{2+}/\text{l}$  (short term exposure) - **72 hours**: nuclear hypertrophy (white arrows), irregular - shaped nuclei (green arrows), biliary duct (blue arrow), melanomacrophage centres (\*), intercellular spaces (\*), x 40.



Although high prevalence of hepatic MMG centres may occur in fish collected in contaminated sites due to the storage of foreign material, they can not be linked to direct contaminant exposure since they are also involved in other metabolic functions (Stentiford et al., 2003).

In our long term exposure, at  $100\text{ }\mu\text{g Cu}^{2+}/\text{l}$ , the connective tissue around the large blood vessels increased in thickness, after 14 days of treatment. But the most severe histopathological changes induced by copper intoxication registered after 21 days of treatment in the second concentration ( $250\text{ }\mu\text{g Cu}^{2+}/\text{l}$ ). These consisted of cytoplasmic vacuolation, irregular shaped cells, cellular rupture, blood vessels congestion, which impair the normal functioning of the hepatic tissue.




**Fig.4 - (A)** *Carassius auratus gibelio* liver intoxicated with  $100\text{ }\mu\text{g Cu}^{2+}/\text{l}$  (long term exposure) – **14 days**: large thickness of connective tissue around blood vessels (black arrows), x 20; **(B)** *Carassius auratus gibelio* liver intoxicated with  $250\text{ }\mu\text{g Cu}^{2+}/\text{l}$  (long term exposure) – **21 days**: blood congestion (black arrows), large intercellular spaces (\*), x 40.

Thus, sublethal levels of copper sulphate affect the efficiency of tissue metabolites and cause pathological changes in the liver of *Carassius auratus gibelio*.

## CONCLUSIONS

1. The goldfish showed various stress responses to  $100\text{ }\mu\text{g Cu}^{2+}/\text{l}$  and  $250\text{ }\mu\text{g Cu}^{2+}/\text{l}$  exposure.
2. Taking in to account that the liver is the first organ exposed to toxic compounds, the biochemical and histological alterations produced by  $\text{Cu}^{2+}$  were studied.
3. The increase in LDH activity may reflect an increased dependence on anaerobic carbohydrate metabolism by the liver exposed to copper.
4. Our results provide evidence that LDH can be used as a sensitive indicator of aquatic pollution.

5. Generally, the histopathological changes were related to the toxicant dose and exposure time. Thus, sublethal levels of copper caused pathological modifications in the liver of *Carassius auratus gibelio*.

## REFERENCES

1. Almeida J.A., Novelli E.L.B., Silva M.D., Alves, R.: Environmental cadmium exposure and metabolic responses of the Nile tilapia, *Oreochromis niloticus*, Environ. Pollut., 2001, 114, 169-175.
2. Bergmeyer H.U., Bernt E.: Lactate dehydrogenase, p.574-579, in „Methods of Enzymatic Analysis”, Vol.2, (Edited by Hans Ulrich Bergmeyer), Acad. Press. Inc., New York, San Francisco, London, 1974.
3. Das P.C., Ayyappan S., Das B.K., Jena J.K.: Nitrite toxicity in Indian major carps: sublethal effect on selected enzymes in fingerlings of *Catla catla*, *Labeo rohita*, *Cirrhinus mrigala*, Comp. Biochem. Physiol., 2004, 138(C), 3-10.
4. Elumalai M., Antunes C., Guilhermino L.: Effects of single metals and their mixtures on selected enzymes of crab, Air Soil Pollut., 2001, 141, 273-280.
5. Fanta E., Rios F.S., Romao S., Vianna C.C., Freiburger S.: Histopathology of the fish *Corydoras paleatus* contaminated with sublethal levels of organophosphorus in water and food, Ecotox. Environ. Saf., 2003, 54, 119-130.
6. Lai J. C. K., Blass J. P.: Neurotoxic effects of copper: inhibition of glycolysis and glycolytic enzymes, Neurochemical Research, 1984, 9, 1699-1710.
7. Lopes P.A., Pinheiro T., Santos M.C., Mathias M., Colares-Pereira M.J., Viegas A.M.: Response of antioxidant enzymes in freshwater fish populations to inorganic pollutants exposure, Sci. Total Environ., 2001, 280, 153-163.
8. Lowry O.H., Rosenbrough, N.J., Farr, A.L., Randall, B.J.: J. Biol. Chem., 1951, 193, 265-275.
9. Mishra R., Shukla S.P.: Impact of endosulfan on LDH from the freshwater catfish *Clarias batrachus*, Pest. Biochem. Physiol., 1997, 57, 220-234.
10. Sastry K.V., Gupta P.K.: Alterations in the activities of a few dehydrogenases in the digestive system of 2 teleost fishes exposed to lead nitrate, Ecotox. Environ. Safe., 1980, 4, 232-239.
11. Sastry K.V., Rao D.R.: Effect of mercuric chloride on some biochemical and physiological parameters of *Channa punctatus*, Environ. Res., 1984, 34, 343-350.
12. Stadtman E.R., Levine R.L.: Protein oxidation, Ann. New York Acad. Sci., 2001, 191-208.
13. Stentiford G.D., Longshaw M., Lyons B.P., Jones G., Green M., Feist S.W.: Histopathological biomarkers in estuarine fish species for the assessment of biological effects of contaminants, Mar. Environ. Res., 2003, 55, 137-159.
14. Teh S.J., Adams S.M., Hinton D.E.: Histopathologic biomarkers in feral freshwater fish populations exposed to different types of contaminant stress, Aquat. Toxicol., 1997, 37, 51-70.
15. Varanka Z., Rojick I., Abraham M.: Biochemical and morphological changes in carp liver following exposure to  $\text{Cu}_2\text{SO}_4$  and tannic acid, Comp. Biochem. Physiol., 128, 467-478.

# K<sup>+</sup>-SELECTIVE ELECTRODE BASED ON A CALIX[6]ARENIC COMPOUND

Lidia VARVARI, Sorin-Aurel DORNEANU, Cătălin-Ionel POPESCU \*

Department of Physical Chemistry, Faculty of Chemistry and Chemical Engineering,  
Babeş-Bolyai University, Str. Arany Janos Nr. 11, RO-400 028 Cluj-Napoca, Roumania

\* E-mail: cpopescu@chem.ubbcluj.ro

## ABSTRACT

A calix[6]arenic compound (C6Cr3Am3) was tested as ionophore for cation-selective electrodes (ISE), based on PVC membrane. Potentiometric measurements were performed in Na<sup>+</sup>, K<sup>+</sup>, Ca<sup>2+</sup>, Mg<sup>2+</sup> and Ba<sup>2+</sup> standard solutions. The best response was recorded for K<sup>+</sup>: a low detection limit and large linear range were obtained.

**Key words:** calix[6]arene, PVC-based ISE, potassium selective electrode

## INTRODUCTION

Ion-selective electrodes (ISE) are well-known for their wide applications in important fields such as clinical, food and environmental chemistry (Bakker et al., 1999). For example, it was estimated that over a billion clinical analyses are performed annually in laboratories all over the world using ISE (Bakker et al., 1997).

ISE are usually made of a polymer matrix incorporating an ionophore. Macrocyclic compounds are among the most frequently used ionophores, they selectively bind different ions by entrapping them in their cavity. The selectivity of the membrane is thus strongly influenced by the size match between the guest ion and the host cavity. A very widely used class of ionophores are calixarene derivatives, due to their cup-like shape (Arora et al., 2007, McMahon et al., 2003, Ludwig 2002), as shown in figure 1.

This paper aims at testing a new calix[6]arenic compound as ionophore for cation-selective membranes. The structure of the ionophore, p-tert-butyl-tricrothyl-triamide-calix[6]arene (C6Cr3Am3), is shown in Figure 2.




**Fig. 1.** The general structure of a calix[4]arene compound




**Fig. 2.** The structure of the C6Cr3Am3 ionophore. Crothyl: -CH<sub>2</sub>-CH=CH-CH<sub>3</sub>;  
Amide: -CH<sub>2</sub>-C(O)N(CH<sub>2</sub>CH<sub>3</sub>)<sub>2</sub>

## MATERIALS AND METHODS

### Materials

The C6Cr3Am3 ionophore was kindly provided by dr. Jeanne-Elisabeth Popovici from the Institute of Chemistry "Raluca Ripan" Cluj-Napoca (Romania). All used reagents were of analytical grade. Calcium chloride, ammonium chloride, lithium acetate, 2-nitrophenyloctylether (NPOE), high molecular weight polyvinyl chloride (PVC) and tetrahydrofuran (THF) were purchased from Fluka (Buchs, Switzerland). Potassium chloride was from Riedel-de Haën (Seelze, Germany), magnesium chloride was purchased from Chimopar (Bucharest, Romania) and sodium chloride was from Merck (Darmstadt, Germany). Barium chloride was from Silal Trading (București - România).

### Membrane preparation

The investigated cation-selective membrane contained 1% (w/w) C6Cr3Am3 as ionophore, 33% (w/w) PVC as polymer matrix and 66% (w/w) NPOE as plasticizer. The membrane weighted ~0.3 g.

The ion-selective membrane was prepared according to the following procedure. The ionophore, plasticizer and polymer matrix were successively dissolved in THF, under stirring. After dissolution, the mixture was poured into a glass cylinder, in THF atmosphere, in order to avoid pores formation. The membrane was dried and stored in dark. Before use, it was conditioned for at least 24 hours in the solution containing the cation to be determined.

### Experimental setup

Measurements were performed using a PC-controlled setup (Dorneanu et al., 2005). The system control, as well as data acquisition, was performed using the LabView 5.1 software. Data treatment was done by using the Origin 5.0 software.

Two similar electrodes were prepared by fixing an 8 mm diameter disc membrane at the bottom end of a plastic syringe body. As internal reference, an Ag/AgCl system was used. The inner electrolytes contained 5 mM NaCl. Both electrodes were tested simultaneously, and each measurement was repeated at least two times in the same working conditions. A double-junction saturated calomel electrode was used as external reference. The external liquid junction was filled with CH<sub>3</sub>COOLi 0.1 M.




Fig. 3. Records of electrodes potentials during additions of standard solutions of investigated cations

## Experimental procedure

The experimental procedure consisted in “batch” potentiometric measurements, performed using addition of standard solutions of the investigated cations. In order to assess the ISE sensitivity towards  $\text{Na}^+$ ,  $\text{K}^+$ ,  $\text{Ca}^{2+}$ ,  $\text{Mg}^{2+}$  and  $\text{Ba}^{2+}$  cations, the ISE potential was recorded for three concentrations ( $10^{-4}$  M,  $10^{-3}$  M and  $10^{-2}$  M), and the sensitivity was estimated as the potential difference observed between  $10^{-3}$  M and  $10^{-2}$  M (see figure 3). Then, calibration curves were drawn only for the cation for which a quasi-nernstian response was obtained. The calibration curves, served also for evaluation of the linear domain and the detection limit.

## RESULTS AND DISCUSSIONS

### ISE sensitivities for $\text{Na}^+$ , $\text{K}^+$ , $\text{Ca}^{2+}$ , $\text{Mg}^{2+}$ and $\text{Ba}^{2+}$

As a preliminary test, the sensitivities of the electrode, based on C6Cr3Am3 ionophore, were evaluated for various cations, using the method described in the previous section. The reported values are the mean for the two measurements performed with the two different ISEs, in similar experimental conditions (Table 1). As can be observed, a quasi-nernstian response was obtained only for  $\text{K}^+$ . All other cations gave under-nernstian responses.

**Table 1:** Sensitivities for different cations recorded for C6Cr3Am3-based electrode.

| Cation           | Sensitivity (mV/decade) |
|------------------|-------------------------|
| $\text{Na}^+$    | $44.0 \pm 2.0$          |
| $\text{K}^+$     | $53.8 \pm 1.2$          |
| $\text{Ca}^{2+}$ | $8.0 \pm 0.0$           |
| $\text{Mg}^{2+}$ | $7.5 \pm 0.5$           |
| $\text{Ba}^{2+}$ | $11.5 \pm 1.0$          |




**Fig. 4.** Calibration curve recorded for  $\text{K}^+$  using C6Cr3Am3-based ISE

Based on the preliminary tests, the calibration curves were drawn for  $\text{K}^+$  cation. In order to maintain a constant ionic strength, the calibration solutions contained 1 mM NaCl. An example of calibration curves is given in figure 4. The ISE sensitivity for  $\text{K}^+$ , recalculated from the calibration curve, was 50 mV/decade, the mean detection limit was  $5 \cdot 10^{-1}$  mM, and the mean linear range was  $\sim 2.5$  decades.

### Electrodes reproducibility

Two types of reproducibility were studied: (1) inter-electrodes reproducibility, i.e., the reproducibility observed between two measurements performed with similar electrodes in the same experimental conditions; (2) inter-measurements

reproducibility, i.e., the reproducibility observed for three measurements performed with the same electrode, in the same experimental conditions. As figure 5 shows, in both cases a very good reproducibility was observed.




**Fig. 5.** Inter-electrodes and inter-measurements reproducibility. The error bars correspond to the standard deviations of the measurements.

## CONCLUSIONS

1. This study aimed at characterizing the calix[6]arene compound C6Cr3Am3 as ionophore for potentiometric ISE based on a PVC membrane.
2. A preliminary study was performed to assess the ISE sensitivities for different cations:  $\text{Na}^+$ ,  $\text{K}^+$ ,  $\text{Ca}^{2+}$ ,  $\text{Mg}^{2+}$  and  $\text{Ba}^{2+}$ . Results showed a quasi-nernstian response only for  $\text{K}^+$ .
3. Calibration curves were recorded for  $\text{K}^+$  and good electroanalytical parameters were obtained: a mean slope of 50 mV/decade of concentration, a linear range of  $\sim 2.5$  decades and a detection limit of  $5 \cdot 10^{-1}$  mM.
4. The obtained results suggest that C6CrAm3 could be used as ionophore for  $\text{K}^+$ -selective ISE. A study to evaluate the interference of cations, usually accompanying  $\text{K}^+$ , is under progress.

## REFERENCES

1. Arora V., Chawla H.M., Singh S.P.: Calixarenes as sensor materials for recognition and separation of metal ions, *ARKIVOC*, 2007, ii, 172-200
2. Bakker E., Diamond D., Lewenstam A., Pretsch E: Ion sensors: current limits and new trends, *Anal. Chim. Acta*, 1999, 393, 11-18
3. Bakker E., Buhlmann P., Pretsch E.: Carrier-Based Ion-Selective Electrodes and Bulk Optodes. 1. General Characteristics, *Chem. Rev.*, 1997, 97, 3083-3132
4. Dorneanu S.A. Coman V., Popescu I.C., Fabry P.: Computer-controlled system for ISEs automatic calibration, *Sensors and Actuators B*, 2005, 105, 521-531
5. Ludwig R., Dzung N. T. K.: Calixarene-based molecules for cation recognition, *Sensors*, 2002, 2, 397-416
6. McMahon G., O'Malley S., Nolan K.: Important calixarene derivatives – their synthesis and applications, *ARKAT USA Inc.*, 2003, vii, 23-31

# IN VIVO EFFECTS OF CIS-PLATINUM ON MAGNESIUM LEVELS IN BLOOD, MUSCLES AND HEART SAMPLES

Ariana-Bianca VELCIOV <sup>1\*</sup>, Petru-Robert VELCIOV <sup>2</sup>, Gabriela GARBAN <sup>3</sup>,  
Adina-Elena AVACOVICI <sup>2</sup>, Mirela AHMADI <sup>1</sup>, Ludovic SAYTI <sup>4</sup>

*1 Department of Biochemistry and Molecular Biology, Faculty of Food Products Technology, University of Agricultural Sciences and Veterinary Medicine, Calea Aradului Nr. 119, RO-300 645 Timișoara, Roumania; 2. Faculty of Industrial Chemistry and Environmental Engineering, University „Politehnica”, Bd. Victoriei Nr.2, RO-300 006 Timisoara, Roumania; 3. Institute of Public Health, Bd. Dr. V.Babes Nr.16, RO-300 226 Timișoara, Roumania; 4. Department of Inorganic Chemistry, Chemistry Institute of the Romanian Academy, 24, Mihai Viteazu Blvd., RO- 300223 Timisoara, Roumania;*

*\* E-mail: martauariana@yahoo.com*

## ABSTRACT

*The purpose of this study was to evaluate the concentration of magnesium in blood, muscles and heart samples during cis-platinum administration on Wistar rats. For this experiment, we used adult animals from Wistar rats strain divided in two experimental groups and one control group. Wistar rats from the experimental groups were injected intraperitoneally (i.p.) with the drug called Sin-Platinum (doses 2.0 mg/kg b.w. respectively 8.0 mg/kg b.w.), and the animals from control group were injected with saline solution (placebo). The results showed that the concentration of metal elements from blood and tissues samples is changed proportionally with the administered doses.*

*We have observed the consequences of accumulation of cis-platinum in several different samples, (in this experiment blood samples, muscles and heart tissues) and we evaluated the modifications which appeared after repeated doses simulating chemotherapeutic treatment in tumor-free rats.*

**Key words:** cis-platinum, Wistar rats, magnesium

## INTRODUCTION

Cis-platinum, as a compound was the first time described by Peyrone in 1845, and the structure was elucidated in 1895 by Alfred Werner. In 1960s, Rosenberg and his team research discovered that electrolysis products from a platinum electrode inhibited mitosis in *Escherichia coli* bacteria (Rosenberg, 1985). Cis-platinum is known as a chemotherapeutic agent often used in the treatment of cancer. Literature data show that cis-platinum administration affects the biochemical homeostasis (Velciov et al., 2007). Studies regarding the consequences of cis-platinum administration in blood serum and some tissues samples in case of experimental animals, present interest for biochemistry, nutrition, pharmacology and toxicology.

Cis-platinum (cis-diamminedichloroplatinum) abbreviated cDDp or cis-DDP is a planar complex containing a central platinum atom surrounded by two ammonia

molecules and two chloride atoms arranged in cis position (Riley and Sternson, 1985, Lippert, 1999, Threkeld, 1999). Cis-platinum mechanism of action is based on binding to DNA which determined intra-strand cross—links and formation on DNA adducts.

This drug has a broad spectrum of activity against many different solid tumors such as lung, ovary, testis, bladder and head and neck cancers and is also an effective agent in treatment of some hematological malignancies such as refractory lymphomas (Lippert, 1999; Baba, 2002). The adverse effects of c-DDP include nausea, nephrotoxicities, ototoxicities, neurotoxicities and myelosuppression.

Many nutritional problems in cancer patients are caused by alterations following chemotherapy. During cytostatic therapy the metabolism of carbohydrate, lipid and protein is disturbed. Metal elements are implicated in maintenance of acido-basic balance, osmotic and colloid-osmotic balance. Without an adequate supply of magnesium, the toxic effects cannot be prevented (Links and Lewis, 1999; Ghizdavu, 2000; Garban, 2005b).

## MATERIALS AND METHODS

For the experimental part, we used adults Wistar strain rats, maintained on pathogen-free conditions, at 22–25°C room temperature, at 55–65% relative air humidity, fed on normal rhythm and standard breeding food and water.

The animals were randomly divided in three groups: one control (C) and two experimental groups ( $E_1$  and  $E_2$ ). Each group contained 10 animals (males and females) with an average body weight (b.w.) of  $200 \pm 20$  g.

Cis-platinum used for this study was the commercially available Sin-Platin (Sindan, Romania). The animals of experimental group  $E_1$  were injected i.p. with 2.0 mg/kg b.w. Sin-Platin and with 8.0 mg/kg b.w. the animals of  $E_2$  group. The animals from C group, were injected i.p. with saline solution.

On the 15<sup>th</sup> day of the experiment, after 12 hours of fasting (overnight), and Ketanest anesthesia, the rats were killed, and blood and tissue samples from skeletal muscles and heart were removed for analyses. The blood samples were taken after laparotomy and puncture of vena cava caudalis. Magnesium concentration in blood was determined by using an analyser Hospitex Screen Master. The contents of magnesium in tissues were determined by flameless atomic absorption spectrophotometry of samples. Results are expressed as means (X) and standard deviations (SD) and to calculate the statistical significance, the *t* test was used as appropriate and *p* values.

## RESULTS AND DISCUSSIONS

Macrobioelements and trace elements have a very important role in material and energetic metabolism, and are responsible for biochemical homeostasis (Ghergariu, 1980, Garban, 1999).

The studies regarding chemotherapy are developed especially using experimental studies on laboratory animals (Ciudin și Marinescu, 1996).

Our results concerning the concentration of magnesium in blood, skeletal muscle and heart samples after cis-platinum administration in Wistar rats are presented in Table 1.



**Table 1.** Mean concentration of Mg in blood, muscles and heart samples of Wistar rats after cis-platinum administration.

| Specifications       | n  | Mg in blood<br>mg/dL | Mg in muscle<br>( $\mu$ g/g tissue) | Mg in heart<br>( $\mu$ g/g tissue) |
|----------------------|----|----------------------|-------------------------------------|------------------------------------|
|                      |    | $\bar{X} \pm SD$     | $\bar{X} \pm SD$                    | $\bar{X} \pm SD$                   |
| Group C              | 10 | 2.12 $\pm$ 0.18      | 258,28 $\pm$ 43,01                  | 235,91 $\pm$ 39,20                 |
| Group E <sub>1</sub> | 10 | 0.93 $\pm$ 0.59      | 248,56 $\pm$ 41,42                  | 227,41 $\pm$ 37,15                 |
| $\Delta \bar{X}$     |    | -1.19                | -9,72                               | -8,50                              |
| Grup E <sub>2</sub>  | 10 | 0.52 $\pm$ 0.06      | 227,38 $\pm$ 37,58*                 | 207,61 $\pm$ 33,60*                |
| $\Delta \bar{X}$     |    | -1.60                | -30,90                              | -28,30                             |

n– number of animals in case of each group

\*p< 0,01;\*\* p<0,05

From this data it is obviously that mean value of Mg in blood after administration of cis-platin are lower compared to the values of control group (Sartori, 1991) – see fig. 1.




**Fig. 1** Concentration of magnesium in blood of Wistar rats

Depressions of trace elements can be correlated with some changes of acid - base balance, osmotic and osmotic – colloidal balance (e.g. magnesium can realise stabile combinations with aminoacids).

From the obtained data we can observe that Mg concentration in experimental groups is lower as compared with the values obtained in the control

group. Magnesium plays a structural and regulatory role in living organisms (Ghergariu, 1980, Buckley et al., 1984, Seelig M., 1993, Lajer et al., 2003, Garban et al., 2005). It has an essential role in metabolism of several minerals, through he's function as an activator of all enzymatic systems. Magnesium deficiency is linked with cardiovascular, renal digestive, neurological and mostly muscular alterations (Garban, 1999) – see fig. 2.




**Fig. 2.** Concentration of magnesium in tissue samples of Wistar rats

In case of heart tissues samples magnesium values are lower in experimental groups as compared to control groups, and in case of experimental group (E<sub>4</sub>) these difference are significant ( $p < 0.01$ ).

## CONCLUSIONS

1. Studies regarding concentration of some metal elements from blood and tissue samples are very important for defining the influence of chemotherapy on normal health status.
2. In case of blood samples, we can observe a depression of magnesium levels, some even significant, in experimental groups compared to control groups.
3. Administration of cis-platinum in Wistar rats is characterized by magnesium depression in skeletal muscles indirectly proportional with administrated doses.
4. The concentration of magnesium metal elements in heart tissue are decreased after cis-platinum administration maybe because toxic effects of the metallic compound, on particularly nephrotoxic effects, which cause severe depletion magnesium.

5. Metal elements are involved in biochemical homeostasis, which is very important for the normal development of physiological processes, such as: acido-basic balance, osmotic and colloid-osmotic balance. Without an adequate supply of metal elements the toxic effects cannot be prevented.

## REFERENCES

1. Ahmadi-Vincu M., Garban G., Pup M., Garban Z., Clep C., Martău A-B. - Manganese overdose effects on some blood serum electrolytes in Wistar strain rats, 399-402, in *Simpozionul "Euroaliment" Galați*, Romania, 23-25 octombrie, 2003;
2. Baba Al. I. - *Oncologie comparată*, Ed. Academiei Române, București, 2002 ;
3. Buckley J.E., Clark V.L., Meyer T.J., Pearlman N.W. - Hypomagnesemia after cis-platin combination chemotherapy. *Arch. Intern. Med.*, 1984, 144, 2347
4. Ciudin Elena, Marinescu D. - *Animale de laborator*, Vol.1, Ed. All București, 1996;
5. Garban Z. – *Biochemistry – Comprehensive Treatise* (in romanian), Vol I, Ed. Didactică și Pedagogică, București, 1999;
6. Garban Z., Velcirov Ariana–Bianca, Velcirov P.R, Moldovan I., Avacovici Adina, Ahmadi-Vincu Mirela,. Ghibu G.D - Experimental data concerning the action of Cisplatin on metal elements in blood, "*Scientific Researches*" Universitatea "S.A.M.V.B." Timișoara, 2005;XI (1) 147-153
7. Garban Z. – *Xenobiochimie: Tratat comprehensiv*, Vol. IV, Ed. Eurobit, Timișoara, 2005b
8. Ghergariu S. - *Oligominerale și oligomineraloze*, Ed. Acad. RSR, București, 1980;
9. Ghizdavu L. – *Chimie bioanorganică*, Editura Poliam, Cluj – Napoca, 2000;
10. Lajer H., Bundgaard H., Secher N.H., Hansen H.H., Kjeldsen K., Daugaard G. - Severe intracellular magnesium and potassium depletion in patients after treatment with cisplatin, *British Journal of Cancer*, 89, 1633-1637, 2003;
11. Links M., Lewis C., - Chemoprotectants: a review of their clinical pharmacology and therapeutic efficacy, *Drugs*, 57, 293-308, 1999;
12. Lippert B. (Ed.) - *Cisplatin. Chemistry and Biochemistry of a leading anticancer drug*, Verlag Helvetica Chimica Acta, Zürich, Switzerland, 1999
13. Pollera C.F., Ameglio F., Nardi M., Marolla P., Carlini P., Frasca A.M. - Dose and schedule effects of cisplatin on the related acute iron changes. *Oncology*, 1990, 47(2), 133-138;
14. Riley, C.M., Sternson L.A. - Cisplatin, *Analytical Profiles of Drug Substances*, 14, 77, 1985;
15. Rosenberg B.- Fundamental studies with cisplatin, *Cancer*, 55, 2303-2315, 1985;
16. Seelig M. - Magnesium in oncogenesis and in anti-cancer treatment: Interaction with minerals and vitamins, pp.238-318, in „*Adjuvant Nutrition in Cancer Treatment*” (Quillan P., Williams R. M., Eds.), Publ. Cancer Treatment Research Foundation, 1993;
17. Threlkeld D.S. (Ed.) - Antineoplastics, alkylating agents, cisplatin (CDDP). 652a–2d, in „*Facts and Comparisons Drug Information*”, St. Louis, Facts and Comparisons, 1999;

18. Velciov Ariana - Bianca, Velciov Petru Robert, Garban Gabriela, Ravis A., Ahmadi Mirela - Changes of some metal elements concentration in the kidneys of laboratory animals after cis-platinum administration pp. 155-158 in *"Proceedings of the 14<sup>th</sup> Symposium on Analytical and Environmental Problems"*, Szeged, 2007

# MAGNESIUM IN DRINKING WATER OF SOME PUBLIC SYSTEMS AND WELLS IN IASI DISTRICT AND THE DAILY MAGNESIUM REQUIREMENT

Otilia VORONIUC<sup>1\*</sup>, Manuela GHEORGHEȘ<sup>2</sup>, Nicoleta FLORESCU<sup>2</sup>,

Anca CRISTEA<sup>3</sup>, Iulia CASAUX-VORONIUC<sup>4</sup>

1. Department of Preventive Medicine, Faculty of Medicine, University of Medicine and Pharmacy « Gr. T. Popa » Street of the University no. 16, RO-700115 Iași, Romania; 2. The Public Health Institute Iași; 3. The Public Health Authority Iași; 4. University of Medicine, Rouen, France

\* Email : otília\_voroniuc@yahoo.com

## ABSTRACT

Tap water was more effective in meeting magnesium requirements than dietary supplementation. The mineral macroelements concentration: Mg, Ca, Ca/Mg, fixed residuum and total hardness was measured in 62 samples of the drinking water, 38 of the public systems and 24 samples of wells in two periods of the year, July and October in 5 citis of Iasi district. The chemical analyse of the samples point out a low degree of the mineralization and fixed residuum in average below 500 mg/l, variations of the Ca, Mg, total hardness concentration. Mg average concentration was higher in the well water of Tg. Frumos and Hirlau city and lower in depth water of the city Pascani and Tg. Frumos; the mean concentration of the total hardness in drinking water of the public system, was ranged between 12.02 – 16.28 °G, average was higher in the well water of the four citys. Ca/Mg ratio was higher in the drinking water of the public system with variation between 2,91- 16,84. In our study, in 5 citis with an total of 388.650 inhabitants, the Mg content of drinking water may provide below 10% of the daily Mg requirement in public system and between 10,78 % – 48,5 % in wells, with importants variations depending on the soil.

**Key words:** magnesium, daily Mg requirement, Ca/Mg, total hardness, drinking waters

## INTRODUCTION

Several previous epidemiological studies have shown a relation between drinking water quality and death in cardiovascular disease (Rylander, Arnaud, 2004).

Many studies support a statistic correlation between a reduced level of the magnesium and total hardness in the drinking water and the higher cardiovascular diseases mortality (Alexa L., 1996, Cuciureanu R., 2002, Gavăt V., 2007, Ionuț C., 2004, Nebrand C., 2003).

It is possible, then, that waterborne magnesium could correct an insufficient dietary magnesium level.

Magnesium - rich water may contribute to coverage of magnesium requirements by providing significant amounts of natural, energy-free, bioavailable magnesium. Waterborne magnesium could correct an insufficient dietary magnesium level (Durlach, M.J., 1985, Rylander, Arnaud, 2004, Sabatier M., 2002, Seeling M.S., Rosanoff A., 2003)

The aims of the study were to analyse the Mg, Ca, Ca/Mg and total hardness in the drinking water of the towns areas of Iasi county, for evaluate their health hazard. In the same time, were to realise an useful presentation for the therapy of the mean Mg concentration depending of a person's daily magnesium requirement.

## MATERIALS AND METHODS

The mineral macroelements concentration: Mg, Ca, Ca/Mg, fixed residuum and total hardness was measured in 62 samples of the drinking water, 38 of the public systems and 24 samples of wells in two periods of the year, July and October in Iași, Pâncani, Hirleu, Tg. Frumos and Podu Iloaie (*with an total of 388. 650 inhabitants*) areas of Iasi district.

The calculus of probability to determinate the size of the water sample in the four towns was used. On used the standardisation methods.

## RESULTS AND DISCUSSIONS

### Magnesium concentration in drinking water

In the drinking water of the public system of the 5 citis magnesium average concentration was higher at Hirleu (mean 23,74 mg/l in july and 28,75 mg/l in octobre) and in order at Podu Iloaie city (mean 27,18 mg/l on july and 14,33 mg/l in octobre), Iasi city (mean 19,25 mg/l in surface water Prut and 9,46 mg/l in groundwater Timisesti), at Pâncani city (mean 12,23 mg/l on july and 13,52 mg/l on octobre), at Tg. Frumos (mean 10,28 mg/l on july and 5,36 mg/l on octobre). Results are given in Table 1.

In the drinking water of the wells, of the 5 citis, magnesium average concentration was higher in well water of Tg. Frumos, Podu Iloaie and Hirleu city and lower in wells water at the Pâncani city (table 1).

It is advisable to have 30 mg/liter of Mg in drinking water (Durlach, 1989).

**Table 1.** The mean Mg concentration (mg/l) in the samples of the drinking water of the 5 cities in the public system and wells

| Town        | Public System                                                            |         | Wells  |         |
|-------------|--------------------------------------------------------------------------|---------|--------|---------|
|             | july                                                                     | octobre | july   | octobre |
| Iasi        | 19,25 mg/l in surface water Prut<br>9,46 mg/l in grandwater<br>Timisesti |         | -      | -       |
| Pâncani     | 12,23                                                                    | 13,52   | 37,93  | 26,6    |
| Hirleu      | 23,74                                                                    | 28,75   | 94,36  | 74,5    |
| Tg. Frumos  | 10,28                                                                    | 5,36    | 102,95 | 101,56  |
| Podu Iloaie | 27,18                                                                    | 14,33   | 94     | 98      |

In some geographic areas, the magnesium content of drinking water may provide 20% to 40% of a person's daily magnesium requirement (Sabatier, 2002).

In our study the Mg content of drinking water may provide below 10% of a person's daily magnesium requirement from the water of the public system (with variations between 9,62 – 2,17) (fig 1) and between 10,78 % – 48,5 % from the water

of wells, with important variations depending on the soil for the depth water in different cities (fig 2).




**Fig.1.** Percent provide of the person's daily magnesium requirement (370 mg/day) through drinking water of public system




**Fig. 2.** Percent provide of the person's daily magnesium requirement (370 mg/day) through drinking water of wells

For example, a 1and1/2 liter of water with a magnesium content of 100 mg per liter contains 40,54 % of the daily magnesium requirement of 370 mg per day. However, a liter of water that is low in magnesium (eg, <10 mg per liter) provides less than 3% of the daily requirement.

In a previous study on rural area of Iasi county, (430.402 inhabitants in 2007) the results showed that  $Mg^{++}$  in wells water was the only mineral which proved to have a protective effect against cardio-vascular diseases; the interrelation between the ratio of Mg and the mortality from cardiovascular disease was statistically significant (Alexa, 1996).

We found important variation of the Mg and Ca concentration depending of the season.

Corrosivity of the water in public system is important. Magnesium appears as a competitive inhibitor of the two main noxious polluting agents: Pb and Cd.

### **Mg and Ca/Mg ratio in drinking water**

An optimal water would also have a ratio of Ca/Mg of not more than 2:1. (Durlach, 1985). Results are given in Table 2.

**Table 2.** The mean Ca/Mg ratio in the samples of the drinking water of the 5 cities in the public system and wells

| Town        | Public System                                              |         | Wells |         |
|-------------|------------------------------------------------------------|---------|-------|---------|
|             | july                                                       | octobre | july  | octobre |
| Iasi        | 4,63 in surface water Prut<br>4.5 in groundwater Timisesti |         | -     | -       |
| Pascani     | 8,55                                                       | 11,96   | 3,46  | 8,49    |
| Hirlau      | 5,13                                                       | 2,91    | 1,37  | 2,04    |
| Tg. Frumos  | 11,79                                                      | 16,84   | 0,7   | 0,75    |
| Podu Iloaie | 6,3                                                        | 8,4     | 1,3   | 1,8     |

### **Total hardness in drinking water**

Cardiovascular diseases rates may be inversely related to water hardness. Rates of cardiovascular mortality and sudden death are 10% to 30% greater in soft water areas (low in magnesium or calcium) than in hard water areas (high in magnesium or calcium).

Total hardness is made up by Ca and Mg compounds/ salts of other metals (Al, Ba, Fe, Mn, Sr).

The mean concentration of the total hardness in drinking water of the public system, was ranged between 12.02 – 16. 28<sup>o</sup>G (table 3). We found important variation of the total hardness (Mg and Ca too) concentration depending of the season (table 3 and 4).

**Table 3.** The mean, maximum and minimum of the total hardness (grd. G) in the samples of the drinking water of the 4 cities on the public system

| Town       | july  |         |         | octobre |         |         |
|------------|-------|---------|---------|---------|---------|---------|
|            | mean  | maximum | minimum | mean    | maximum | minimum |
| Iasi*      | 12,02 | 17,78   | 9,32    | -       | -       | -       |
| Iasi**     | 13,10 | 14,11   | 11,42   | -       | -       | -       |
| Pascani    | 13,33 | 15,03   | 12,12   | 16,28   | 25      | 13,5    |
| Hirlau     | 15,66 | 20,2    | 13,91   | 15,15   | 19,5    | 12,6    |
| Tg. Frumos | 14,09 | 15,93   | 9,14    | 13,2    | 14,6    | 11,9    |

\* surface water Prut; \*\*Groundwater Timisesti



The total hardness was higher in the wells water with variations of the mean concentration between 20, 36 – 38, 37 grd.G(table 4).

**Table 4.** The mean, maximum and minimum of the total hardness (grd. G) in the samples of the drinking water in the wells of the 3 cities

| Town       | july  |         |         | octobre |         |         |
|------------|-------|---------|---------|---------|---------|---------|
|            | mean  | maximum | minimum | mean    | maximum | minimum |
| Pascani    | 23,78 | 33,43   | 13,91   | 20,36   | 33,9    | 13,5    |
| Hirlau     | 38,37 | 55,65   | 22,89   | 31,2    | 55,4    | 20,6    |
| Tg. Frumos | 32,91 | 42,64   | 7,28    | 34,21   | 44,9    | 17      |

The chemical analyse of the samples point out a low degree of the mineralization and fixed residuum in average below 500 mg/l.

## CONCLUSIONS

In our study, in Iasi county, in urban area (5 cities with an total of 388. 650 inhabitants) the Mg content of drinking water may provide below 10% of the RDI in public system and between 10,78 % – 48,5 % of the RDI in wells, with important variations.

Mg need to be taken into account as a protective factor in cardiovascular diseases when represent quantitatively a significant part of the Recommended Dietary Allowances or if, quantitatively constitute a higher bioavailable intake.

In the drinking water of the public system of the 5 cities magnesium average concentration was higher in Hirlau (mean 23,74 mg/l on july and 28,75 mg/l on octobere) and in order in Podu Iloaie city (mean 27,18 mg/l on july and 14,33 mg/l on octobere), Iasi city (mean 19,25 mg/l in surface water Prut and 9,46 mg/l on groundwater Timisesti), in Pascani city (mean 12,23 mg/l on july and 13,52 mg/l on octobere), in Tg. Frumos (mean 10,28 mg/l on july and 5,36 mg/l on octobere).

In the drinking water of the wells, of the 5 cities, magnesium average concentration was higher on well water of Tg. Frumos, Podu Iloaie and Hirlau city and lower in depth water of the city Pascani

Because Mg concentration in the drinking water in the public system of the towns in Iasi county is below 30 mg/l, Mg is not competitive inhibitor of the two main noxious water polluting agents: Pb and Cd.

The mean Ca/Mg ratio in the water of the public system in urban area of Iasi county varied between 2,91- 16,84 and in wells water between 0.75- 8.49 (optim 2/1).

The chemical analyse of the samples point out a low degree of the mineralization and fixed residuum in average below 500 mg/l.

The mean concentration of the total hardness in drinking water of the public system, was ranged between 12,02 – 16, 28<sup>o</sup>G. The total hardness was higher in the wells water with variations of the mean concentration between 20,36 – 38,37 grd. G.

We found important variation of the total hardness (Mg and Ca too) concentration depending of the season.

It is also the possibility to supplement the drinking water of the public system with magnesium.

Tap water was more effective in meeting magnesium requirements than dietary supplementation.

## REFERENCES

1. Alexa L., Gavăt V, Neamțu G, Voroniuc O, Albu A, Alexa I: Mineralization of the drinking water in two districts from Moldavia. Metal elements in environment Biology and Medicine, Publishing House "Eurobit" Timișoara, pp 109-116, 1996.
2. Cuciureanu R.: Apa – element de mediu, 1-66, in Elemente de igiena mediului și a alimentației, Ed Junimea Iași, 2002,
3. Durlach, M.J., Bara A. Guiet-Bara: Magnesium Level in Drinking Water and Cardiovascular Risk Factor: A Hypothesis, Magnesium 1985, 4: 5-15.
4. Gavăt V.: Igiena mediului, Ed. „ Gr. T.Popa” U.M.F. Iași, 2007.
5. Ionuț C: Compendiu de igienă, Ed. Medicală Universitară „Iuliu Hațieganu” Cluj – Napoca, 2004
6. Nerbrand C, Agrens L, Lenner RA.: The influence of calcium and magnesium in drinking water and diet on cardiovascular risk factors in individuals living in hard and soft water areas with differences in cardiovascular mortality, BMC Public Health. 2003; 18:10-21.
7. Rylander C. R., Arnaud J.M.: Mineral water intake reduces blood pressure among subjects with low urinary magnesium and calcium levels, BMC Public Health 2004, 4:56 4-56
8. Sabatier M., Arnaud J., & co: Meal effect on magnesium bioavailability from mineral water in healthy women American Journal of Clinical Nutrition, 2002, Vol. 75, No. 1, 65-71.
9. Seeling M.S., Rosanoff A.: The Magnesium Factor, Ed. Avery Penguin Group USA, Inc., New York, 2003, 1-375.
10. Voroniuc O, Tarcea M., Buksa C., Rusu I, Gavăt V., Voroniuc I: Variation in Mineral Content of Bottled Romanian Waters Abstract book European Magnesium Meeting, Paris, 2008, P10.

# THE REDOX REACTIVITY OF GLOBINS: THE CHICKEN AND EGG PARADOX

Radu SILAGHI-DUMITRESCU\*, Florina DEAC

*Department of Chemistry, Faculty of Chemistry and Chemical Engineering Babeş-Bolyai University,  
Str. Arany Janos Nr. 11, RO-400 028 Cluj-Napoca, Roumania*

\* E-mail : rsilaghi@chem.ubbcluj.ro

## ABSTRACT

*The role of globins in oxygen transport and storage is well established and has long been known. However, more recently discovered members of the globin family appear to clearly have functions in vivo that do not involve oxygen storage and transport. Such are the globins found in inferior organisms, for which nitric oxide dioxygenase as well as peroxidase functions have been proposed; the roles of neuroglobins, cytoglobins and leghemoglobins are yet to be fully established, with nitrite reduction among the possible candidates. With globins present in organisms that seem to precede aerobic metabolism, it appears most likely that all of these redox activities, which may appear at least exotic in relation to hemoglobin and myoglobin, do in fact precede, in an evolutionary sense, the dioxygen transport function in the globin family.*

**Key words:** globins, oxidative stress, nitrosative stress

Globins are mostly known for their functions in dioxygen transport (hemoglobin) and storage (myoglobin). However, more recently a few other types of globin reactivities, mostly of redox and enzymatic nature, have been found to have biological/physiological/medical relevance. These reactivities are the focus of the current presentation; whereas in relation to the dioxygen handling function they appear as rather exotic and perhaps even fortuitous, the redox reactivities are now well accepted as physiological functions for a range of globins.

The globin family has in recent years expanded to include a number of new members. Among these, **neuroglobin** has been demonstrated to be expressed in human brain.<sup>[1]</sup> Unlike hemoglobin and myoglobin, neuroglobin in its deoxy (ferrous, not bound to oxygen) state appears hexacoordinated endogenously, which suggests its function is distinctly not one involving ligand binding – hence, not dioxygen storage or transport; the two axial ligands are, as expected, the homologues of the famous hemoglobin proximal and distal histidines.<sup>[1]</sup> Moreover, even the ferric form is still hexacoordinated, as a further argument against ligand binding playing any role in neuroglobin's physiological function. On the other hand, both dioxygen and CO can, in vitro, displace the 'distal' histidine; however, the oxygen affinity is too low for this reactivity to be relevant under the low dioxygen concentrations found in neuronal cells, and the autooxidation rate is also distinctly higher than that of myoglobin.<sup>[1-7]</sup> It is thus generally accepted that neuroglobin's in vivo function is not dioxygen binding; neuroglobin is overexpressed under hypoxia and ischemia, and exerts a protective

---


„Metal Elements in Environment, Medicine and Biology”, Tome VIII, pp. 271-276, Publishing House “Eurobit” Timișoara, 2008

Proc. of 8<sup>th</sup> Int. Symp. of Roumanian Academy - Branch Cluj-Napoca and Branch Timișoara, 2008, December 5-6, Timișoara, Roumania

effect under these conditions.<sup>[4-7]</sup> Ferric, but not ferrous, human Ngb has been shown to bind to heterotrimeric G proteins, acting as a guanine nucleotide dissociation inhibitor, eventually protecting against neuronal cell death; within this framework, Ngb would be acting as a redox sensor.<sup>[1]</sup> A physiological role for the ferric (met) form of a globin is certainly at considerable distance from everything known about the 'traditional' members of this family – hemoglobin and myoglobin.<sup>[1]</sup> On the other hand, a role in reactive nitrogen species scavenging (e.g., peroxynitrite) has also been proposed for neuroglobin.<sup>[8]</sup>

Closely related to neuroglobin is **cytoglobin**. It, like neuroglobin, appears hexacoordinated in the deoxy state. However, its dioxygen affinity falls within the physiological range, and subject both to cooperativity (cytoglobin is a dimeric protein) and to control via a disulfide bridge. Interestingly, however, is that cytoglobin appears localized exclusively in cell nuclei, which suggests that its function, while related to dioxygen metabolism, is not simply one of storage, and may in fact be one of transcriptional regulator, still under some redox (via the cysteine disulfide bridge) control.<sup>[4,5]</sup>

A family of slightly smaller ('truncated') globins, occasionally condensed with a flavin-containing redox partner, has been identified in inferior organisms, such as *E. coli*. Of these, **flavoheemoglobin** (Fhb) has probably received most of the attention. One of the roles proposed for Fhb has been the reaction with lipid hydroperoxides; indeed, this protein is reactive towards such peroxides more so than towards H<sub>2</sub>O<sub>2</sub>; moreover, a possible role for the ferrous form of the protein has been invoked in this reaction – which would be at odds with all other reactions of heme proteins with peroxides, known in general to involve ferric, not ferrous, hemes.<sup>[9,10]</sup> In any case, dioxygen storage is distinctly not a physiological function, in view of the high autooxidation rate.<sup>[11-13]</sup> On the other hand, Gardner et al have demonstrated that Fhb protects *E. coli* against toxic levels of nitric oxide, under non-anaerobic conditions, and that, moreover, expression of this protein is induced under these conditions.<sup>[11-13]</sup> The proposed mechanism of this process involves an iron(II)-O<sub>2</sub> + NO reaction<sup>[14-25]</sup>, with initial attack of NO on the (formally) iron(III)-superoxo heme, forming an iron(III)-peroxynitrite species, [Fe-OONO]<sup>2+</sup> (Reaction 1). Formation of this intermediate is controlled only by diffusion of NO into the heme pocket, in line with the superoxo character of the dioxygen ligand; free superoxide indeed reacts with NO at rates approaching the diffusion limit.<sup>[26]</sup> The peroxynitrite adduct then decays to resting iron(III) heme, liberating nitrate.



### Reaction 1

An alternative mechanism for flavoheemoglobins, with O<sub>2</sub> attacking an Fe(II)-NO heme adduct to produce an N-bound iron(III)-peroxynitrite intermediate,<sup>[27]</sup> is disfavoured by DFT calculations.<sup>[10]</sup> Moreover, in mammalian myoglobins, the reaction of Fe(II)-NO with dioxygen is rate-limited by dissociation of nitric oxide and a subsequent mechanism cf. Reaction 1.<sup>[28]</sup>

In relation to the peroxidase function proposed for flavoheemoglobin, we and others have demonstrated that **myoglobin** as well as **hemoglobin** exhibit an ascorbate peroxidase reactivity, which is entirely enzymatic in nature as characterized within the Michaelis-Menten formalisms.<sup>[29,30]</sup> In fact, the K<sub>m</sub> values for ascorbate in this process are in the low-micromolar range for hemoglobin, and well within the range of physiological concentrations of ascorbate, suggesting that

hemoglobin is especially designed to interact with ascorbate, and in fact to our knowledge it is the one protein with the highest affinity for ascorbate, of all those known to us so far, from any species.<sup>[29]</sup> Figure 1 illustrates the type of reactivity entailed by globin reaction with peroxides.




**Fig. 1.** Shown in black and with solid arrows is the peroxidase general mechanism. The electrons required for reduction of Compound I and Compound II are supplied by a reducing substrate (typically a small organic molecule, ascorbate in the globin case). Shown in gray and with a solid arrow is the peroxxygenase variation of the catalytic cycle, which is less common in globins; here, 'R' is an organic substrate and 'RO' is its monooxygenated form. Shown in gray and with dashed arrows is the oxygenase variation, which again is even further less common in globins. 'X' denotes the protein-derived axial ligand which, in globins, is the proximal histidine.

A final type of reactivity taken into discussion here is that of nitrite reductase. This reaction, occurring via mechanisms illustrated in Fig.2, liberates nitric oxide, known for its physiological role in blood among other things.<sup>[31,32]</sup>




**Fig.2** Mechanism of nitrite reductase reaction

Of particular interest to us is the occurrence of nitrite linkage isomerism in this reaction, which was introduced by us based on density functional theory (DFT) calculations, and has since been demonstrated experimentally by crystal structures.<sup>[33]</sup> On the other hand, with this reaction contributing to the regulation of the NO concentration in blood, significant interest has recently been shown towards

the possibility of nitrite being used as a drug for cardiovascular issues.<sup>[34-41]</sup> The key importance, and medical relevance, of this and of the peroxidase reactivity of globins,<sup>[29,30]</sup> do question any attempt to view these redox reactivities as mere essentially fortuitous side-reactions in hemoglobin and myoglobin. Moreover, as the redox reactivity appears as a predominant function in globins found in inferior organisms, in metabolic systems that are likely to have preceded aerobic life, it appears that the functions of dioxygen transport and storage in globins must have evolved subsequent to those of redox nature, and that, in chronological order, reversible dioxygen handling is in fact a side-reaction of the initial, redox based, physiological function(s) of globins in ancient organisms.

## REFERENCES

1. Wakasugi, K., Kitatsuji, C. and Morishima, I., Possible neuroprotective mechanism of human neuroglobin, *Ann N Y Acad Sci*, 2005, 1053, 220-30.
2. Dewilde, S., Kiger, L., Burmester, T., Hankeln, T., Baudin-Creux, V., Aerts, T., Marden, M.C., Caubergs, R. and Moens, L., Biochemical characterization and ligand binding properties of neuroglobin, a novel member of the globin family, *J Biol Chem*, 2001, 276, 38949-55.
3. Dewilde, S., Mees, K., Kiger, L., Lechaue, C., Marden, M.C., Pesce, A., Bolognesi, M. and Moens, L., Expression, purification, and crystallization of neuro- and cytoglobin, *Methods Enzymol*, 2008, 436, 341-57.
4. Fago, A., Hundahl, C., Malte, H. and Weber, R.E., Functional properties of neuroglobin and cytoglobin. Insights into the ancestral physiological roles of globins, *IUBMB Life*, 2004, 56, 689-96.
5. Fago, A., Hundahl, C., Dewilde, S., Gilany, K., Moens, L. and Weber, R.E., Allosteric regulation and temperature dependence of oxygen binding in human neuroglobin and cytoglobin. Molecular mechanisms and physiological significance, *J Biol Chem*, 2004, 279, 44417-26.
6. Fago, A., Mathews, A.J., Moens, L., Dewilde, S. and Brittain, T., The reaction of neuroglobin with potential redox protein partners cytochrome b5 and cytochrome c, *FEBS Lett*, 2006, 580, 4884-8.
7. Fago, A., Mathews, A.J., Dewilde, S., Moens, L. and Brittain, T., The reactions of neuroglobin with CO: evidence for two forms of the ferrous protein, *J Inorg Biochem*, 2006, 100, 1339-43.
8. Herold, S., Fago, A., Weber, R.E., Dewilde, S. and Moens, L., Reactivity studies of the Fe(III) and Fe(II)NO forms of human neuroglobin reveal a potential role against oxidative stress, *J Biol Chem*, 2004, 279, 22841-7.
9. Silaghi-Dumitrescu, R., Heme ferrous-hydroperoxo complexes, *Arch. Biochem. Biophys.*, 2004, 424, 137-140.
10. Silaghi-Dumitrescu, R., A density functional study of heme-peroxynitrite adducts., *J. Mol. Struct. THEOCHEM*, 2005, 722, 233-237.
11. Gardner, P.R., Gardner, A.M., Martin, L.A., Dou, Y.D., Li, T., Olson, J.S., Zhu, H. and Riggs, A.F., Nitric-oxide dioxygenase activity and function of flavohemoglobins, *J. Biol. Chem.*, 2001, 275, 31581-31587.
12. Gardner, A.M. and Gardner, P.R., Flavohemoglobin detoxifies nitric oxide in aerobic, but not anaerobic, *Escherichia coli*. Evidence for a novel inducible anaerobic nitric oxide-scavenging activity, *J. Biol. Chem.*, 2002, 277, 8166-71.

13. Gardner, A.M., Helmick, R.A. and Gardner, P.R., Flavorubredoxin, an inducible catalyst for nitric oxide reduction and detoxification in *Escherichia coli*, J. Biol. Chem., 2002, 277, 8172-7.
14. Herold, S., Kinetic and spectroscopic characterization of an intermediate peroxynitrite complex in the nitrogen monoxide induced oxidation of oxyhemoglobin, FEBS Lett, 1999, 443, 81-4.
15. Herold, S., Kinetic and spectroscopic characterization of an intermediate peroxynitrite complex in the nitrogen monoxide induced oxidation of oxyhemoglobin, FEBS Lett, 1998, 439, 85-8.
16. Herold, S., Exner, M. and Nauser, T., Kinetic and mechanistic studies of the NO\*-mediated oxidation of oxymyoglobin and oxyhemoglobin, Biochemistry, 2001, 40, 3385-95.
17. Herold, S., Kalinga, S., Matsui, T. and Watanabe, Y., mechanistic studies of the isomerization of peroxynitrite to nitrate catalyzed by distal histidine metmyoglobin mutants, J. Am. Chem. Soc., 2004, 126, 6945-6955.
18. Herold, S., Matsui, T. and Watanabe, Y., Peroxynitrite isomerization catalyzed by His64 myoglobin mutants, J Am Chem Soc, 2001, 123, 4085-6.
19. Herold, S. and Shivashankar, K., Metmyoglobin and methemoglobin catalyze the isomerization of peroxynitrite to nitrate, Biochemistry, 2003, 42, 14036-46.
20. Herold, S., Shivashankar, K. and Mehl, M., Myoglobin scavenges peroxynitrite without being significantly nitrated, Biochemistry, 2002, 41, 13460-72.
21. Blomberg, L.M., Blomberg, M.R.A. and Siegbahn, P.E.M., A theoretical study of myoglobin working as a nitric oxide scavenger, J. Biol. Inorg. Chem., 2004, 9, 923-935.
22. Olson, J.S., Foley, E.W., Rogge, C., Tsai, A.L., Doyle, M.P. and Lemon, D.D., No scavenging and the hypertensive effect of hemoglobin-based blood substitutes, Free Radic Biol Med, 2004, 36, 685-97.
23. Eich, R.F., Li, T., Lemon, D.D., Doherty, D.H., Curry, S.R., Aitken, J.F., Mathews, A.J., Johnson, K.A., Smith, R.D., Phillips, G.N., Jr. and Olson, J.S., Mechanism of NO-induced oxidation of myoglobin and hemoglobin, Biochemistry, 1996, 35, 6976-83.
24. Moller, J.K. and Skibsted, L.H., Nitric oxide and myoglobins., Chem. Rev., 2002, 1167-78.
25. Moller, J.K. and Skibsted, L.H., Mechanism of nitrosylmyoglobin autoxidation: temperature and oxygen pressure effects on the two consecutive reactions, Chemistry, 2004, 10, 2291-300.
26. Beckman, J.S. and Koppenol, W.H., The good, the bad, and the ugly, Am. J. Physiol., 1996, 271, C1424-37.
27. Hausladen, A., Gow, A. and Stamler, J.S., Flavohemoglobin denitrosylase catalyzes the reaction of a nitroxyl equivalent with molecular oxygen, Proc Natl Acad Sci U S A, 2001, 98, 10108-12.
28. Herold, S. and Koppenol, W.H., Peroxynitritometal complexes, Coord. Chem. Rev., 2005, 249, 499-506.
29. Cooper, C.E., Silaghi-Dumitrescu, R., Rukengwa, M., Alayash, A.I. and Buehler, P.W., Peroxidase activity of hemoglobin towards ascorbate and urate: a synergistic protective strategy against toxicity of Hemoglobin-Based Oxygen Carriers (HBOC), Biochim Biophys Acta, 2008, 1784, 1415-20.
30. Silaghi-Dumitrescu, R., Reeder, B.J., Nicholls, P., Cooper, C.E. and Wilson, M.T., Ferryl haem protonation gates peroxidatic reactivity in globins, Biochem J, 2007, 403, 391-5.

31. Silaghi-Dumitrescu, R., The nitric oxide adducts of cytochrome cd1 nitrite reductase, *Rev. Chim.*, 2004, 55, 496-498.
32. Silaghi-Dumitrescu, R., Nitrite linkage isomerism in cytochrome cd1 nitrite reductase, *Inorg. Chem.*, 2004, 43, 3715-3718.
33. Copeland, D.M., Soares, A.S., West, A.H. and Richter-Addo, G.B., Crystal structures of the nitrite and nitric oxide complexes of horse heart myoglobin, *J Inorg Biochem*, 2006, 100, 1413-25.
34. Gladwin, M.T., Crawford, J.H. and Patel, R.P., The biochemistry of nitric oxide, nitrite, and hemoglobin: role in blood flow regulation, *Free Radic Biol Med*, 2004, 36, 707-17.
35. Gladwin, M.T., Haldane, hot dogs, halitosis, and hypoxic vasodilation: the emerging biology of the nitrite anion, *J Clin Invest*, 2004, 113, 19-21.
36. Gladwin, M.T., Hemoglobin as a nitrite reductase regulating red cell-dependent hypoxic vasodilation, *Am J Respir Cell Mol Biol*, 2005, 32, 363-6.
37. Gladwin, M.T., Nitrite as an intrinsic signaling molecule, *Nat Chem Biol*, 2005, 1, 245-6.
38. Gladwin, M.T., Role of the red blood cell in nitric oxide homeostasis and hypoxic vasodilation, *Adv Exp Med Biol*, 2006, 588, 189-205.
39. Gladwin, M.T., Raat, N.J., Shiva, S., Dezfulian, C., Hogg, N., Kim-Shapiro, D.B. and Patel, R.P., Nitrite as a vascular endocrine nitric oxide reservoir that contributes to hypoxic signaling, cytoprotection, and vasodilation, *Am J Physiol Heart Circ Physiol*, 2006, 291, H2026-35.
40. Gladwin, M.T. and Kim-Shapiro, D.B., The functional nitrite reductase activity of the heme-globins, *Blood*, 2008, .
41. Gladwin, M.T., Evidence mounts that nitrite contributes to hypoxic vasodilation in the human circulation, *Circulation*, 2008, 117, 594-7.



# THE SIGNIFICANCE OF CYTOMORPHOMETRY IN PERITONEAL EFFUSIONS CYTOLOGY PRACTICE

Ana-Maria CREȚU<sup>1\*</sup>, Mariana ASCHIE<sup>2</sup>, Gabriela PREDA<sup>1</sup>, Natalia ROȘOIU<sup>3</sup>

1\*. Faculty of Biology, University „Ovidius”, Bdl. Mamaia 124, RO-900 527 Constanta, Roumania;

2. Pathological Anatomy Department, Emergency Clinical Hospital, RO-900 591 Constanta, Roumania; 3. Faculty of Medicine, University „Ovidius”, RO-900 527 Constanta, Roumania

\* E-mail : cretu\_anamaria@yahoo.com

## ABSTRACT

The cytomorphological evaluation of effusion is a challenging aspect of non-gynecology cytology. The definite diagnosis of malignant effusion is often complicated by the presence of inflammation, exuberant mesothelial hyperplasia and a paucity of malignant cells. Computerised image analysis can rapidly digitise the image of the cells and thereby can estimate various morphometric parameters of the cell. Cytological samples suitable for image were obtained from cytocentrifuge. The previous study showed that existed significant differences between morphometric variables derived from nuclear and cytoplasmic outlines in benign and malignant effusions. In this retrospective study based on 58 available cases with a histological diagnosis, we have examined the peritoneal effusions from routine morphologic features. Cytologically, 34 (58,62%) peritoneal fluids smears were classified as benign (negative cytology), 18 (31.03%) were malignant (positive cytology) and 8 (13.79%) were considered suspicious. The various cellular parameters that were measured by cytomorphometry included nuclear as well as cytoplasmic diameter, area, ratio of nucleus to cytoplasmic area. Cytomorphometry helps in better and quicker sorting for malignant cells. Finally, 34 cases were found to be benign, 22 malignant and 4 in the category of atypical cases.

**Key Words:** cytomorphometry, peritoneal effusions, benign, malignant

## INTRODUCTION

Because of its complexity, cytopathologic evaluation of effusions fluids can be challenging. However, because effusions fluids are relatively easy to collect, any pathology laboratory may be confronted with such specimens, regardless of its level of expertise. In comparison to other common specimens, effusions fluids need special and unique approaches for cytopathologic evaluation.

The cytological diagnoses of serous effusions are usually made by routine cytomorphology with certainty, allowing treatment decisions. Various studies have shown a sensitivity of 57.3% and specificity of 89% by conventional cytology for the detection of malignant cells in effusion samples (Thunnissen et al., 1993). Studies have shown that positive and negative predictive values for detection of malignancy by cytomorphology are 89.3% and 69.4% respectively. However, a grey zone always exists, where the cytopathologist encounters problems in determining the nature of the cells whether reactive, atypical, or beyond doubt malignant (Bedrossian, 1994;

Motherby et al., 1999). Therefore various ancillary techniques should be used to increase the diagnostic accuracy of malignancy in serous effusions.

The conventional cytology rate for identification of neoplastic cells in effusions is about 60%. The rate of diagnostically equivocal effusions in routine cytology is dependent on the volume of effusion examined, type of preparation and staining, experience of the examiner, and application of ancillary methods. Peritoneal effusions are a frequently encountered clinical manifestation of metastatic disease, with breast, ovarian, and lung carcinomas and malignant mesothelioma leading the list (Redman et al., 1991; Creasman and Rutledge, 1971).

Neoplastic cells that disseminate into cavities containing effusions are highly metastatic and possess a strong autonomous proliferative drive while concurrently being stimulatory of exudative effusions.

The diagnosis of a malignant effusion signifies disease progression and is associated with a worse prognosis regardless of the tumor site of origin. Furthermore, cancer cells of different origins differ considerably in their biology and have unique phenotypic and genotypic characteristics (Johnson, 1966).

Primary cytomorphologic criteria of malignancy include cellular aggregates, pleomorphism (variable cellular appearance), anisocytosis (variation in cell size), anisokaryosis (variation in nuclear size), multinucleation, prominent to irregular nucleoli, increased nuclear to cytoplasmic ratio, monomorphic cellular appearance, and increased mitotic figures. Hyperplastic mesothelial cells also may exhibit anisocytosis, anisokaryosis, increased nuclear to cytoplasmic ratio, binucleate and multinucleate, and scattered mitoses. Any situation that results in fluid accumulation within the body cavities can induce mesothelial cell hyperplasia and exfoliation with an abnormal cellular morphology (Runyon, 1999). Therefore, the differentiation between mesothelial cell hyperplasia and mesothelioma may be difficult or impossible.

This study was carried out with the aim to analyse not only the mean nuclear variables from the sample of cells in each specimen, but also the outlying values, represented by the largest nucleus found in each, and to confirm the role of cytomorphometry on peritoneal effusions.

This study used computerized interactive morphometry to evaluate the differential characteristics of mesothelial normal cells, mesothelial hyperplastic cells, and carcinomatous cells.

## **MATERIALS AND METHODS**

This was hospital based prospective study carried out in Emergency Clinical Hospital of Constanta – Pathological Anatomy Department (SCJUC) from January 1, 2008 to 31 October, 2008. Clinical charts of all the patients whose peritoneal fluid samples were sent for cytological examination during the study period were retrieved for relevant information.

Histopathologic examination of tissue was used as gold standard to diagnose malignancy, in cases where this was available. Where histopathological examination was not done, diagnosis of malignancy was based on strong clinical suspicion, radiological methods and direct visualization of tumour.

The fluid for cytological analysis was collected during laparotomy from the abdominal cavity. If no fluid was present, the peritoneal cavity was lavaged with saline solution, and the fluid was then collected for analysis.

Follow-up data were obtained from the Tumor Registry at SCJUC.

For small fluid accumulations the entire specimen was submitted for laboratory evaluation. For larger effusions, 50-200 mL of well-mixed fluid was sent for cytologic examination; however, the entire specimen is also acceptable.

Giemsa stained and Papanicolaou stained slides were prepared from sediment obtained by centrifuging the peritoneal liquid samples at 1500 rpm for 5 minutes, using Shandon Cytospin preparations. After the centrifugation, the stained is fixed using alcohol (95% ethyl alcohol) as the fixative.

The Papanicolaou stain uses a standard nuclear stain, hematoxylin, and two cytoplasmic

Counterstains, OG-6 and EA. The outcome of this method is crisp nuclear detail and transparency of the cytoplasm, which allows the examiner to clearly visualize cellular morphology. The analysis of false positive and false negative peritoneal fluid sample shows the importance of using both the Giemsa stain and the Pap stain.

Effusion cytology was studied from 18 malignancy associated and 34 non malignancy associated ascitic fluid samples.

The smears collected were submitted to the image analysis. The morphometric differences between benign and malignant serous effusions, as diagnosed by standard cytologic criteria in 58 cases (34 benign and 18 malignant), were studied using the LUCIA Net semi-automated image analysis system, which calculates various parameters from tracings of cellular and nuclear outlines.

The procedure included extraction of multiple descriptors of the nuclear profile: perimeter length, area and distribution table of the nuclear area. Morphometric measurements of cellular and nuclear area, nuclear perimeter, nuclear/cytoplasmic ratio (N/C) and mean nuclear diameter of mesothelial cells were performed on single cells in peritoneal effusions from 18 patients with effusions caused by different types of carcinoma. The results were compared with corresponding measurements on mesothelial cells in peritoneal effusions from 34 patients without malignant disease.

Measurements were performed on the 50 cells in each specimen, excluding multinucleation and polymorphonuclear leucocytes.

## RESULTS AND DISCUSSIONS

Effusions associated with malignancy contained tumor cells and occasionally small fragments of neoplastic tissue in which may be recognized the relations of the tumor cells to their stroma. Fluids containing tissue fragments or groups of tumor cells offer little difficulty and ordinarily permit a positive diagnosis of malignancy. The cells of these clumps usually are larger than the other non-neoplastic cells of the sediment and because of their deeper staining give the slide a mottled appearance even under the low powers of the microscope.

The differentiation of neoplastic cells from mesothelial cells in such sediments is based on marked variation in size and shape, or the presence of large numbers of unusual-appearing cells, as well as on cytological factors such as multinucleation, giant nuclei and the presence of mitotic figures. Vacuolization of the cytoplasm is more common in tumor cells, but is also seen in the mesothelial cells of long standing effusions.

Some different cytologic findings, including smaller cell and nuclear size, larger relative nuclear area, fewer cells with large discrete cytoplasmic vacuoles and

macronucleoli, more frequent multinucleation and larger closely packed clusters of cells, were shown.

#### Cytologic features.

**Benign mesothelial cells** - when mesothelial cells are aspirated by a needle from the surface of the abdominal cavity, they appear as sheets of polygonal cells, about 15-30  $\mu\text{m}$  in diameter (1.5-2 times the size of neutrophils, but they may vary significantly and may range up to 50  $\mu\text{m}$  in diameter), that are usually separated from each other by clear „windows“. The cells have a delicate, yet sharply demarcated, cyanophilic or eosinophilic cytoplasm and round or oval nuclei. The nuclei, generally located in the center of the cell, are of even size (about 10  $\mu\text{m}$  in diameter), sharply demarcated, slightly granular and contain one or two, centrally located, readily visible nucleoli (Fig.1).

**Reactive mesothelial cells** - benign mesothelial cells may proliferate and display a spectrum of reactive changes. These range from minimal simple reactive changes to highly atypical reactive changes that enter into the differential diagnosis for malignancy. The mesothelial cells may maintain the morphologic features of normal mesothelial cells (round to oval nuclei with smooth nuclear contours and fine chromatin) (Fig.2).




**Fig.1.** Benign mesothelial cells. The nucleus is usually central or near central (b), but may be eccentric(c). Papanicolaou stained, 100X zoomed.




**Fig.2.** Reactive mesothelial cells (RM) with inflammatory cells (E- eosinophil, L- lymphocyte, N- neutrophil), 100x zoomed.

**Atypical mesothelial cells** - cytologic changes may include enlarged nuclei, prominent nucleoli, frequent mitosis. Nuclear contours may show some variation, but are still generally round to oval with smooth nuclear membranes (fig 3.).

Atypical mesothelial hyperplasia can mimic many of these cytologic and architectural features.

Cancer cells in malignant effusions are most often seen in separate individual cancer cells, in sheets and in spherical three-dimensional highly cellular




**Fig.3.** Multinucleation is common in benign conditions. The nuclei in benign multinucleated cells are similar in size, shape, and chromatin pattern in contrast to those of malignant giant cells which show marked variation in these features, ob.40X

clusters called morulae. Classic cytologic features of individual cancer cells in malignant peritoneal effusion include enlarged cells with high N:C ratio, coarse chromatin, enlarged and multiple nucleoli and irregular or indented nuclear contours (Fig. 4, 5 ).




**Fig.4.** Mesothelial cells (a, b, c) versus adenocarcinoma cells (d, e, f) with eccentric nuclei.




**Fig.5.** Cancer mesothelial cells are generally larger than normal counterparts with larger nuclei, marked hyperchromasia in well preserved cells suggest malignancy; ob.40X

On cytomorphometry, values for mean nuclear area and mean cytoplasmic area in benign cases were found to be  $11.16\mu\text{m}^2$  and  $60.70\mu\text{m}^2$ , while for malignant cases were  $20.31\mu\text{m}^2$  and  $68.66\mu\text{m}^2$ , respectively (Table 1, Table 2).

**Table 1.** Results of mean nuclear variables

| Nuclear data      |                               |                                |                              |                              |            |                             |
|-------------------|-------------------------------|--------------------------------|------------------------------|------------------------------|------------|-----------------------------|
|                   | Mean Area [ $\mu\text{m}^2$ ] | Eq. Diameter [ $\mu\text{m}$ ] | Max. Feret [ $\mu\text{m}$ ] | Min. Feret [ $\mu\text{m}$ ] | Elongation | Perimeter [ $\mu\text{m}$ ] |
| <b>B</b>          | 1116                          | 37.70                          | 49.00                        | 33.00                        | 1.48       | 126.95                      |
| <b>R</b>          | 1585                          | 44.92                          | 58.00                        | 40.00                        | 1.45       | 151.17                      |
| <b>M</b>          | 2031                          | 50.85                          | 59.00                        | 50.00                        | 1.18       | 166.08                      |
| Global statistics |                               | Mean                           | StDev                        | Min                          | Max        |                             |
| Area              |                               | 1577                           | 457.55                       | 1116                         | 2031       |                             |
| Eq. Diameter      |                               | 44.49                          | 6.59                         | 37.70                        | 50.85      |                             |
| Max. Feret        |                               | 55.33                          | 5.51                         | 49.00                        | 59.00      |                             |
| Min. Feret        |                               | 41.00                          | 8.54                         | 33.00                        | 50.00      |                             |
| Elongation        |                               | 1.37                           | 0.17                         | 1.18                         | 1.48       |                             |
| Perimeter         |                               | 148.07                         | 19.75                        | 126.95                       | 166.08     |                             |

The evaluation of nuclear/nucleolar size was accomplished by the use of a micrometer at X40 magnification using as reference nearby red blood cells, which measured approximately 7  $\mu\text{m}$ .

**Table 2.** Results of mean cytoplasmic variables

| Cytoplasmic data  |                          |                                |                              |                              |            |                             |
|-------------------|--------------------------|--------------------------------|------------------------------|------------------------------|------------|-----------------------------|
|                   | Area [ $\mu\text{m}^2$ ] | Eq. Diameter [ $\mu\text{m}$ ] | Max. Feret [ $\mu\text{m}$ ] | Min. Feret [ $\mu\text{m}$ ] | Elongation | Perimeter [ $\mu\text{m}$ ] |
| <b>B</b>          | 6070                     | 87.91                          | 113.00                       | 83.00                        | 1.36       | 323.05                      |
| <b>R</b>          | 6300                     | 89.56                          | 116.00                       | 74.00                        | 1.57       | 307.81                      |
| <b>M</b>          | 6866                     | 93.50                          | 121.00                       | 108.00                       | 87.00      | 310.37                      |
| Global statistics |                          | Mean                           | StDev                        | Min                          | Max        |                             |
| Area              |                          | 6412                           | 409.65                       | 6070                         | 6866       |                             |
| Eq. Diameter      |                          | 90.32                          | 2.87                         | 87.91                        | 93.50      |                             |
| Max. Feret        |                          | 112.33                         | 4.04                         | 108.00                       | 116.00     |                             |
| Min. Feret        |                          | 81.33                          | 6.66                         | 74.00                        | 87.00      |                             |
| Elongation        |                          | 1.39                           | 0.16                         | 1.24                         | 1.57       |                             |
| Perimeter         |                          | 313.74                         | 8.16                         | 307.81                       | 323.05     |                             |

Significant differences were found between the mean values from cellular area, nuclear area, and nuclear: cytoplasmatic ratios (N:C) in mesothelial cells from benign effusion versus malignant cells from effusions associated with tumour (Table 3).

**Table 3.** Results of N/C ratios and standard deviation

|                                       | Benign   | Reactiv  | Malign   |
|---------------------------------------|----------|----------|----------|
| <b>N/C [<math>\mu\text{m}</math>]</b> | 1/5      | 1/4      | 1/3      |
| <b>Std.dev.</b>                       | 499,0904 | 708,8335 | 908,2908 |

The nuclear size was determined by the assessment and measurement of the predominant nuclear size found throughout the multiple specimens examined in each case. The largest dimension of the nucleus repetitively seen was scored as the nuclear size.

Cytomorphometry helps in better and quicker sorting for malignant cells. Finally, 34 cases were found to be benign, 22 malignant and 4 in the category of atypical cases.

The results indicated that the range of the values for suspicious cells is closer to the range of the benign cells (Fig.5).




**Fig.5.** Mean nuclear and cytoplasmic area of mesothelial cells

## CONCLUSIONS

1. The previous study showed that existed significant differences between morphometric variables derived from nuclear and cytoplasmic outlines in benign and malignant effusions. In conclusion, these results confirm data from literature and indicate that cytometric analysis of nuclear parameters represent a useful marker for identification of malignant cells in equivocal effusions and can be used to increase the cytological sensitivity in doubtful mesothelial proliferations.



2. Morphometric measurements of peritoneal fluids can be a valuable adjunct to routine light microscopy of cytology specimens and may provide important information for the differentiation of atypical atypical mesothelial cells from malignant adenocarcinoma cells. Increasing the sample size may improve test performance. Morphometry is of value in several areas of diagnostic histopathology. Its application to cytology would be of considerable benefit both in terms of diagnostic accuracy and as the basis of an automated screening programme.

3. Ascitic fluid cytology is a simple and useful procedure and should be routinely requested. In a set up like ours where definite involvement of peritoneal liquid cytology still can detect malignant cells. As presence or absence of malignant cells is useful in predicting the prognosis of patients and sometimes it can be the only clue to presence of malignancy, ascitic fluid cytology should be routinely requested.

## REFERENCES

1. Bedrossian C.W.M. - Malignant effusions: A multimodal approach to cytologic diagnosis, Igaku-Shoin, New York, 1994.
2. Creasman W.T., Rutledge F. - The prognostic value of peritoneal cytology in gynecologic malignant disease, *Am. J. Obstet. Gynecol.*, 1971, 110, 773-781.
3. Johnson W.D. - The cytological diagnosis of cancer in serous effusion, *Acta cytological*, 1966, 10, 161-172.
4. Motherby H., Nadjari B., Friegel P., Kohaus J., Ramp U., Böcking A. - Diagnostic accuracy of effusion cytology, *Diagn. Cytopathol*, 1999, 20, 350-351.
5. Redman C.W., Chapman S.E., Chan S.Y., Buxton E.J., Blackledge G., Luesley D.M. - Out-patient peritoneal lavage cytology in the detection of residual epithelial ovarian cancer, *Cytopathology*, 1991, 2, 291-298.
6. Runyon B. - Approach to the patient with ascites, pp. 966-991, In: *Textbook of Gastroenterology* (eds: Yamada T., Alpers D.H., Laine L., Owyang C., Powell D.W), 3rd ed., Philadelphia: lippincott Williams & Wilkins, 1999.
7. Thunnissen F.B., Peterse J.L., Van Pel R., Jansen J.W., Laddé B.E., Arends J.W., Volovics A. - Reliability of fine needle aspiration cytology for distinguishing between carcinoma, lymphoma and sarcoma: the influence of clinical information, *Cytopathology*, 1993, 4(2), 107-114.



# AMBROSIA MOWING IN NOVI SAD AND THE NUMBER OF CHILDREN WITH ALLERGIC DISEASES REGISTERED IN PRIMARY CARE

Mila HADNADJEV <sup>1\*</sup>, Darka HADNADJEV <sup>2</sup>

1. Health Center Novi Sad, Department of Children's Health Protection, Bulev. Cara Lazara 75, 21 000 Novi Sad, Serbia; 2. Health Center Indjija, Emergency Department, Srpskokrkvena 5, Serbia

\* E-mail : darka@eunet.yu

## ABSTRACT

*Purpose was to find out the difference in the number of outpatient registered allergic children, after Ambrosia mowing in 2007, comparing to the year 2006, considering the third(III), fourth(IV) quarter and the whole year. Method. Ambrosia Eradication Headquarters conducted Ambrosia mowing in 2006 and in 2007. We collected the data on all allergic diseases in 2006 and 2007 from the Health Center official records, before and after Ambrosia mowing and compared them. Results. The number of the outpatient registered allergic small children (aged 0-6years) showed statistically significant differences in the III quarter of the year 2006 to the same period of 2007 ( $p=0.000$ ) and the number of the outpatient registered school children with all allergic diseases showed statistically significant differences ( $p=0.000$ ) in III and in IV quarters of each year and in the whole year of 2007 ( $p=0.000$ ). Conclusion: There was significantly smaller number of outpatient registered small children with allergic diseases only in III quartal of the year 2007, and significantly lower number of school children with allergic diseases in III, IV quartal and the whole year of 2007. Total number of outpatient registered children was significantly smaller in the year 2007.*

**Key words:** Ambrosia, mowing, allergic diseases, children

## INTRODUCTION

Ambrosia artemisifolia L. is one year, foreign, invasive ragweed which was brought from North America, at the end of the XIX century. It spreads from the west to the east part of Europe and it came to Serbia from Hungary, mainly along the main roads.

Ambrosia pollen is harmful for human health and the main thing is the number of pollen grains in the air per cubic meter. It is known that 30 grains harm respiratory system in sensitized persons causing, itchy, running, sneeze, stuffy nose i.e. allergic rhinitis and wheezing, tight chest, lack of air, i.e. bronchial asthma. But, the whole plant can also provoke eczema or atopic dermatitis (1). As avoidance of allergen is the first step of therapy to relieve symptoms and Ambrosia eradication before its flowering period, is of the main significance.

Purpose was to find out the difference in the number of outpatient registered allergic children, after Ambrosia mowing in 2007, comparing to the year 2006, considering the third, fourth quarter and the all twelve months of the each year.

## **MATERIALS AND METHODS**

Ambrosia Eradication Center Headquarters was conducting Ambrosia mowing from the end of June till the beginning of November in 2006 and in 2007. In the year of 2006, mowing of ragweed was conducted on 500 ha and in the year of 2007. on the 600 ha (urban area). Mowing was conducted mechanically and chemically (herbicide glyphosate in the rate of 1200-2400 g a.i/ha was applied). During 2007 only, on the territory of the City of Novi Sad ragweed mechanical control was performed at 367 030 m<sup>2</sup>, and at suburban areas herbicide treatments were applied on about 325 000 m<sup>2</sup> (2). We considered allergic diseases classified in International Statistical Classification on Diseases and Related Health Problems as: L10-L99, where are ciphers for allergic skin diseases implicated, as Dermatitis atopica (L20), Dermatitis contacta allergica (L23) and Urticaria (L50), and for allergic respiratory diseases as Morbus pulmonis obstructivus chronicus alius (J44), Asthma bronchiale (J45) and Rhinitis vasomotoria et rhinitis allergica (J30) (3). We collected the data on all allergic diseases from the Health Center official records, before and after Ambrosia mowing and compared them to the data of the same period (III, IV quarters and the whole year) of the year before. The source of number of children was last census from 2002 (Population: Census, 2002) (4). Statistical method: t test.

## **RESULTS AND DISCUSSIONS**

Ragweed have flowering period from July to October each year. The number of pollen grains depends on several factors. The greatest number is in mid August and in early September (peak days), in the morning of dry, hot, mild windy day. High pollen loads in Vienna occur with winds coming from Hungary (south-eastern directions) (5,6). Although Ambrosia spp. mainly inhabits disturbed soil, such as agroecosystems, no significant difference has been observed between ragweed airborne pollen characteristics in rural and urban areas (7). During the period of 2000-2004 both the annual sums of daily averages and count of peak days changed with fluctuations. Diurnal ragweed pollen counts were over 30 pollen grains for 6-57 days of its 2-3 months long season, which means severe pollen release of the air (8). During the period of 2005-2007 ragweed pollen mean concentrations were increased considering the year 2005 (9). In the year 2007 a larger Ambrosia area eradication was conducted than it was in 2006 so were expecting to find a lower number of outhospital registered allergic children in 2007.

**Table 1** The number of small and school outhospital registered allergic children in the years 2006 and 2007 and quarter distribution

| Quater      | I    | II   | III  | IV   | I- XII mo | I- XII mo 0-19years |
|-------------|------|------|------|------|-----------|---------------------|
| 2006 small  | 2821 | 1339 | 2002 | 1806 | 7968      |                     |
| 2006 school | 1761 | 1431 | 2270 | 1736 | 7198      | 15166               |
| 2007 small  | 2812 | 2086 | 1663 | 1764 | 8325      |                     |
| 2007 school | 1750 | 2213 | 776  | 1119 | 5858      | 14183               |

Small=small children

School=school children

In the III quarter of 2006 (Table 1), there were more registered allergic small children (n=2002) and school children (n=2270) than in the same period of 2007 (n=1663, n=776, respectively). The same situation is in the IV quarter. Considering the number of registered allergic children for the whole year (I-XII) there were more small allergic children in 2007 than in 2006, but not school allergic children. In total (0-19 years), in 2007 there were less allergic children than in 2006.

**Table 2.** Comparison of the third and fourth quarters of the years 2006. and 2007.- outhospital registered small allergic children

| Quarters | m1(2006) | n1    | m2(2007) | n2    | t      | P     |
|----------|----------|-------|----------|-------|--------|-------|
| III      | 2002     | 20300 | 1663     | 20300 | 5.871  | 0.000 |
| IV       | 1806     | 20300 | 1764     | 20300 | 0.736  | 0.462 |
| I-XII mo | 7968     | 20300 | 8325     | 20300 | -3.615 | 0.000 |

M1= number of small allergic children (0-6years) in the III quarter of the year 2006

N1, N2= Number of small children (0-6years) in Novi Sad (Census 2002)

M2= number of allergic small children (0-6 years) in the IV quarter of the year 2007

Matching the number of the outpatient registered children with allergic disease (Table 2) (aged 0-6years) very high statistically significant differences were seen in the III quarter of the year 2006 to the same period of 2007 in the benefit of the year 2007 ( $p=0.000$ ), while comparing to the number in the IV quarters of each year there was no statistically significant difference ( $p=0.462$ ). Comparing the total number of the small allergic children in the whole year (I-XII) of 2006 to the whole year 2007, there were significantly smaller number in the year of 2006 ( $p=0.000$ ).

**Table 3** Comparison of the third and fourth quarters of the years 2006. and 2007.- school and total number of allergic children (0-19 years)

| Quarters  | M1    | N1    | M2    | N2    | t      | P     |
|-----------|-------|-------|-------|-------|--------|-------|
| III       | 2270  | 46373 | 776   | 46373 | 27.526 | 0.000 |
| IV        | 1736  | 46373 | 1119  | 46373 | 11.729 | 0.000 |
| I-XII mo  | 7198  | 46373 | 5858  | 46373 | 12.652 | 0.000 |
| Wh.yr0-19 | 15166 | 66673 | 14183 | 66673 | 6.497  | 0.000 |

M1= number of allergic school children (7-19years) in the III quarter of the year 2006

N1, N2= Number of school children (7-19 years) in Novi Sad (Census 2002)

M2= number of allergic school children (7-19 years) in the IV quarter of the year 2007

There were (Table 3) significant differences in the number of outhospitaly registered allergic school children in the third and in the fourth quarter of the year 2006 to the same period in the year 2007, in the benefit of the year 2007 ( $p=0.000$ ). In the whole year of 2007 there were statistically significantly less number of allergic school children (7-19 years) and of the total number of children (0-19 years) ( $p=0.000$ ) comparing to the same period of the year 2006.

## CONCLUSIONS

1. Matching the III and IV quarters of the years 2006 and 2007, during and after Ambrosia mowing, there was significantly lower number of outpatient registered small children with allergic diseases in the III quarter of 2007 and not significantly lower in the IV quarter of 2007.
2. The total number of outpatient registered small children in the all twelve months in the year 2007 was significantly higher than in 2006.
3. The number of outpatient registered school children with allergic diseases in the both quarters of the year 2007 was significantly lower than in 2006.
4. Total number of outpatient registered school children was significantly lower in all twelve months of the year 2007 comparing to the same period of the year 2006.
5. Total number of all outpatient registered allergic children (0-19years) was lower in all twelve months of 2007 comparing to all twelve months of 2006.

## REFERENCES

1. Jarić D, Radišić P, Šikoparija B: Sublingvalna imunoterapija u terapiji astme i alergijskog rinitisa. Zbornik sažetaka I kongresa alergologa i kliničkih imunologa Srbije, Novi Sad, 16-19. Novembar 2008.
2. Jovanović M, Mimica-Dukić N, Poljački M, Boža P: Erythema multiforme due to contact with weeds: a recurrence after patch testing. *Contact Dermatitis*, 2003, 48, 17-25. Blackwell Munksgaard.
3. Juhasz M, Juhasz IE, Gallovich E, Radišić P, Ianovici N, Peternel R, Kofol-Seliger A. Last year's ragweed pollen concentrations in the southern part of the Carpathian Basin. The 11th Symposium on Analytical and Environmental problems, Szeged, 24 September 2004, 339-343.
4. Konstantinović B, Meseldžija M, Konstantinović B.B.: Possibilities of Ambrosia *Artemisifolia* L. control in Northern Serbia. In: Book of Abstracts of The first international ragweed conference, 10-13 September 2008, Budapest, Hungary.
5. Šikoparija B, Radišić P, Pejak T, Simić S, Boža P: Ambrosia spp. Pollen in southern Panonnian valley. Abstracts of The 8th International Congress on Aerobiology "Towards a comprehensive vision", Neuchatel, Switzerland, 21-25 August 2006;48:2006
6. \*\*\*International Statistical Classification of Diseases and Related Health Problems: tenth revision Volume 1, World Health Organization, 1992, Geneva.
7. \*\*\*Population: Census, households and flats in 2002, 2, Gender and Age: settlements data. Belgrade Republic Statistic Bureau of Serbia, 2003 Belgrade: Republic Statistics Bureau of Serbia, 565, 2003.
8. \*\*\*[www.nspolen.com](http://www.nspolen.com)
9. \*\*\*[www.ragweed: Polleninfo.org](http://www.ragweed: Polleninfo.org)

## AUTHORS INDEX

### A

Acatincăi S. 217  
 Ahmadi M. 217  
 Albu G. 205  
 Alexandrova R. 105  
 Ardelean R. 127  
 Aschie M. 277  
 Avacovici A.E. 61, 217, 259

### B

Bandur G. 127  
 Baltă C. 163  
 Bankó S. 239  
 Băghină N. 113  
 de Benoist B. 87, 123  
 Bodescu F. 73  
 Bogdan A.T, 217  
 Borcean A. 113  
 Borcean I. 113  
 Brie I. 141  
 Buglyo P. 35  
 Bura M. 163

### C

Capo I. 81  
 Casaux-Voroniuc I. 265  
 Cernea V. 141  
 Ciopec M.E. 173  
 Cîmpeanu C. 205  
 Contrea A. 117  
 Costache M. 231  
 Costisor O. 105  
 Creanga A. 123  
 Crețu A.M. 277  
 Cristea A. 265  
 Csapo E. 35  
 Culic M. 81  
 Culita D.C. 105

### D

David Gh. 113  
 Davidescu C. M. 127  
 Deac F. 271  
 Dehelean C.A. 133  
 Delange F. 87  
 Dinischiotu A. 231, 249

Djujic I. 15, 95  
 Dorneanu S.A. 255  
 Dragan S. 133

### F

Farkas E. 35  
 Fischer-Fodor E. 5, 141  
 Florescu N. 205, 265

### G

Garban G. 147, 163, 259  
 Garban Z. 5, 61, 153, 163, 217  
 Gergen I. 133, 169, 195  
 Gheorgheș M. 265  
 Ghibu G.D. 163  
 Gogoasă I. 195  
 Grbic G. 81

### H

Hadnadjev D. 285  
 Hadnadjev M. 285  
 Haiduc I. 5  
 Halasi T. 81  
 Hădărugă D.I. 173  
 Hădărugă N.G. 173  
 Hărmănescu M. 169  
 Hegedűs A. 199, 225  
 Henin N. 123  
 Horhoi D.E. 163

### I

Ianos Gh. 181  
 Ilia Gh. 127  
 Iordache V. 41, 73

### K

Kalfin R. 105  
 Kantouri M.L. 105  
 Kastori R. 55  
 Kiss A.S. 211  
 Kiss S. 243  
 Koczka N. 189

### L

Lalic T. 95  
 Lalosevic D. 81

Lăcătușu R. 41  
Lăzureanu A. 195

## M

Marinescu G. 105  
Măruțoiu C. 195  
Meșter E.L. 249  
Miklasova N. 141  
Moigrădean D. 195  
Moisuc A. 169  
Moșoarcă E.M. 105  
Munteanu M.C. 231, 249

## N

Nagy A. 199  
Neagoe A. 73  
Nechifor M. 27  
Negrea A. 173  
Negrea P. 127, 173  
Neményi A. 199  
Némethy U.H. 211  
Niță L. 113  
Niță S. 113  
Nyéki J. 225

## O

Orosz F. 211

## P

Palamaru I. 205  
Papadopol V. 205  
Papp A. 67, 239  
Papp N. 225  
Patron L. 105  
Pedryc A. 225  
Perde M. 141  
Pétersz D. 189  
Petruse C. 163  
Podia-Igna C. 87  
Popa A. 127  
Popa E. 205  
Popescu C.I. 255  
Popovici D. 169  
Postolache C. 231  
Preda G. 277  
Princz E. 243

Pucarevic M. 55

## R

Rădulescu H. 243  
Roșoiu N. 277

## S

Sárközi L. 67  
Sayti L. 259  
Sava M. 87  
Sekulic S. 81  
Silaghi-Dumitrescu I. 5  
Silaghi-Dumitrescu L. 141  
Silaghi-Dumitrescu R. 5, 271  
Simescu M. 87  
Skoric T.Z. 55  
Slezák K. 211  
Socaciu C. 133  
Staicu A.C. 231, 249  
Stefanovits-Bányai E. 189, 199, 211  
225, 243  
Szabó Z. 225  
Szilvássy B. 225

## T

Takács Sz. 239  
Takacs-Hajos M. 55  
Tatomir C. 141  
Taubert L. 243  
Teodorescu D. 249  
Timcheva K. 105  
Tordai E. 225  
Tudose R. 105

## V

Varvari L. 255  
Velciov A.B. 259  
Velciov P.R. 259  
Virag P. 141  
Voroniuc O. 265

## Y

Yogi L. 133

## Z.

Zeremski Skoric T.


# RETROSPECTIVES

see website: [www.meemb-timisoara.ro](http://www.meemb-timisoara.ro)





**SERIES OF INTERNATIONAL SYMPOSIA ORGANIZED UNDER THE AUSPICES OF ROUMANIAN ACADEMY - BRANCH TIMIȘOARA WITH THE THEME "METAL ELEMENTS IN ENVIRONMENT, MEDICINE AND BIOLOGY" (M.E.E.M.B.)**



ISBN 973 – 578 – 045 - 3


**1. "Metal Elements in Environment, Medicine and Biology" – Tome I (312 p)**  
(Editors Drăgan Petru, Gârban Zeno)

Foreword by Prof. Mihai Șerban, PhD – corresponding member of Roumanian Academy, president of Comission of Biochemistry of Roumanian Academy București

*Proceedings of 1st International Symposium, October 21-23, 1993*

Organized under the auspices of Roumanian Academy – Branch Timișoara with collaboration of University of Medicine and Pharmacy Timișoara, University of Agricultural Sciences and Veterinary Medicine Timișoara

Publishing House "Mirton" Timișoara (First Edition - 1994, Second Edition - 1995)



ISBN 973 - 9336 -15 - 9


**2. "Metal Elements in Environment, Medicine and Biology" – Tome II (396 p)**  
(Editors Gârban Zeno, Drăgan Petru)

Foreword by Acad. Prof. Ionel Haiduc, PhD – vicepresident of Roumanian Academy București

*Proceedings of 2nd International Symposium, October 27-29, 1996*

Organized under the auspices of Roumanian Academy – Branch Timișoara and Comission of Biochemistry of Roumanian Academy București

Publishing House "Eurobit" Timișoara (First Edition - 1996, Second Edition - 1997)



**3. “Metal Elements in Environment, Medicine and Biology” – Tome III (392 p)**  
(Editors Gârban Zeno, Drăgan Petru)


Foreword by Acad. Prof. Toma Dordea, PhD, Eng. – president of Branch Timișoara – Roumanian Academy

*Proceedings of 3rd International Symposium, October 26-28, 1998*

Organized under the auspices of Roumanian Academy – Branch Timișoara and Comission of Biochemistry of Roumanian Academy București

Publishing House “Eurobit” Timișoara (1998)

ISBN 973 – 9336 – 19 – 1  
ISSN 1583 – 4204



**4. “Metal Elements in Environment, Medicine and Biology” – Tome IV (362 p)**  
(Editors Gârban Zeno, Drăgan Petru)


Foreword by Acad. Prof. Toma Dordea, PhD, Eng. – president of Branch Timișoara - Roumanian Academy

*Proceedings of 4<sup>th</sup> International Symposium, November 6-8, 2000*

Organized under the auspices of Roumanian Academy – Branch Timișoara and Comission of Biochemistry of Roumanian Academy București

Publishing House “Eurobit” Timișoara (2000)

ISBN 973 – 8181 – 39 – 9  
ISSN 1583 – 4204



**5. “Metal Elements in Environment, Medicine and Biology” – Tome V** (368 p)  
(Editors Gârban Zeno, Drăgan Petru, Gârban Gabriela)


Foreword by Acad. Prof. Toma Dordea, PhD, Eng. –  
president of Branch Timișoara - Roumanian Academy

*Proceedings of 5th International Symposium,  
November 4-6, 2002*

Organized under the auspices of Roumanian Academy –  
Branch Timișoara and Comission of Biochemistry of  
Roumanian Academy București

Publishing House “Eurobit” Timișoara (2002)

ISBN 973 – 620– 035– 3  
ISSN 1583 – 4204



**6. “Metal Elements in Environment, Medicine and Biology” – Tome VI** (366 p)  
(Editors Gârban Zeno, Drăgan Petru)

in collaboration with

Avacovici Adina, Davidescu Corneliu, Drăgan Simona,  
Gârban Gabriela, Lăzureanu Aurel, Vaszilcsin Nicolae


Foreword by Prof. Costin Cernescu, MD, PhD – corre-  
sponding member of the Roumanian Academy; Director of  
the “Ștefan Nicolau” Institute of Virology Bucuresti

*Proceedings of 6th International Symposium,  
November 8-10, 2004*

Organized under the auspices of Roumanian Academy –  
Branch Timișoara and Comission of Biochemistry of  
Roumanian Academy București

Publishing House “Eurobit” Timișoara (2004)

ISBN 973– 620– 119– 8  
ISSN 1583 – 4204



**7. “Metal Elements in Environment, Medicine and Biology” – Tome VII (356 p)**

(Editors Gârban Zeno, Drăgan Petru)

in collaboration with

Davidescu Corneliu, Drăgan Simona, Gârban Gabriela,  
Gergen Iosif, Vaszilcsin Nicolae

Foreword by Prof. Costin Cernescu, MD, PhD – corresponding member of the Roumanian Academy; Director of the “Ștefan Nicolau” Institute of Virology București

*Proceedings of 7th International Symposium,  
November 6-8, 2006*

Organized under the auspices of Roumanian Academy – Branch Timișoara and Comission of Biochemistry of Roumanian Academy București

Publishing House “Eurobit” Timișoara (2006)


ISBN (10) 973-620-238-0

ISBN (13) 978-973-620-238-4

ISSN 1583-4204

***IN MEMORIAM***



**PROF. PETRU DRĂGAN, MD, PhD****Member of the Academy of Medical Sciences****February 2, 1932 - November 12, 2007**

On November 12th passed away Prof. Petru Drăgan, MD, PhD. distinguished personality of medical sciences – with remarkable contributions in the domain of urological surgery, general surgery and the application of molecular medicine in urology.

Prof. Petru Drăgan had a prodigious and prestigious professional activity working at the University Clinic of Urology of the Faculty of Medicine – University of Medicine and Pharmacy “Victor Babeș” Timișoara.

Born on February 2, 1932 in town Jimbolia - Timiș county, he started his studies in a school from his native town and continued at the “C.D. Loga” high school in Timișoara (graduated in 1951). His university studies were performed at the Faculty of General Medicine – University of Medicine and Pharmacy Timișoara (graduated in 1957).

At the beginning of his professional career Petru Drăgan, MD worked as a general practitioner physician in town Bocșa Română (1957-1959). Then, after a competition, he occupied a job of assistant professor in the University Clinic of Urology Timișoara – Faculty of Medicine. His doctoral thesis had as subject “Intrarenal circulation in shock”. He achieved, successively, all the university degrees up to university professor (1990).

During his qualification he attended numerous specializations in Romania and abroad. Among them one can mention his specialization in U.S.A. at the University of California Los Angeles (UCLA) – Department of Urology (1969-1970) with a Fulbright scholarship. At UCLA he worked under the direction of Prof. W. Goodwin, MD, PhD and Prof. Kaufmann, MD, PhD.

During the period of specialization in U.S.A. he sustained and graduated all the exams – very rarely in case of Fulbright’ scholars - equalizing the Diploma of physician revealing the high degree of his competence.

Returned in Romania he contributed to the development of the «School of Urology» from Timișoara, dedicating for this purpose a part of his existence and offering it a world-wide prestige.

In this framework there are mentioned some of his essential contributions:

- introducing of over 15 new surgical interventional technics in urology;
- promoting endourology in Timișoara and later on being taken over in Romania
- performing the first renal transplant with kidneys from corpse (1981) – being a pioneer of renal transplant in Romania
- contributing to the development of concepts and applications of Molecular medicine in Urology in Romania (starting from 1980). In this context there are mentioned

contributions on molecular mechanisms of the heterogenous nucleation in uroconcrements biogenesis.

- publishing as first author or editor of over 15 books with monographic character
- author of over 200 scientific papers published in various specialty journals in Romania and abroad
- editor of seven volumes from the Series of International Symposia “Metal Elements in Environment Medicine and Biology” (initiated in 1993)
- coordinator, as chief editor of the journal «Timișoara Medicală» – official periodical of the University of Medicine and Pharmacy “Victor Babeș” Timișoara
- laid the basis of the “Urological Foundation Timișoara” – under its auspices were organized some National Symposia of Urology
- as a real leader of the School of Urology Timișoara he trained over 60 specialists in urology and as a mentor he had the capacity to coordinate, to stimulate and to promote valuable personalities in urology who are working now either in Romania or abroad – in different European states
- as advisor for doctoral thesis in medicine he coordinated the elaboration of numerous doctoral thesis by young physicians, approaching complex surgical, physiological, biochemical, physiopathological, histological problems with application in urology

Recognition of his praiseworthy activity was also proved by his affiliation to various *national scientific societies*, e.g.: Romanian Association of Urology; Romanian Association of Endourology (as founding member); Romanian Society of Biochemistry (by his contributions in Molecular medicine); National Council for Prostate Diseases (founding member); Society of Urologists from Moldavia a.o. and *international scientific societies*, e.g.: European Association of Urology; Britain Association of Urological Surgeons; Society of Urology from South-East Germany; American Association of Urology a.o.

Regarding his professional activity one must mention that he worked as professor at the University of Monrovia – Liberia (1975-1976) being also the private urologist of William Tolbert Jr. – president of the state Liberia.

Merits of Prof. Dr. Petru Drăgan were recognized at national level, too, being rewarded with various state prizes such as : Work Medal Class III (1979); National medal „For Merit” in Great Officer rank (2000).

At local level he was awarded with the titles : Honorary Citizenship of Jimbolia; Honorary Citizenship of Timișoara (2002).

He was titular member of the Academy of Medical Sciences and of Romanian Academy of Scientists.

A wide culture in the domain of history - from Antiquity to Contemporaneity, in the domain of arts, history of culture, economical geography, prove the huge assimilation capacity of a brilliant scientific and cultural mind.

He was appreciated by Acad. Nicolae Cajal, Acad. Pius Brânzeu, Acad. Ionel Haiduc, Acad. Toma Dordea – personalities who honoured by their presence the scientific meetings where Prof. Dr. Drăgan was the chairman.

Many times the scientific preoccupations and, especially, those related to medical practice did not permit him to elaborate manuscripts but each discussion with his students and friends was an intellectual revelation.

Passing into eternity of Prof. Petru Drăgan, MD, PhD – a well known personality, is a great loss not only for the medical sciences but also for the Romanian science and culture.



## LIST OF PUBLICATIONS TEATISES, BOOKS, UNIVERSITY COURSES

**PROF. PETRU DRĂGAN MD, PhD**

1. **Curs de urologie** - în colab. cu Ștefanovici A.B., David M., Litografie Institutul de Medicină Timișoara, 1979
2. **Urologie – curs**, Litografie Institutul de Medicină Timișoara, 1979
3. **Curs de urologie** - în colab. cu Miclea F., Plăcintă A., Litografie Institutul de Medicină Timișoara, 1979
4. **Metal Elements in Environment, Medicine and Biology, Tome I** - in collab. with Gârban Z., Romanian Academy – Branch Timisoara, Proceedings of the 1st International Symposium, Timișoara, October 21-23, 1993, Publishing House “Mirton” Timișoara, 1994
5. **Cancerul prostatei** - în colab. cu Miclea F., Raica M., Editura Almanahul Banatului, Timișoara, 1996
6. **Metal Elements in Environment, Medicine and Biology, Tome II** - in collab. with Gârban Z., Romanian Academy – Branch Timisoara, Proceedings of the 2nd International Symposium, Timisoara, October 27-29, 1996, Publishing House “Eurobit” Timișoara 1996
7. **Afecțiuni ale aparatului urogenital inferior. Tratamentul chirurgical endoscopic** - în colab. cu Miclea F., Boiboreanu P., Almanahul medical, Editura Filantropia și Europa Nova, 1996
8. **Tratamentul endoscopic al tumorilor vezicale** (pp.141-161) - în colab. cu Bucuraș, Botoca M., publicat în “Bazele chirurgiei endourologice” (editor Nicolescu D.), Editura Eurobit Timișoara 1997.
9. **Metal Elements in Environment, Medicine and Biology, Tome III** - in collab. with Gârban Z., Romanian Academy – Branch Timisoara, Proceedings of the 3rd International Symposium, Timișoara, October 27-29, 1998, Publishing House “Eurobit” Timișoara 1998
10. **Urgențe chirurgicale, Vol.1** - în colab. cu Nica C., Bota N., Costea C., Miclea F., Nediglea I., Nica C., Nica C., Nodiți Gh., Poenaru D., Vâlceanu D., Editura „Mirton” Timișoara, 1999.
11. **Metal Elements in Environment, Medicine and Biology, Tome IV** - in collab. with Gârban Z., Romanian Academy – Branch Timisoara Proceedings of the 4th International Symposium, Timișoara, November 6-8, 2000, Publishing House “Eurobit” Timișoara 2000
12. **Adenomectomia Millin**, publicat în “Tratat de tehnici chirurgicale urologice” (editor Lucan M.), Editura Infomedica București 1996 (ed. 1-a), Editura Infomedica București 2001 (ed. 2-a).
13. **Metal Elements in Environment, Medicine and Biology, Tome V** - in collab. with Gârban Z., Gârban Gabriela, Romanian Academy – Branch Timisoara, Proceedings of the 5th International Symposium, Timișoara, November 4-6, 2002, Publishing House “Eurobit” Timișoara 2002

14. **Metal Elements in Environment, Medicine and Biology, Tome VI** - in collab. with Gârban Z., Romanian Academy – Branch Timisoara, Proceedings of the 6th International Symposium, Timișoara, November 8-10, 2004, Publishing House “Eurobit” Timișoara 2004
15. **Metal Elements in Environment, Medicine and Biology, Tome VII** - in collab. with Gârban Z., Romanian Academy – Branch Timisoara, Proceedings of the 7th International Symposium, Timișoara, November 6-8, 2006, Publishing House “Eurobit” Timișoara 2006

## PUBLISHED AND COMMUNICATED PAPERS

PROF. PETRU DRĂGAN MD, PhD

1. *Aplicații ale spectroscopiei în infraroșu la decelarea expeditivă a compoziției calculilor renali* - în colab. cu Gârban Z., Szabadai Z., Cristescu Lucia, Stoicoiu Maria, Timișoara Medicală, 1981, XXVI, 4, 41-48.
2. *Considerații asupra mecanismului bioanorganic în patogeniza urolitiazelor purinice* - în colab. cu Gârban Z., Cristescu Lucia, Mihalca Victoria, Timișoara Medicală, 1981, XXVI, 4, 54-61.
3. *Utilizarea spectroscopiei de absorbție atomică la decelarea compoziției cationice a calculilor urinari - Notă preliminară* - în colab. cu Gârban Z., Nemeș Gisela, Martoni Livia, Daranyi Gabriela, în "A XII-a Sesiune Stiintifică a cadrelor didactice", Institutul de Medicină Timișoara, Volum Rezumate, p.79, Timișoara 15 mai 1982.
4. *Posibilități de utilizare a calculatorului la interpretarea spectrelor în infraroșu pentru urolitiază* - în colab. cu Gârban Z., Ciubotariu D., Holban Șt., Laslău A., Mihalca Victoria, Cristescu Lucia, Timișoara Medicală, 1983, XXVIII, 1, 13-18.
5. *Microcalculatoarele aplicate la interpretarea spectrelor în infraroșu pentru determinări de compoziție în urolitiază* - în colab. cu Gârban Z., Holban Șt., Ciubotariu D., Nica Lucia, Giurgiulescu Elena, Sayti L., Cristescu Lucia, în "Al VII-lea Simpozion de informatică medicală", pp.93-94, Timișoara 8-9 iunie 1984.
6. *Tehnici de recunoaștere a formelor utilizate pentru evaluarea spectrelor în infraroșu la investigarea urolitiazelor* - în colab. cu Vancea R., Ciubotariu D., Holban Șt., Gârban Z., Gogonea V., Manu Rodica, Panaitescu Valeria, în "Al VII-lea Simpozion de informatică medicală", pp. 95-96, Timișoara 8-9 iunie 1984.
7. *Considerations on the metal composition of the calculi in urolithiasis* - in collab. with Gârban Z., Nemeș Gisela, Daranyi Gabriela, Popeti Doina, Martoni Livia, in "Meeting of the Federation of European Biochemical Sciences: Metal Ions, Proteins and Membranes", Abstracts Volume, p.81, Algarve, April 21-26, 1985.
8. *Application of the computer in the interpretation of the infrared spectra to determine the composition in urolithiasis* - in collab. with Gârban Z., Ciubotariu D., Holban Șt., Giurgiulescu Elena, "Baki. Hamaabada", Israel J. Clin. Biochem. and Lab. Sci., 1985, 4(1-3), 12.
9. *Investigations on urolithiasis by means of infrared spectroscopy and computer* - in collab. with Gârban Z., Ciubotariu D., Giurgiulescu Elena, Holban Șt., Bucuraș V., Rev. roum. Biochim., 1986, 23 (3), 161-169.
10. *Considerations concerning the etiopathogeny of urolithiasis and the composition of the spectroscopically investigated calculi. II. Oxalic lithiasis* - in collab. with Gârban Z., Nemeș Gisela, Giurgiulescu Elena, Mihalca Victoria, Timișoara Medicală, Supplement, 1986, XXXI, 10-19.
11. *Biochemical studies on some metals present in urolithiasis. II. Determination of the main alkaline and alkaline-earth metals* - in collab. with Gârban Z., Nemeș Gisela, Daranyi Gabriela, Sigartău Gr., Șișu E., Mureșan Codruța, Rev. roum. Biochim., 1987, 24 (3), 221-227.
12. *Spectroscopy and computer in the investigation of urolithiasis* - in collab. with Gârban Z., Ciubotariu D., Giurgiulescu E., Nemeș G., Holban Șt., in "18<sup>th</sup> FEBS Meeting", Abstracts Volume, p.181, Ljubljana, June 28-July 3, 1987.

13. ***Oxalic and phosphatic urolithiasis investigated by spectrometric methods*** - in collab. with Gârban Z., Neamțu Lucia, Nemeș Gisela, Giurgiulescu Elena, "14<sup>th</sup> International Congress of Biochemistry", Abstracts, Vol.II, p.157, Prague, July 10-15, 1988.
14. ***Implicații ale interacției acidului deoxiribonucleic și a unor derivați purinici cu ioni metalici*** - în colab. cu Gârban Z., Văcărescu G., Eremia Iulia, Bucuraș V., Precob V., in "Al III-lea Congres Național de Chimie", Rezumate, Vol. III, pp. 811-812, București 21-24 septembrie 1988.
15. ***Diagnosticul urolitiazelor purinice cu ajutorul spectroscopiei în infraroșu*** - în colab. cu Gârban Z., Miclea F., Ciubotariu D., Peter F., Mureșan Codruța, Miloș M., în "A XV-a Sesiune Stiințifică a cadrelor didactice", Institutul de Medicină Timișoara, Volum Rezumate, p.54, Timișoara 12 mai 1989.
16. ***Investigations on the metal composition and etiopathogeny of urolithiasis*** - in collab. with Gârban Z., Ciubotariu D., Nemeș G., "Sixth International Symposium on Molybdenum, Vanadium and Other Trace Elements", P.XVIII.61, 1-9, Jena, July 3-6, 1989.
17. ***Investigations on the composition and etiopathogeny of lithiasis. I. Purine urolithiasis*** - in collab. with Gârban Z., Ciubotariu D., Peter F., Mureșan Codruța, Miloș M., Rev.roum. Biochim., 1989, 26 (3), 201-208.
18. ***Comparative data concerning magnesium and calcium in simple urolithiasis*** - in collab. with Gârban Z., Daranyi Gabriela, Nemeș Gisela, Mureșan Codruța, Manu Rodica, in "International Congress on Magnesium Research", Abstracts Volume, p.12, Târgu Mureș, September 21-24, 1989.
19. ***Investigations on the composition and etiopathogeny of lithiasis. III. Phosphatic urolithiasis*** - in collab. with Gârban Z., Bordoș D., Peter F., Martoni Livia, in "20<sup>th</sup> FEBS Meeting", Abstracts Volume, p.146, Budapest, August 19-24, 1990.
20. ***Etudes biochimiques sur quelques métaux presents dans les urolithiases. II. Determination de quelques métaux transitionnels*** - avec collab. des Gârban Z., Daranyi Gabriela, Nemeș Gisela, Șişu E., Șoșdean Doina, Martoni Livia, Rev. roum. Biochim., 1990, 27 (3), 209-216.
21. ***The effects of cis-platinum and cyclophosphamide on deoxyribonucleic acid*** - in collab. with Gârban Z., Daranyi Gabriela, Riviș I., Mareș A., Eremia Iulia, in "XI<sup>th</sup> International Symposium on Medicinal Chemistry", Abstracts Volume, p.52, Jerusalem, September 2-7, 1990.
22. ***Investigations on the composition and etiopathogeny of lithiasis. VII. Some metal trace elements in simple urolithiasis*** - in collab. with Daranyi Gabriela, Matusz-Vâlceanu Anca, Peter F., Bene Cristina, Gârban Z., Publ. in "Review of trace- and macroelement research", Cluj-Napoca, 1992, 1, 67.
23. ***Investigații asupra compoziției și etiopatogeniei litiazelor. IX. Particularități ale urolitiazelor la vârste tinere*** - în colab. cu Daranyi Gabriela, Șoșdean Doina, Negrea Gertrude, Gârban Z., Popescu Delia, Analele Institutului de Sănătate Publică și Cercetări Medicale, Timișoara 1992, XVIII, 1 (10), 76-79.
24. ***Investigations on the composition and etiopathogeny of lithiasis. X. Alkaline and alkaline-earth metals in mixed urolithiasis*** - in collab. with Daranyi Gabriela, Gârban Z., Sayti L., in "Proceedings of the Third Symposium of the Romanian Society of Biominerals", Programme and Abstracts Volume, p.9, Odorheiu Secuiesc, October 1-3, 1992.
25. ***Investigations on the composition and etiopathogeny of lithiasis. XI. Alkaline and alkaline-earth metals in uroconcrements and biochemical homeostasis*** - in collab. with Gârban Z., Daranyi Gabriela, Peter F., Popescu Delia, Timișoara Medicală, 1993, XXXVIII, (1-2), 43-46.

26. ***Investigations on the composition and etiopathogeny of lithiasis. XII. Determination of some divalent metals in simple urolithiasis*** - in collab. with Daranyi Gabriela, Gârban Z., Peter F., pp. 904-905, in "Proceedings of the 8<sup>th</sup> International Symposium on Trace Elements in Man and Animals", Dresden-Germany, May 16-21, 1993 (Eds. Anke M., Meissner I.D., Mills C.F.), Verlag Media Touristik, Gersdorf, 1993.
27. ***Investigations on the composition and etiopathogeny of lithiasis. XV. Calcium and magnesium homeostasy in urolithiasis*** - in collab. with Daranyi Gabriela, Gârban Z., Șoșdean Doina, Popescu Delia, Annals of the Institute of Public Health and Medical Research, Timișoara, New Series 1993, XIX, 2 (11) 301-304.
28. ***Investigations on the composition and etiopathogeny of lithiasis. XVI. Metallic composition of cystinic uroconcrements*** - in collab. with Daranyi Gabriela, Bucuraș V., Matusz Anca, Papoe Gabriela, Martoni Livia, Gârban Z., pp.77-80, Metal Elements in Environment, Medicine and Biology, Tome I (Eds. Drăgan P., Gârban Z.), Publishing House Mirton Timișoara, 1994.
29. ***Investigations on the composition and etiopathogeny of lithiasis. XVII. Metallic bioelements implicated in molecular pathology - Reference to the pathobiology of urolithiasis*** - in collab. with Gârban Z., Miclea F., Holban Șt., pp.85-90, Metal Elements in Environment, Medicine and Biology, Tome I (Eds. Drăgan P., Gârban Z.), Publishing House Mirton Timișoara, 1994.
30. ***Investigations on the composition and etiopathogeny of lithiasis. XVIII. Dishomeostasy of some alkaline and alkaline-earth metals in pre-, postsurgical and ESWL treatment of urolithiasis*** - in collab. with Daranyi Gabriela, Bucuraș V., Agapie D., Botoca M., Patroi Delia, Gârban Z., pp.91-94, Metal Elements in Environment, Medicine and Biology Tome I (Eds. Drăgan P., Gârban Z.), Publishing House Mirton Timișoara, 1994.
31. ***Investigations on the composition and etiopathogeny of lithiasis. XXIII. Sodium and potassium homeostasis in patients with urolithiasis*** - in collab. with Daranyi Gabriela, Șoșdean Doina, Gârban Z., Annals of the Institute of Public Health and Medical Research, Timișoara, New Series 1994-1995, XX, 3 (12), 91-96.
32. ***Investigations on the composition and etiopathogeny of lithiasis. XXVI. Investigations concerning the magnesium, calcium, zinc and copper concentration in binary urolithiasis*** - in collab. with Daranyi Gabriela, Gârban Z., Riviș A., in "6<sup>th</sup> Magnesium Symposium", Abstracts Volume, p.79, Balatonszéplak, June 17-19, 1996.
33. ***Investigations on the composition and etiopathogeny of lithiasis. XXVII. Thermal analysis and atomic absorption spectroscopy in the identification of urolithiasis*** - in collab. with Gârban Z., Daranyi Gabriela, Sigartău Gr., Avacovici Adina, Urzică A., pp.307-312, in New Perspectives in the Research of Hardly Known Trace Elements and Their Role in Life Science, (Ed. Pais I.), University Press of U.H.F. Budapest, 1996.
34. ***Investigations concerning the composition and etiopathogeny of urolithiasis*** - in collab. with Daranyi Gabriela, Gârban Z., Annals of the Institute of Public Health and Medical Research, Timișoara, New Series 1996, XXI, 4 (13), 51-59.
35. ***Purine urolithiasis: peculiarities of the etiopathogeny and composition of uroconcrements*** - in collab. with Gârban Z., pp.89-98, in Metal Elements in Environment, Medicine and Biology, Tome II (Eds. Gârban Z., Drăgan P.), Publishing House Eurobit Timișoara, 1996.
36. ***Research on urolithiasis composition by infrared spectroscopy, thermal analysis and atomic absorption spectroscopy*** - in collab. with Daranyi Gabriela, Gârban Z., Sigartău Gr., Avacovici Adina, Sarafolean S., pp.139-144, in Metal Elements in Environment,

- Medicine and Biology, Tome II (Eds. Gârban Z., Drăgan P.), Publishing House Eurobit Timișoara, 1996.
37. ***Investigations on the composition and etiopathogeny of lithiasis. XXVIII. Status of blood and urinary metallogram in relationship with ESWL treatment in urolithiasis*** - in collab. with Gârban Z., Daranyi Gabriela, Bucuraș V., Botoca M., Minciu R., Urzică A., Patroi-Popescu Delia, Ehling Șt., pp.173-176, Metal Elements in Environment, Medicine and Biology, Tome II (Eds. Gârban Z., Drăgan P.), Publishing House Eurobit Timișoara, 1996.
  38. ***Methodological principles in the computerized evaluation of the infrared spectra of some urolithiasis and metal composition of uroconcrements*** - in collab. with Holban Șt., Gârban Z., Holban Nina, Fântână N., Peter F., Găzdoiu Șt., Urzică H., Sarafolean S., pp.237-242, in Metal Elements in Environment, Medicine and Biology, Tome II (Eds. Gârban Z., Drăgan P.), Publishing House Eurobit Timișoara, 1996.
  39. ***Cytologic regression index in hormone sensitive and hormone sensitive carcinoma of the prostate*** – in collab. with Raica M., Bucuraș V., Herman I. „Progress and Controversies in Oncological Urology IV”, Rotterdam, Nederlanden, April 11-13, 1996
  40. ***Hormone-insensitive carcinoma of the prostate: is it predictable? A histochemical and imunohistochemical retrospective analysis*** – in collab. with Bucuraș V., Raica M., Miclea F. „14th World Congress on Endourology and SWL”, Melbourne, Australia, November 11-14, 1996
  41. ***Tratamentul calculilor ureterali lombari prin abord percutanat*** - în colab. cu Boiborean P., Miclea F., Bucuraș V., Botoca M., Claici D., Suflea D. „Al VI-lea Simpozion Național de Endourologie”, Oradea-Felix 1-4 mai 1996
  42. ***Rezultate imediate și tardive ale uretrotomiei interne pentru stricturi de uretrale*** - în colab. cu Boiborean P., Miclea F., S. Ghișoiu, Pupca G. „Al VI-lea Simpozion Național de Endourologie”, Oradea-Felix, 1-4 mai 1996
  43. ***Strategia tratamentului endoscopic al stricturilor uretrale complexe*** - în colab. cu Boiborean P., Miclea F., Minciu R., Jahansouz S. „Al VI-lea Simpozion Național de Endourologie”, Oradea-Felix 1-4 mai 1996
  44. ***Nefrostomia percutanată, prim gest terapeutic în anuria obstructivă*** - în colab. cu Miclea F., Boiborean P., Claic D., Herman I., Librimir C. „Al VI-lea Simpozion Național de Endourologie”, Oradea-Felix 1-4 mai 1996
  45. ***Iatrogenoze ginecologice întâlnite în practica urologică*** - în colab. cu Claici D., Almăjan G., Librimir C. „Al VI-lea Congres Național de Urologie”, Sinaia, 25-28 septembrie, 1996
  46. ***Valoarea metodelor morfo și imunohistochimice pentru predicția hormonorezistenței în cancerul de prostată*** - în colab. cu Raica M., Bucuraș V., Botoca M. „Al VI-lea Congres Național de Urologie”, Sinaia, 25-28 septembrie 1996
  47. ***Peculiarities of the relationship nutrition - metaphylaxis of urolithiasis. I. Nutritional aspects in oxalic urolithiasis*** - in collab. with Daranyi Gabriela, Gârban Z., Annals of the Institute of Public Health and Medical Research, Timișoara, New Series 1997, XXII, 5 (14), 61-65.
  48. ***Valoarea terapeutică a ciprofloxacinului în infecțiile urinare înalte - studiu clinicobiologic*** - în colab. cu Miclea F., Bucuraș V., Ciobănaș D., Popa R. „Al 7-lea Simpozion Național de Endourologie cu participare internațională”, Timișoara, 7-10 mai 1997

49. ***Ureteroscopia retrogradă pentru calculoza versus ureterolitotomie*** - în colab. cu Boiborean P., Miclea F., Bucuras V., Minciu R., Pupca Gh. 7-lea Simpozion Național de Endourologie cu participare internațională, Timișoara 7-10 mai 1997
50. ***The relationship between immunohistochemical expression of Prostate Specific Antigen and Gleason score in prostate carcinoma*** – in collab. with Miclea F., Boiborean P., Bucuraș V. „15th World Congress on Endourology and SWL”, Edinburgh, United Kingdom, August 31- September 3, 1997
51. ***The Relationship Between the Immunohistochemical Expression of Prostate Specific Antigen (PSA) and Gleason Score in Prostate Carcinoma*** - Bucuraș V., Raica M., Miclea F. „15th World Congress on Endourology and SWL”, Edinburgh, United Kingdom, August 31 – September 3, 1997
52. ***Conservative surgical treatment in urothelial tumors of the upper urinary tract*** - in collab. with Miclea F., Minciu R., Botoca M., Pupca Gh. „24<sup>th</sup> World Congress, 7-11 September 1997.
53. ***Immunohistochemistry of estrogen and androgen receptors in carcinoma of the prostate*** – in collab with Miclea F., Raica M., Dema A. „24<sup>th</sup> World Congress”, SIU Montreal, September 7-11, 1997.
54. ***Immunohistochemistry of PSA in prostate carcinoma. A relationship with Gleason score*** - in collab. with Raica M., Miclea F., Dema A., Toma M., Herman I. „24<sup>th</sup> World Congress”, SIU Montreal, September 7-11, 1997.
55. ***Advantages of the direct internal urethrotomy – un guided – in posterior urethral strictures*** - in collab. with Miclea F., Raica M., Bucuraș V., Herman I. „The Xth Congress of the Hungarian Society of Urology”, Debrecen, Hungary, October 16-18, 1997
56. ***Metallograms in simple and mixed urolithiasis*** - in collab. with Gârban Z., Daranyi Gabriela, Avacovici Adina, J. Trace Elements in Experimental Medicine, 1998, 11, 376.
57. ***Clinical aspects and biochemical pathology in cystinic urolithiasis - peculiarities of the metallic composition*** - in collab. with Gârban Z., Holban Șt., pp. 27-34, in Metal Elements in Environment, Medicine and Biology, Tome III (Eds. Gârban Z., Drăgan P.), Publishing House Eurobit Timișoara, 1998
58. ***TURP versus chirurgie deschisă pentru adenoame periuretrale la pacienții peste 80 de ani*** - în colab. cu Herman I., Minciu R., Corcan J. „Al VIII-lea Simpozion Național de Endourologie cu participare internațională”, București, 6-9 mai 1998
59. ***Epidemiology of kidney tumors in our region*** – „Urologia Triangularis”, Szeged, Hungary, May 14-15, 1998 ***Surgery of renal cell carcinoma. Which approach is “of choice”?*** - „Urologia Triangularis”, Szeged, Hungary, May 14-15, 1998
60. ***Is conservative surgical treatment of renal tumors a fair option ?*** - „Urologia Triangularis”, Szeged, Hungary, May 14-15, 1998
61. ***Conservative surgical treatment of multiple tumors in solitary kidney*** – Bucuraș V., Covaciu Alina, Bardan R., Corcan J. „Urologia Triangularis”, Szeged, Hungary, May 14-15, 1998
62. ***Valoarea scorului simptomatologic prostatic IPSS în stabilirea opțiunii terapeutice*** - în colab. cu Botoca M., Bardan R., Mureșan C. „Al VIII-lea Congres Național de Urologie”, Iași, 30 septembrie – 3 octombrie 1998
63. ***Tumorectomie multiplă pe rinichi unic chirurgical (Film Video)*** - în colab. cu Bucuraș V., Covaciu Alina, Bardan R., Corcan J. „Al VIII-lea Congres Național de Urologie”, Iași, 30 septembrie – 3 octombrie 1998

64. ***Experiența Clinicii Urologice Timișoara în tratamentul chirurgical al afecțiunilor suprarenalei*** - în colab. cu Miclea F., Bucuraș V., Covaciu A. „Al VIII-lea Congres Național de Urologie”, Iași, 30 septembrie – 3 octombrie 1998
65. ***Surgical treatment in congenital obstructing urological malformations – personal experience*** - în colab. cu Miclea F., Boiborean P. „Symposium of Pediatric Urology”, București 28 –30 octombrie 1998
66. ***Conservative Surgical treatment of urotelial upper urinary tract*** - UCLA Urological day, Los Angeles, USA, 05.06.1998
67. ***Internal optical urethrotomy, 17 years of experience*** - in collab. with Boiborean P., Miclea F., Minciu R., Botoca R., Bucuraș V., Herman I., Pupca Gh., Claiți D. „16th World Congress on Endourology and SWL”, New York, September 3-6, 1998
68. ***Profile of metallograms in phosphatic uroconcrements*** - in collab. with Daranyi Gabriela, Bucuraș V., Avacovici Adina, Vincu Mirela, Gârban Z., pp. 888-895, in Mengen- und Spurenelemente, 19. Arbeitstagung 1999, Friedrich-Schiller-Universität Jena, (Hrsg. Anke M. et al.), Verlag Harald Schubert, Leipzig, 1999.
69. ***Complicațiile intraoperatorii și postoperatorii imediate în chirurgia endoscopică a tumorilor vezicale*** - în colab. cu Bucuraș V., Herman I., Georgescu M., Shokri J. „Simpozionul Național de Endourologie”, Oradea-Felix 12-15 mai 1999
70. ***TUR – P la adenoamele peste 100g*** - în colab. cu Boiborean P., Miclea F., Herman A., Minciu R. „Simpozion Național de Endourologie”, Oradea-Felix 12-15 mai 1999
71. ***Valvă uretrală posterioară la copil – rezolvare endoscopică*** - în colab. cu Bardan A. „Simpozion Național de Endourologie”, Oradea-Felix 12-15 mai 1999
72. ***Tumora gonadica operata la un hermafrodit”(Film Video)*** - în colab. cu Bucuraș V., Bardan R., Corcan J. „Al IX-lea Congres Național de Urologie”, Sibiu 22-25 septembrie 1999
73. ***Leziuni ureterale In chirurgia cancerelor recto-colice*** - în colab. cu Miclea F., Bucuraș V., Botoca M., Braham A., Al IX-lea Congres Național de Urologie, Sibiu 22-25 septembrie 1999
74. ***Metalograma serica, urinara și a calculilor la pacienții cu litiaza urinara*** - în colab. cu Miclea F., Daranyi Gabriela, Garban Z., Bucuraș V. „Al IX-lea Congres Național de Urologie”, Sibiu 22-25 septembrie 1999
75. ***Fine needle aspiration biopsy in the follow-up patients with prostate cancer*** – in collab with Miclea F., Raica M., Bucuraș V., Herman I., Minciu R. „First Congress of the Central European Association of Urology”, Viena, Austria November 5-6, 1999
76. ***TUR-P for prostates over 100 grams*** – in collab. with Boiborean P., Miclea F., Herman I., Minciu R., Botoca M., Bucuraș V., Sandesc D. „First Congress of the Central European Association of Urology”, Viena, Austria, November 5-6 1999
77. ***Particular lesions associated with benign prostatic hyperplasia*** – in collab. with Demă A., Bucuraș V., Raica M., Botoca M., Herman I., Minciu R., Boiborean P., Miclea F., „First Congress of the Central European Association of Urology”, Viena, Austria, November 5-6 1999
78. ***Incidence of prostatic tumors in Timisoara County Hospital, Romania*** – in collab. with Bucuraș V., Demă A., Raica M., Botoca M., Herman I., Minciu R., Boiborean P., Miclea F. „First Congress of the Central European Association of Urology”, Viena, Austria, November 5-6, 1999
79. ***Concentration of magnesium in relation with other metals in serum, urine and uroconcrements of the surgically treated kidney stone patients*** - in collab. with Daranyi Gabriela, Gârban Z., Bucuraș V., Avacovici Adina, Popa M., Vincu Mirela, in



- “9<sup>th</sup> International Magnesium Symposium”, Abstract Volume, p.163, Vichy-France, September 10-15, 2000.
80. ***Clinical aspects and biochemical pathology in oxalic urolithiasis - peculiarities of the metallic composition*** - in collab. with Gârban Z., pp.45-58, in Metal Elements in Environment, Medicine and Biology, Tome IV (Eds. Gârban Z., Drăgan P.), Publishing House Eurobit, Timișoara, 2000.
  81. ***Profile of uroconcrements metallogram in simple lithiasis with oxalates, phosphates and urates*** - in collab. with Daranyi Gabriela, Bucuraș V., Gârban Z., Vincu Mirela, Popescu Georgeta-Sofia, pp.1101-1108, in Mengen- und Spurenelemente, 20. Arbeitstagung 2000, Friedrich-Schiller-Universität Jena, (Hrsg. Anke M. et al.), Verlag Harald Schubert, Leipzig, 2000.
  82. ***Bladder preservation. Conservative surgery and TUR in T2-T3 bladder tumors*** - în colab. cu Miclea F., Boiborean P., Minciu R., Bucuras V., Botoca M. „4th Urologia Triangularis”, Timișoara 24-25 martie 2000
  83. ***Curba de invatare in resectia transuretrala a adenomului de prostata. Experienta Clinicii Urologice Timisoara*** - în colab. cu Claici D., Bahram A., Botoca M., Bucuras V., Georgescu M., Herman I., Minciu R., Pupca Gh. „Al X-lea Simpozion National de Endourologie”, Targu-Mures 10-13 mai 2000
  84. ***Nefrolitotomia percutana. Impactul asupra tratamentul litiazei coraliforme*** - în colab. cu Miclea F., Boiborean P., Botoca M., Bucuras V., Minciu R., Herman I., Claici D., Pupca Gh., Georgescu M., Cumanas A., El Houbi M. „Al X-lea Simpozion National de Endourologie”, Targu-Mures 10-13 mai 2000, Targu-Mures
  85. ***Traumatismele ureterale dupa interventiile chirurgicale din obstetrica si ginecologie*** - în colab. cu Miclea F., Boiborean P., Bucuras V., Botoca M. „Al X-lea Congres National de Urologie”, București 27-29 septembrie 2000
  86. ***Nefrectomia in cancerul renal (Film Video)*** - în colab. cu Bucuras V., Botoca M., Bardan R., Muresan A., Miclea F. „Al X-lea Congres National de Urologie”, Bucuresti, 27-29 septembrie 2000
  87. ***Rata de proliferare in neoplazia intraepiteliala prostatica (PIN)*** - în colab. cu Raica M., Toma M., Dema A., Miclea F., Bucuras V. „Al X-lea Congres National de Urologie”, București 27-29 septembrie 2000
  88. ***Vasculita sistemica ce intereseaza prostata si testicolul*** - în colab. cu Dema A., Bucuras V., Tudose N., Taban Sorina, Alexandrescu Lavinia „Al X-lea Congres National de Urologie”, București 27-29 septembrie 2000
  89. ***Malignant dysgerminoma in a true hermaphrodite (Video)*** – in collab. with Bucuras V., Bardan R., Corcan J. „XVth Congress of the European Association of Urology”, Bruxelles, Belgie, April 12-15, 2000
  90. ***Immunohistochemistry in the diagnosis of hormone-insensitive prostate carcinoma*** - in collab. with Raica M., Miclea F., Bucuras V., Botoca M. „Second Congress of the Balkan's Urologists”, Struga, Macedonia, May 25–27, 2000
  91. ***Late complications after TUR*** – P - V. Bucuraș, Gh. Pupca, Botoca M. „Second Congress of the Central European Association of Urology (CEAU)”, Budapest, Hungary, October 13-14, 2000
  92. ***Increasing role of percutaneous approach in the management of staghorn calculi*** – in collab. wiht Miclea F., Boiborean P., Botoca M., Bucuraș V., Minciu R., Herman I., Claici D. „Second Congress of the Central European Association of Urology (CEAU)”, Budapest, Hungary, October 13-14, 2000

93. ***Management of ruptured posterior urethra – Modalities and results*** – in collab. with Boiborean P., Miclea F., Bucuras V., Botoca M., Minciu R. „Congress of the International Society of Urology”, Singapore, China, October 2000
94. ***Management of staghorn calculi: percutaneous vs. Surgical approach*** - in collab. with Miclea F., Bucuraş V., Mureşan A. „Urologia Triangularis”, Novi Sad, Yugoslavia, 2001
95. ***Ureteroscopy in the treatment of ureteric stones: can better equipment lead to better results?*** - in collab. with Botoca M., Herman I., Corcan J. “Urologia Triangularis “, Novi Sad, Yugoslavia, 2001
96. ***ESWL-Metodă terapeutică de elecție în litiaza renală*** - în colab. cu Miclea F., Boiboreanu P., Botoca M., Bucuraş V., Claici D., Herman I., Pupca Gh., Cumpănaş A., Bardan R. „Simpozionul național de ESWL”, Constanța 1-3 iunie 2001
97. ***New data regarding the metallogram of the patients with urolithiasis before and after surgical and ESWL treatment*** - in collab. with Gârban Z., pp.77-82, in Metal Elements in Environment, Medicine and Biology, Tome V (Eds. Gârban Z., Drăgan P.), Publishing House Eurobit, Timișoara, 2002.
98. ***Magnesium concentration in purinic uroconcrements*** - in collab. with Gârban Gabriela, Gârban Z., Avacovici Adina, Popa M., Simionică E., Popescu Sofia-Georgeta, pp.191-199, in Magnesium involvements in biology and pharmacotherapy (Eds. Nechifor M., Porr P.J.), Casa Cărții de Știință, Cluj-Napoca, 2003.
99. ***Oxalate urolithiasis: peculiarities of the etiopathogeny and composition of uroconcrements*** - in collab. with Popescu Sofia-Georgeta, Gârban Z., Gârban Gabriela, Ionescu I., pp.453-470, in Trace Elements in Human: New Perspectives, Vol. I (Eds. Ermidou-Pollet Sophie, Pollet S.), Printed by Entypossis, Athens, 2003.
100. ***Serum and urinary magnesium and calcium concentration in kidney stone patients before and after extracorporeal shock-wave lithotripsy*** - in collab. with Gârban Gabriela, Popescu Sofia-Georgeta, Gârban Z., Ionescu I., Martău Ariana-Bianca, Avacovici Adina, in “8<sup>th</sup> European Magnesium Congress”, Abstract Volume, p.64, Cluj-Napoca, May 25-28, 2004.
101. ***Oxalic urolithiasis: peculiarities of etiopathogeny and metallograms*** - in collab. with Gârban Z., pp. 121-130, in Metal Elements in Environment, Medicine and Biology, Tome VI (Eds. Gârban Z., Drăgan P.), Publishing House Eurobit, Timișoara, 2004.
102. ***Alkaline metals in serum and urine and urolithiasis disease*** - in collab. with Popescu Sofia, Gârban Gabriela, Gârban Z., pp. 267-270, in Macro and Trace Elements, Mengen- und Spurenelemente, 23. Workshop 2006, Friedrich-Schiller-Universität Jena (Hrsg. Anke M. et al.), Verlag Harald Schubert, Leipzig, 2006.
103. ***Peculiarities of metallograms in mixed urolithiasis*** - in collab. with Gârban Gabriela., pp.103-108, in Metal Elements in Environment, Medicine and Biology, Tome VII (Eds. Gârban Z., Drăgan P.), Publishing House Eurobit Timișoara, 2006.
104. ***Biogenesis of urolithiasis. Profile of uroconcrements metallogram in urolithiasis*** - in collab. with Popescu Sofia, Gârban Gabriela, Grecu Rodica, Gârban Z., pp.271-278, in Macro and Trace Elements, Mengen- und Spurenelemente, 23. Workshop 2006, Friedrich-Schiller Universität Jena (Hrsg. Anke M. et al.), Verlag Harald Schubert, Leipzig, 2006.